

Acoustic Voice Quality Index fordiscriminating Normal and different Vocal Pathological Conditions

by Author Author

Submission date: 28-Jul-2020 11:57AM (UTC+0530)

Submission ID: 1363147466

File name: REVISED_MANUSCRIPT.docx (716.76K)

Word count: 5618

Character count: 31230

Acoustic Voice Quality Index for discriminating Normal and different Vocal Pathological Conditions

Abstract

The Acoustic Voice Quality Index (AVQI) is a multiparametric measure to assess the overall quality of voice using both *sustained and continuous speech*. The study was aimed to compare the AVQI values across normal voice and dysphonic voice due to different pathological conditions, and to compare the values obtained from constituent parameters of AVQI across normal and different vocal pathological conditions. Seventy-four participants in the dysphonic group and 28 in the normal voice group were considered. Phonation samples of /a/ and reading samples were recorded using the *Praat program*, and AVQI was calculated using *Praat AVQI script v.2.03*. The perceptual analysis was done by three Speech-Language Pathologists using the GRBAS scale. The results revealed that among the dysphonic group, higher AVQI values were seen in unilateral vocal cord (VC) palsy, followed by bilateral mass lesion, unilateral mass lesion, MTD-I, MTD-II & III, and acute laryngitis. CPPs and HNR values were found to be least in the unilateral VC palsy group indicating high breathiness and noise component, respectively. Shimmer local and shimmer local dB values were high in the unilateral VC palsy group and mass lesion groups suggesting maximum aperiodic vibration of vocal folds in these groups. To conclude, AVQI and constituent parameters might help in discriminating vocal pathological conditions acoustically. As the present study is preliminary in nature, future studies can be carried out with greater sample size, restricted age range, and by considering the perceptual dysphonia severity.

Introduction

Mostly when an individual's voice quality, loudness, and pitch vary from those of people of similar age, gender, geographical location, and cultural background, it suggests voice disorder (Aronson, 1980; Boone, 1977). The Diagnostic Classification System of Voice Disorders (DCSVC) grossly divides the voice disorders into two groups, i.e., Organic voice disorder (OVD) and Functional voice disorder (FVD). Further, FVD consists of two groups, i.e., psychogenic voice disorders (PVD) and muscle tension voice disorder (MTVD) (Baker, Ben-Tovim, Butcher, Esterman, McLaughlin, 2007). The prevalence of voice disorders among communication disorders found to be around 4-7% in the Indian context (Sinha, Shivaswamy, Barman, Seth, Seshadri & Savithri, 2017; Konadath, Chatni, Lakshmi, & Saini, 2017). The prevalence rate of voice disorders among the professional voice users is quite high; 86% of the politicians, 74% vendors, 59% singers, and 49% of teachers found to have voice problems (Boominath, Rajendran, Nagarajan, Seethapathy & Gnanasekar, 2008). These studies suggest that voice disorder is a prevalent condition in the Indian context and hence needs attention regarding precise diagnosis and effective intervention.

The perceptual and acoustic analyses of voice are a vital part of the voice evaluation carried out by Speech-Language Pathologists as they provide excellent measures of intervention outcome (Stemple, Roy, & Klaben, 2014). Perceptual analysis of voice is a process of listening to recorded voice samples or live patient's voice and describing the abnormalities of a voice, specifically the deviations in terms of pitch, loudness, and quality. Acoustic analysis of voice provides quantitative data on vocal fold vibration in terms of pitch and amplitude, perturbation measures, harmonics to noise ratio, spectral, and cepstral measures, which in turn provide a better understanding of the pattern of vocal fold vibrations (Maryn, Roy, De Bodt, Van Cauwenberge, & Corthals, 2009).

Maryn, Corthals, Van Cauwenberge, Roy, De Bodt (2010) developed AVQI, a multiparametric acoustic model to assess voice quality, which uses both *sustained and continuous speech* to improve the ecological validity, perceptual, and instrumental assessment of dysphonia. For this purpose, sustained and continuous speech (reading phonetically balanced text) samples were collected from the 251 participants (229 with dysphonia and 22 without dysphonia) and were linked together. Then the samples were given to five experience voice clinicians for the perceptual rating of overall voice quality. The non-voiced segments

within the continuous speech were removed using a custom voicing detection algorithm, and concatenated samples were analyzed using 13 acoustic parameters based on spectral and cepstral analyses, amplitude perturbation, and fundamental frequency perturbation. The AVQI equation consists of six acoustic parameters (smoothened cepstral peak prominence, shimmer local, harmonics-to-noise ratio, shimmer local dB, general slope of the spectrum, the tilt of the regression line through the spectrum).

$$AVQI = [3.295 - (0.111 * CPPs) - (0.073 * HNR) - (0.213 * shimmer\ local) + (2.789 * shimmer\ local\ dB) - (0.032 * slope) + (0.077 * tilt)] * 2.571.$$

Heman-Ackah, Michael, Goding (2002), and Maryn, et al. (2010) report the diagnostic efficacy of combining both continuous speech and sustained vowel samples in the acoustic and perceptual assessment of dysphonic voice. Studies have also reported that AVQI has diagnostic accuracy, concurrent validity, and responds to changes following intervention (Heman-Ackah, et al., 2002; Maryn, et al., 2010; Maryn, De Bodt, Roy, 2010). AVQI has been validated across different languages: Dutch, Lithuanian, Japanese, Korean, German, Spanish, and Kannada and is found it to be reliable (Maryn, et al., 2010; Uloza, Petrauskas, Padervinskis, Ulozaitė, Barsties, Maryn, 2017; Hosokawa, Barsties, Iwahashi, Iwahashi, Kato, Iwaki, Sasai, Miyauchi, Matsushiro, Inohara, Ogawa, 2017; Kim, Barsties, Lee, 2019; Barsties, Lehnert, Janotte, 2020; Delgado, Leon, Jiménez, Izquierdo, Barsties, 2018; Benoy, 2017; Pebbili, Shabnam, Pushpavathi, Rashmi, Sankar, Nethra, Shreya, Shashish, 2019). AVQI is found to be useful in discriminating normophonic and dysphonic voices. Though, there is a scarcity of research on whether AVQI can be a useful tool to discriminate across the different vocal pathological conditions. Currently, laryngeal/ vocal imaging is considered as a standard tool for understanding vocal fold physiology, its' pathologies, and in the differential diagnosis. Visual examination using endoscopy or stroboscopy are commonly used instruments for vocal imaging.

Many vocal pathologies can be treated through voice therapy, and the prognosis can be assessed through acoustic evaluation and vocal imaging. However, for monitoring the prognosis of voice therapy, frequent evaluations are required. In this scenario, the feasibility of endoscopy/stroboscopy reduces because of high operational time and cost factors. Also, most of the time, younger children with voice problems do not cooperate for endoscopic procedures. Hence, to overcome the concerns mentioned above, a preliminary

attempt was made to investigate if AVQI and its components can help discriminate normal voice and different vocal pathological conditions. Acoustic measures provide information about different vocal aspects like reduced ease of mucosal vibration, phonatory gap, irregularity in vocal cord vibration, and vocal fatigue. Hence, it is hypothesized that AVQI and the constituent parameters can also help in differentiating various vocal pathological conditions. The present study was aimed to investigate AVQI and the constituent parameters for discriminating normal voice and different vocal pathological conditions. The specific objectives of the study were to compare the AVQI values across normal voice and dysphonic voice due to different pathological conditions and to compare values obtained from constituent parameters of AVQI across normal and different vocal pathological conditions.

Methods

Participants

There were 74 participants in the dysphonic group (51 males and 23 females, Age range=11 to 82 years, mean age= 39.4±15.5 years). The dysphonic group had Bilateral mass lesion=13; unilateral mass lesion=16; unilateral palsy=10; muscle tension dysphonia type II and III (MTD II & III) = 9; muscle tension dysphonia type I (MTD-I) =18; and Acute laryngitis=8. The bilateral and unilateral mass lesion group majorly consisted of individuals with vocal nodule and vocal polyp. There were 28 in normal voice group with 13 males and 15 females (Age range= 19 to 39 years; Mean age= 24.7±4.2 years).

All the participants considered for the study were native Kannada speakers. The individuals under the dysphonic group had to undergo the routine clinical examination, which involved detailed case history perceptual evaluation, acoustic & aerodynamic evaluation, and vocal imaging. The underlying vocal pathology was diagnosed by the team involving a Speech-Language Pathologist (SLP), and an Otolaryngologist using the videostroboscopy Xion Endostrob E with a 70-degree rigid scope and the Xenon R-180 LED light source for illumination. The individuals with normal hearing abilities, with dysphonia ranging from slight to severe, and with organic and functional voice disorders were included in the dysphonic group. For the normal voice group, the individuals had to undergo a perceptual examination of voice by an experienced SLP. The individuals with perceptually normal voice (G=0 on GRBAS scale; Hirano, 1981); individuals with no complaints of voice problems or upper respiratory tract infections,

asthma, or allergic disease on the day of recording and with normal hearing and cognitive abilities were considered for the normal group. The written consent was obtained from each participant, where information regarding the aim, objectives, method of the research, and approximate duration of the procedure was mentioned.

1 Procedure for voice recording


For the acoustic analysis, the voice recording was done in a sound-treated room where the average ambient noise level was 25 dB. The participants were made to sit comfortably, and the table-mounted dynamic microphone Shure SM48 (Shure Incorporated Product Support Niles, IL) was placed at a distance of 4 to 5 cm and 30° angle from the participant's mouth. All the recordings were done at a sampling frequency of 44.1 kHz, 16-bit resolution, in the mono channel, using the program *Praat* v. 6.0.40, and were saved in .wav format. For AVQI calculation, both phonation, as well as continuous speech sample, was required. For this purpose: the participants were instructed to phonate vowel /a/ for more than three seconds, and three trials were taken at their comfortable pitch and loudness. The most stable recording was considered for analysis. An interval of 2 minutes was given between each recording. Next, they were asked to read the first paragraph of the standardized Voiced Kannada passage (Shasidhar, 1984) at their comfortable pitch and loudness. As the AVQI requires sustained phonation of three seconds to be named as 'sv' and continuous speech sample to be named as 'cs,' the obtained samples opened in the *Praat* program and were truncated, renamed and were saved .wav format accordingly. The second sentence of the Voiced Kannada passage (/i: u:rannu namma ra:dzjaða bamba:I ennuvaru/) was considered for the continuous speech sample.

1 Acoustic Analysis of voice samples

For the calculation AVQI, both 'sv' and 'cs' were opened in the *Praat* program, the *Praat* script of AVQI version 02.03 (Maryn, 2013) was run in the *Praat* program, and then AVQI value and values of constituent parameters are obtained on output window (Fig 1).

Figure 1

An image of the AVQI 02.03 output



Perceptual analysis of voice samples

The perceptual analysis was done by three SLPs (raters). They had a minimum of 5 years of clinical experience in dealing with the diagnosis and management of voice disorder. The GRBAS scale was used for the perceptual analysis where the overall grade (severity of dysphonia) was rated on 0-3 scale (0, 1, 2, 3 representing normal, slight, moderate, and severe, respectively). Based on the consensus across at least two of the three raters, a particular grade was assigned to each sample. The voice samples were based on overall Grade of dysphonia (G): 28 samples under the normal, 40 under slight, 25 under moderate, and nine under

the severe category. Also, to confirm the consistency among the raters, inter-rater agreement⁴⁴ was assessed for each pair by using Cohen's Kappa.

²⁷ Statistical Analysis

The Shapiro -Wilks' test⁵ was carried out to test the assumption of normality for AVQI and its constituent parameter across all the normal and vocal pathological groups. Descriptive statistics were done to obtain the mean and standard deviation (SD) values for AVQI and its constituent parameters across all the groups. A¹⁹ one-way analysis of variance (ANOVA) with Tukey post hoc analysis was carried out to determine the group mean difference for AVQI. MANOVA and²⁴ Tukey post hoc test was carried out to investigate the main effect of type of pathology on the acoustic measures.⁶ Mann Whitney U test was used to observe the effect of type of pathology on shimmer local and shimmer local dB.

Results

¹ Inter-rater agreement

Cohen's Kappa coefficient was used to assess the inter-rater agreement for the overall grade (G) of dysphonia severity. The coefficient for Rater 1 vs. Rater 2 = 0.66; Rater 1 vs. Rater 3 = 0.64, and for Rater 2 vs. Rater 3 = 0.76, signifying good agreement among the raters.

⁶ Test of Normality

The results of the Shapiro-Wilks' test revealed that all the measures except for shimmer local and shimmer local dB followed a normal distribution (p -value > 0.05) across all the vocal pathological and normal groups. The measures shimmer local and shimmer local dB did not follow normal distribution for the normal and unilateral mass lesion groups (p -value < 0.05), while for rest of the groups followed a normal distribution (p -value > 0.05).⁹

Comparison of AVQI values across different groups:

The results of descriptive statistics suggest that higher AVQI values were obtained for the dysphonic group compared to normal. Among the dysphonic group, higher values were obtained for unilateral VC

palsy, followed by bilateral mass lesion, unilateral mass lesion, MTD-I, MTD-II & III, and laryngitis (Table 1). Higher the AVQI value indicates poorer the overall voice quality. The unilateral mass lesion, MTD-I, MTD-II & III, and laryngitis found to have similar AVQI values. The results of ANOVA showed that there is a significant effect of pathologies on AVQI values ($F(6, 95) = 11.77, P < 0.001$). The result of the Tukey post hoc test suggested that the normal group had significantly lower AVQI values compared to unilateral VC palsy, bilateral mass lesion, and unilateral mass lesion groups ($P < 0.05$). Moreover, unilateral VC palsy had significantly higher AVQI values compared to other pathological groups ($P < 0.05$).

Table 1

Mean (SD) AVQI values across different pathological conditions and Normal

Groups	Mean	Lowest	Highest
Unilateral VC palsy	5.62 (1.56)	2.81	8.26
Bilateral mass lesion	4.07 (1.41)	2.27	6.69
Unilateral mass lesion	3.60 (1.68)	1.49	6.50
MTD-I	3.48 (1.32)	1.06	5.72
MTD II & III co-existed	3.08 (1.14)	1.06	4.91
Laryngitis	3.06 (1.12)	1.25	4.60
Normal voice	1.94 (0.83)	0.26	3.50

Comparison of CPPs, HNR, Slope, and Tilt values across different groups:

The CPPs values were found to be least for unilateral VC palsy, and the value increases respectively in the bilateral mass lesion, unilateral mass lesion, MTD-I, acute laryngitis, MTD-II & III, and maximizes for the normal group (Table 2). Similarly, HNR values found to be least for unilateral VC palsy, followed by bilateral mass lesion, unilateral mass lesion, MTD-I. MTD-II & III, acute laryngitis, and maximum for normal group. CPPs and HNR values found to be least for the unilateral VC palsy group. The values obtained for slope and tilt did not vary much across the groups. The MANOVA showed an overall significant main effect of type of pathology on the acoustic measures, Wilks' Lambda = 0.38, $F(24, 322.16) = 4.20, P < 0.001$. The subsequent ANOVA result for each parameter has been summarized in Table 3.

Table 2

Mean (SD) of CPP, HNR, Slope values across different pathological conditions and Normal

Groups	CPPs	HNR	Slope	Tilt
Unilateral VC palsy	8.39 (3.35)	12.15 (5.73)	-23.24 (4.09)	-11.49 (2.34)
Bilateral mass lesion	10.74 (2.86)	18.35 (5.23)	-24.12 (7.02)	-11.44 (1.62)
Unilateral mass lesion	11.85 (2.73)	18.85 (6.03)	-25.35 (4.19)	-12.43 (0.72)
MTD-I	11.96 (2.89)	19.58 (4.17)	-25.79 (4.72)	-12.84 (0.64)
MTD II & III co-existed	12.90 (1.57)	21.95 (3.64)	-27.67 (6.10)	-11.45 (1.39)
Laryngitis	13.74 (1.04)	20.63 (2.82)	-24.01 (4.06)	-10.59 (2.51)
Normal	14.82 (1.71)	22.87 (3.38)	-23.42 (5.61)	-12.53 (1.25)

Table 3

ANOVA results for CPPs, HNR, Slope, and Tilt differentiating across the groups

Parameters	F (6, 95)	Sig.	Partial Eta Squared
CPPs	10.86	.000	.407
HNR	7.67	.000	.326
Slope	1.08	.375	.064
Tilt	3.85	.002	.196

² The results of the Tukey post hoc test indicated that CPPs values were significantly high for the normal group compared to unilateral VC palsy, unilateral mass lesion, and bilateral mass lesion ($P < 0.05$). The unilateral VC palsy group found to have significantly lesser CPPs values compared to other pathological groups ($P < 0.05$). The CPPs values obtained for MTD-I, MTD-II & III, acute laryngitis, and normal did not differ significantly. ⁷ HNR values found to be significantly less in the unilateral VC palsy group compare to normal and other pathological groups ($P < 0.05$). The values obtained for slope found to have no significant difference across the groups. Tilt values varied significantly across the groups ($P < 0.05$), but no definite pattern could be obtained.

¹⁰ **Comparison of Shimmer local and shimmer local dB values across different groups:**

The shimmer local and shimmer local dB values were high for the unilateral VC palsy group, followed by bilateral mass lesion, unilateral mass lesion, MTD-I, MTD-II & III co-existed, acute laryngitis and least for the normal group (Table 4). The MANOVA showed a significant main effect of type of pathology on shimmer local and shimmer local dB, Wilks' Lambda = 0.67, $F(8, 104) = 2.84, P = 0.007$. The subsequent ANOVA result for shimmer local and shimmer local dB is summarized in Table 5. The results of the Tukey post hoc test suggested that shimmer local found to be significantly high for unilateral VC palsy compared to acute laryngitis and MTD type II & III ($P < 0.05$). There was no significant difference across the bilateral mass lesion, MTD type I, MTD type II & III, and acute laryngitis groups for shimmer local. Shimmer local dB found to be significantly high for unilateral VC palsy compared to acute laryngitis, MTD-I, and MTD-II & III co-existed. At the same time, there was no significant difference between unilateral palsy and bilateral mass lesion groups.

Table 4

Mean (SD) and Median values for across Shimmer local and shimmer local dB different pathological conditions and Normal

Groups	Shimmer local		Shimmer local dB	
	Mean (SD)	Median	Mean (SD)	Median
Unilateral VC palsy	10.51 (8.03)	10.13	.98 (.52)	0.91
Bilateral mass lesion	6.94 (3.52)	6.27	.66 (.29)	0.55
Unilateral mass lesion	6.47 (4.69)	4.80	.66 (.43)	0.51
MTD Type I	5.87 (2.80)	5.32	.58 (.21)	0.53
MTD Type II & III	4.89 (2.07)	4.75	.48 (.15)	0.46
Laryngitis	4.39 (1.36)	4.07	.46 (.14)	0.44
Normal	3.22 (1.18)	3.13	.35 (.46)	0.32

Table 5

ANOVA results for shimmer local and shimmer local dB differentiating across the groups

Parameter	F (4, 53)	Sig.	Partial Eta Squared
shimmer local	3.328	.017	.201
shimmer local dB	4.863	.002	.268

The results of the Mann Whitney U test indicated that the unilateral mass lesion group has a significantly higher value compared to the normal group. At the same time, there was no significant difference observed between unilateral mass lesion when compared with other pathological conditions for both shimmer local and shimmer local dB. Also, the normal group found to have significantly lower shimmer local and shimmer local dB values compared to all pathological conditions ($*p < 0.05$).

Discussion

Comparison of AVQI values across different groups

The first objective of the present study was to compare the AVQI values across normal voice and dysphonic voices due to different pathological conditions. The results of the present study suggest that the normal group had significantly lower AVQI values compared to the dysphonic group. These results are in consensus with previous studies where they had obtained significantly lower AVQI values for normophonic group compared to the dysphonic group (Pebbili et al., 2019; Benoy, 2017).

The unilateral VC palsy group had significantly higher AVQI values compared to other pathological groups. This can be due to the manifestation of larger glottic chink and asynchronous vocal fold vibration compared to other pathological conditions. This result can be supported by literature, which reports that the majority of palsy cases have type 4 voice quality wherein, there is a wide phonatory gap, and voice quality is extremely breathy (Dedo, 1992). The unilateral mass lesion, MTD-I, MTD-II & III, and acute laryngitis found to have similar AVQI values. AVQI values in these groups were significantly lower than the unilateral VC palsy group and significantly higher than the normal group. The other pathological groups had lesser AVQI values in comparison to unilateral VC palsy, which can be attributed to a lesser extent of phonatory gap and irregularity in vocal fold adduction in them. For example, nodules and polyps will have increased mass and stiffness of the vocal folds, as well as hourglass-shaped glottic closure with reduced vibratory amplitude and mucosal wave (Hirano and Bless, 1993). Acute laryngitis is reported to have generalized edema, decreased or absent mucosal wave, and slightly decreased in vibratory amplitude (Sapienza, & Hoffman-Rudy, 2009). MTD will have an excessive glottic and supraglottic medial

contraction, anterior-posterior contraction of the supraglottic musculature, decreased vibratory amplitude, or psychogenic bowing of vocal folds (Altman, Atkinson, & Lazarus, 2005; Lee, & Son, 2009). While unilateral VC palsy is characterized with weakened or bowed vocal fold, presence of passive vibration around the paralyzed vocal fold, arytenoid cartilage on the affected side will not abduct or adduct; also there will be asymmetry characterized by slower initiation of the mucosal wave on the affected side along with a slower period and reduced amplitude of vibration (Sercarz, Berke, Gerratt, Ming, & Natividad, 1992). Hence, the extent of pathology seems to be more in unilateral VC palsy resulting in higher AVQI value in them compared to other pathological conditions.

Comparison of CPPs, HNR, shimmer local, shimmer local dB, Slope, and Tilt values across different groups

The next objective of the study was to compare values obtained from constituent parameters of AVQI across normal and different vocal pathological conditions. The CPPs values were significantly high for the normal group compared to unilateral VC palsy, unilateral mass lesion, and bilateral mass lesion. Literature reports high CPP value for normophonic individuals due to the presence of well-defined harmonic structure, and low in severe dysphonic voices as the harmonic formation is restricted by irregular adduction of vocal folds (Heman-Ackah et al., 2002). The unilateral VC palsy group found to have significantly lesser CPPs values compared to other pathological groups. Lesser CPPs values can be because CPPs have a high correlation with breathiness (Hillenbrand and Houde, 1996), and unilateral VC palsy will have a high breathiness component due to a large phonatory gap. The CPPs values obtained for MTD-I, MTD-II & III; acute laryngitis and normal did not differ significantly, which can be due to lesser severity of dysphonia and lesser extent of pathology in MTD and acute laryngitis compared to palsy and mass lesion conditions.

HNR values found to be significantly less in the unilateral VC palsy group compare to normal and other pathological groups due to the presence of high noise components in palsy conditions. Even this can be because of the presence of a wide phonatory gap in unilateral VC palsy. At the same time, other conditions would have better vocal fold closure as both the vocal cords would have mobility. The high noise component results from incomplete glottal closure that creates excess air during phonation, which increases the noise amplitude, in turn, lowers the HNR (Hartl, Hans, Vaissiere, Riquet, Brasnu, 2001; Oguz, Demirci, Safak, Arslan, Islam, Kargin, 2007).

⁹ Shimmer local and shimmer local dB found to be significantly high value for unilateral VC palsy compared to other groups suggesting maximum aperiodic vibration of vocal folds in them. Patel and Parsram (2005) had reported significantly higher shimmer values in individuals with vocal cord paralysis compare to normophonic individuals, which results from asynchronous vibration of vocal cords. The study reports higher shimmer values in the mass lesion group compared to normal. This can be attributed to the inflammation or ¹⁶ small masses on vocal folds leading to inconsistent glottal closure, and poorer ³³ vocal fold median edge contact (Oguz, Tarhan, Korkmaz, Yilmaz, Safak, Demirci, & Ozluoglu, 2007). The result of the study agrees with Davis (1979), where amplitude perturbation quotient (APQ) values were higher for unilateral paralysis followed by nodules and then laryngitis. Lieberman (1963) reports that inflammation and very small growth on vocal folds only minimally affects the perturbation measures, while larger masses can produce increased perturbation.

Spectral slope and spectral tilt are measures obtained from Long-term Average Spectrum (LTAS) analysis. The signal attained through LTAS represents the vocal function taking place at larynx as sound and transfer through the vocal tract (Lofqvist and Mandersson, 1987). The spectral slope has been identified as a correlate of hoarseness in the voice. The smaller values of the spectral slope values indicate a slower decline of energy with frequency, which is associated with vocal fold hyperfunction. In comparison, larger values of spectral slope indicate a faster decline of energy with frequency, which can be associated with vocal hypofunction (Ludlow, Kent, & Gray, 2018).

Similarly, spectral tilt was found to be associated with glottal closure during phonation. A reduction in spectral tilt value is associated with hyperadduction and high values associated with hypoadduction (Ludlow, Kent, & Gray, 2018). Although, in the current study, the values obtained for slope and tilt did not vary much across the groups, indicating that slope and tilt might not help discriminate the pathological conditions when considered in isolation.

Conclusion

AVQI and constituent parameters might help in discriminating pathological conditions acoustically. The results of the study have shown that CPPs, HNR, and Shimmer parameter values are well discriminated

across the pathological conditions. The values obtained for palsy, mass lesion, and muscle tension dysphonia are well demarcated. However, the current study is preliminary and hence future studies can consider following points such as (i) higher ⁷ and an equal number of participants in each group; (ii) restricting the age range, as age affect the acoustic measures (children, individuals in pubertal age, adults and geriatric population vary in their acoustic norms); (iii) overall perceptual dysphonia severity; and (iv) the size of mass lesions also might provide us with some remarkable and supporting results. Further studies in this regard can assist Speech-language pathologist in screening and diagnosis of voice disorders, and monitoring the prognosis during the voice therapy effectively. AVQI 02.03 is a non-commercial tool, that runs in the *Praat* program, making it cost-effective; also, it is less time consuming, and non-invasive.

References

- Altman, K. W., Atkinson, C., & Lazarus, C. (2005). Current and emerging concepts in muscle tension dysphonia: A 30-month review. *Journal of Voice*, *19*(2): 261-267.
- Aronson, A. E. (1980). *Clinical Voice Disorders. An Interdisciplinary Approach*. New York, NY: Thieme-Stratton, Inc.
- Baker, J., Ben-Tovim, D. I., Butcher, A., Esterman, A., & McLaughlin, K. (2007). Development of a modified diagnostic classification system for voice disorders with inter-rater reliability study. *Logopedics Phoniatrics Vocology*, *32*(3): 99-112.
- Benoy, J. J. Acoustic Voice Quality Index (AVQI) and perceptual measures in the Indian population. Unpublished Master's Dissertation submitted to University of Mysore, 2017.
- Boominathan, P., Rajendran, A., Nagarajan, R., Seethapathy, J., & Gnanasekar, M. (2008). Vocal abuse and vocal hygiene practices among different level professional voice users in India: A survey. *Asia Pacific Journal of Speech, Language and Hearing*, *11*(1): 47-53.
- Boone, D. R. (1977). *The Voice and Voice Therapy*. Englewood Cliffs, NJ: Prentice Hall.
- Davis, S.B. (1979). Acoustic characteristics of normal and pathological voices. *Speech and language*, *1*: 271-335.
- Dedo, H. H. (1992). Injection and removal of Teflon for unilateral vocal cord paralysis. *Annals of Otolaryngology, Rhinology & Laryngology*, *101*(1): 81-86.
- Delgado Hernandez, J., Leon Gomez, N. M., Jiménez, A., Izquierdo, L. M., & Barsties v. Latoszek, B. (2018). Validation of the acoustic voice quality index version 03.01 and the acoustic breathiness index in the Spanish language. *Annals of Otolaryngology, Rhinology & Laryngology*, *127*(5): 317-326.
- Hartl, D. M., Hans, S., Vaissière, J., Riquet, M., & Brasnu, D. F. (2001). Objective voice quality analysis before and after onset of unilateral vocal fold paralysis. *Journal of Voice*, *15*(3): 351-361.
- Heman-Ackah, Y. D., Michael, D. D., & Goding Jr, G. S. (2002). The relationship between cepstral peak prominence and selected parameters of dysphonia. *Journal of Voice*, *16*(1): 20-27.
- Hillenbrand, J., Cleveland, R. A., & Erickson, R. L. (1994). Acoustic correlates of breathy vocal quality. *Journal of Speech, Language, and Hearing Research*, *37*(4): 769-778.

- Hirano, M. (1981). "GRBAS" scale for evaluating the hoarse voice & frequency range of phonation. In M. Hirano (Ed.), *Clinical Examination of Voice*, (pp. 83-84). New York, NY: Springer-Verlag/Wein.
- Hirano, M., & Bless, D. M. (1993). *Videostroboscopic Examination of the Larynx*. San Diego, CA: Singular.
- Hosokawa, K., Barsties, B., Iwahashi, T., Iwahashi, M., Kato, C., Iwaki, S., ... & Ogawa, M. (2017). Validation of the acoustic voice quality index in the Japanese language. *Journal of Voice*, 31(2): 260-e1.
- Kim, G. H., von Latoszek, B. B., & Lee, Y. W. (2019). Validation of acoustic voice quality index version 3.01 and Acoustic Breathiness Index in Korean Population. *Journal of Voice*.
<https://doi.org/10.1016/j.jvoice.2019.10.005>.
- Konadath, S., Chatni, S., Lakshmi, M. S., & Saini, J. K. (2017). Prevalence of communication disorders in a group of islands in India. *Clinical Epidemiology and Global Health*, 5(2): 79-86.
- Latoszek, B. B. V., Lehnert, B., & Janotte, B. (2020). Validation of the acoustic voice quality index version 03.01 and acoustic breathiness index in German. *Journal of Voice*, 34(1): 157-e17.
- Lee, E. K., & Son, Y. I. (2005). Muscle tension dysphonia in children: Voice characteristics and outcome of voice therapy. *International Journal of Pediatric Otorhinolaryngology*, 69(7): 911-917.
- Lieberman, P. (1963). Some acoustic measures of the fundamental periodicity of normal and pathologic larynges. *The Journal of the Acoustical Society of America*, 35(3): 344-353.
- Löfqvist, A., & Mandersson, B. (1987). Long-time average spectrum of speech and voice analysis. *Folia Phoniatica et Logopaedica*, 39(5): 221-229.
- Ludlow, C. L., Kent, R. D., & Gray, L. C. (2018). *Measuring Voice, Speech, and Swallowing in the Clinic and Laboratory*. San Diego, CA: Plural Publishing.
- Maryn, Y., Corthals, P., Van Cauwenberge, P., Roy, N., & De Bodt, M. (2010). Toward improved ecological validity in the acoustic measurement of overall voice quality: Combining continuous speech and sustained vowels. *Journal of Voice*, 24(5): 540-555.
- Maryn, Y., De Bodt, M., & Roy, N. (2010). The acoustic voice quality index: Toward improved treatment outcomes assessment in voice disorders. *Journal of Communication Disorders*, 43(3): 161-174.
- Maryn, Y. (2013). *De Acoustic Voice Quality Index in het programma Praat: een praktische handleiding*. Belsele: Vlaamse Vereniging voor Logopedisten
- Maryn, Y., Roy, N., De Bodt, M., Van Cauwenberge, P., & Corthals, P. (2009). Acoustic measurement of overall voice quality: A meta-analysis. *The Journal of the Acoustical Society of America*, 126(5): 2619-2634.
- Oguz, H., Demirci, M., Safak, M. A., Arslan, N., Islam, A., & Kargin, S. (2007). Effects of unilateral vocal cord paralysis on objective voice measures obtained by Praat. *European Archives of Oto-Rhino-Laryngology*, 264(3): 257-261.
- Oguz, H., Tarhan, E., Korkmaz, M., Yilmaz, U., Safak, M. A., Demirci, M., & Ozluoglu, L. N. (2007). Acoustic analysis findings in objective laryngopharyngeal reflux patients. *Journal of Voice*, 21(2): 203-210.
- Patel, R., & Parsram, K. S. (2005). Acoustic analysis of subjects with vocal cord paralysis. *Indian Journal of Otolaryngology and Head and Neck Surgery*, 57(1): 48-51.
- Pebbili, G. K., Shabnam, S., Pushpavathi, M., Rashmi, J., Sankar, R. G., Nethra, R., ... & Shashish, G. (2019). Diagnostic Accuracy of Acoustic Voice Quality Index Version 02.03 in Discriminating across the Perceptual Degrees of Dysphonia Severity in Kannada Language. *Journal of Voice*.
<https://doi.org/10.1016/j.jvoice.2019.07.010>

- Sapienza, C., & Hoffman-Rudy, B. (2009). *Voice Disorders*. San Diego, CA: Plural Publishing.
- Sercarz, J. A., Berke, G. S., Gerratt, B. R., Ming, Y., & Natividad, M. (1992). Videostroboscopy of human vocal fold paralysis. *Annals of Otology, Rhinology & Laryngology*, 101(7): 567-577.
- Shasidhar, K.N. "Voiced Kannada Passage" developed as a part of unpublished Master's Dissertation titled "Analysis of Speech of Stutterers" submitted to University of Mysore, 1984.
- Sinha, S. K., Shivaswamy, J., Barman, A., Seth, D., Seshadri, D., & Savithri, S. R. (2017). Prevalence of communication disorders in a rural population at taluq level of Gujarat, India. *Clinical Epidemiology and Global Health*, 5(2): 73-78.
- Stemple, J. C., Roy, N., & Klaben, B. K. (2018). *Clinical Voice Pathology: Theory and Management*. San Diego, CA: Plural Publishing.
- Uloza, V., Petrauskas, T., Padervinskis, E., Ulozaitė, N., Barsties, B., & Maryn, Y. (2017). Validation of the acoustic voice quality index in the Lithuanian language. *Journal of Voice*, 31(2): 257-e1.

Table 1

Mean (SD) AVQI values across different pathological conditions and Normal

Groups	Mean	Lowest	Highest
Unilateral VC palsy	5.62 (1.56)	2.81	8.26
Bilateral mass lesion	4.07 (1.41)	2.27	6.69
Unilateral mass lesion	3.60 (1.68)	1.49	6.50
MTD-I	3.48 (1.32)	1.06	5.72
MTD II & III co-existed	3.08 (1.14)	1.06	4.91
Laryngitis	3.06 (1.12)	1.25	4.60
Normal voice	1.94 (0.83)	0.26	3.50

Table 2

Mean (SD) of CPP, HNR, Slope values across different pathological conditions and Normal

Groups	CPPs	HNR	Slope	Tilt
Unilateral palsy	8.39 (3.35)	12.15 (5.73)	-23.24 (4.09)	-11.49 (2.34)
Bilateral mass lesion	10.74 (2.86)	18.35 (5.23)	-24.12 (7.02)	-11.44 (1.62)
Unilateral mass lesion	11.85 (2.73)	18.85 (6.03)	-25.35 (4.19)	-12.43 (0.72)
MTD-I	11.96 (2.89)	19.58 (4.17)	-25.79 (4.72)	-12.84 (0.64)
MTD II & III co-existed	12.90 (1.57)	21.95 (3.64)	-27.67 (6.10)	-11.45 (1.39)
Laryngitis	13.74 (1.04)	20.63 (2.82)	-24.01 (4.06)	-10.59 (2.51)
Normal	14.82 (1.71)	22.87 (3.38)	-23.42 (5.61)	-12.53 (1.25)

Table 3

MANOVA results for CPPs, HNR, Slope and Tilt differentiating across the groups

Parameters	F (6, 95)	Sig.	Partial Eta Squared
CPPs	10.86	.000	.407
HNR	7.67	.000	.326
Slope	1.08	.375	.064
Tilt	3.85	.002	.196

Table 4

Mean (SD) and Median values for across ⁴ Shimmer local and shimmer local dB different pathological conditions and Normal

Groups	Shimmer local		Shimmer local dB	
	Mean (SD)	Median	Mean (SD)	Median
Unilateral palsy	10.51 (8.03)	10.13	.98 (.52)	0.91
Bilateral mass lesion	6.94 (3.52)	6.27	.66 (.29)	0.55
Unilateral mass lesion	6.47 (4.69)	4.80	.66 (.43)	0.51
MTD Type I	5.87 (2.80)	5.32	.58 (.21)	0.53
MTD Type II & III	4.89 (2.07)	4.75	.48 (.15)	0.46
Laryngitis	4.39 (1.36)	4.07	.46 (.14)	0.44
Normal	3.22 (1.18)	3.13	.35 (.46)	0.32

Table 5

MANOVA results for ¹⁰ shimmer local and shimmer local dB differentiating across the groups

Parameter	F (4, 53)	Sig.	Partial Eta Squared
shimmer local	3.328	.017	.201
shimmer local dB	4.863	.002	.268

FIGURE LEGENDS

Figure 1

An image of the AVQI 02.03 output

Acoustic Voice Quality Index for discriminating Normal and different Vocal Pathological Conditions

ORIGINALITY REPORT

21%

SIMILARITY INDEX

10%

INTERNET SOURCES

15%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Gopi Kishore Pebbili, Srushti Shabnam, Mariswamy Pushpavathi, Jayaramu Rashmi et al. "Diagnostic Accuracy of Acoustic Voice Quality Index Version 02.03 in Discriminating across the Perceptual Degrees of Dysphonia Severity in Kannada Language", Journal of Voice, 2019
Publication 2%
- 2** fliphtml5.com
Internet Source 2%
- 3** Submitted to Loma Linda University
Student Paper 1%
- 4** biblio.ugent.be
Internet Source 1%
- 5** Submitted to All India Institute of Speech & Hearing
Student Paper 1%
- 6** www.aiishmysore.in
Internet Source 1%

-
- 7 Submitted to Curtin University of Technology 1%
Student Paper
-
- 8 Pebbili, Gopi Kishore, Juhi Kidwai, and Srushti Shabnam. "Dysphonia Severity Index in Typically Developing Indian Children", Journal of Voice, 2016. 1%
Publication
-
- 9 YeonWoo Lee, GeunHyo Kim, KeonTae Sohn, ByungJoo Lee, JinChoon Lee, SoonBok Kwon. "The Usefulness of Auditory Perceptual Assessment and Acoustic Analysis as a Screening Test for Voice Problems", Folia Phoniatica et Logopaedica, 2019 1%
Publication
-
- 10 Ben Barsties, Rudi Verfaillie, Nelson Roy, Youri Maryn. "Do body mass index and fat volume influence vocal quality, phonatory range, and aerodynamics in females?", CoDAS, 2013 1%
Publication
-
- 11 pubs.asha.org 1%
Internet Source
-
- 12 Kwang Jae Cho, Inn Chul Nam, Yeon Shin Hwang, Mi Ran Shim, Jun Ook Park, Jung Hae Cho, Young Hoon Joo, Min Sik Kim, Dong Il Sun. "Analysis of factors influencing voice quality and therapeutic approaches in vocal <1%

polyp patients", European Archives of Oto-Rhino-Laryngology, 2011

Publication

-
- | | | |
|----|---|-----|
| 13 | smd.it
Internet Source | <1% |
|----|---|-----|
-
- | | | |
|----|--|-----|
| 14 | Submitted to Higher Education Commission Pakistan
Student Paper | <1% |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 15 | hdl.handle.net
Internet Source | <1% |
|----|---|-----|
-
- | | | |
|----|---|-----|
| 16 | eprints.iums.ac.ir
Internet Source | <1% |
|----|---|-----|
-
- | | | |
|----|--|-----|
| 17 | Hee-Jeong Shim, Hun Jung, Rajinder Koul, Do-Heung Ko. "Spectral and Cepstral Based Acoustic Features of Voices with Muscle Tension Dysphonia", Clinical Archives of Communication Disorders, 2016
Publication | <1% |
|----|--|-----|
-
- | | | |
|----|--|-----|
| 18 | Wayland, R.. "Acoustic correlates of breathy and clear vowels: the case of Khmer", Journal of Phonetics, 200304
Publication | <1% |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 19 | www.hindawi.com
Internet Source | <1% |
|----|---|-----|
-
- | | | |
|----|---|-----|
| 20 | Saime Sagiroglu, Neslihan Kurtul. "The Effect of Supraclavicular Radiotherapy on Acoustic Voice | <1% |
|----|---|-----|

Quality Index (AVQI), Spectral Amplitude and Perturbation Values", Journal of Voice, 2019

Publication

21

Marina Englert, Ben Barsties v. Latoszek, Youri Maryn, Mara Behlau. "Validation of the Acoustic Voice Quality Index, Version 03.01, to the Brazilian Portuguese Language", Journal of Voice, 2019

Publication

<1%

22

Submitted to University of Ulster

Student Paper

<1%

23

www.karger.com

Internet Source

<1%

24

Submitted to Pennsylvania State System of Higher Education

Student Paper

<1%

25

Sarah McAlister, Irena Yanushevskaya. "Voice assessment practices of speech and language therapists in Ireland", Clinical Linguistics & Phonetics, 2019

Publication

<1%

26

Geun-Hyo Kim, Ben Barsties von Latoszek, Yeon-Woo Lee. "Validation of Acoustic Voice Quality Index Version 3.01 and Acoustic Breathiness Index in Korean Population", Journal of Voice, 2019

Publication

<1%

27	Submitted to BPP College of Professional Studies Limited Student Paper	<1%
28	Staffan Morén, Per Åke Lindestad, Mats Holmström, Maria Mani. "Voice Quality in Adults Treated for Unilateral Cleft Lip and Palate", The Cleft Palate-Craniofacial Journal, 2018 Publication	<1%
29	www.jneurosci.org Internet Source	<1%
30	Submitted to Queen Margaret University College, Edinburgh Student Paper	<1%
31	www.ncbi.nlm.nih.gov Internet Source	<1%
32	Submitted to Texas A & M University, Kingville Student Paper	<1%
33	Submitted to West Texas A&M University Student Paper	<1%
34	Submitted to University of Mary Student Paper	<1%
35	Submitted to University of Sheffield Student Paper	<1%
36	Submitted to National University of Ireland, Galway	<1%

37 Youri Maryn, Paul Corthals, Paul Van Cauwenberge, Nelson Roy, Marc De Bodt. "Toward Improved Ecological Validity in the Acoustic Measurement of Overall Voice Quality: Combining Continuous Speech and Sustained Vowels", *Journal of Voice*, 2010
Publication <1%

38 stimmklinik.de
Internet Source <1%

39 Sujitha P S, Gopi Kishore Pebbili. "Cepstral Analysis of Voice in Young Adults", *Journal of Voice*, 2020
Publication <1%

40 Submitted to University of Utah
Student Paper <1%

41 Leila Susana Finger, Carla Aparecida Cielo, Karine Schwarz. "Acoustic vocal measures in women without voice complaints and with normal larynxes", *Brazilian Journal of Otorhinolaryngology*, 2009
Publication <1%

42 www.scopus.com
Internet Source <1%

43 citeseerx.ist.psu.edu
Internet Source <1%

-
- 44 Barsties, Ben, and Youri Maryn. "The improvement of internal consistency of the Acoustic Voice Quality Index", American Journal of Otolaryngology, 2015. <1%
- Publication
-
- 45 Haldun Oguz, Munir Demirci, Mustafa A. Safak, Necmi Arslan, Ahmet Islam, Selda Kargin. "Effects of unilateral vocal cord paralysis on objective voice measures obtained by Praat", European Archives of Oto-Rhino-Laryngology, 2006 <1%
- Publication
-
- 46 Christopher R. Watts, Shaheen N. Awan, Youri Maryn. "A Comparison of Cepstral Peak Prominence Measures From Two Acoustic Analysis Programs", Journal of Voice, 2017 <1%
- Publication
-
- 47 Submitted to University of Auckland <1%
- Student Paper
-
- 48 Submitted to Australian Catholic University <1%
- Student Paper
-
- 49 Pasquale Bottalico, Juliana Codino, Lady Catherine Cantor-Cutiva, Katherine Marks et al. "Reproducibility of Voice Parameters: The Effect of Room Acoustics and Microphones", Journal of Voice, 2020 <1%
- Publication
-

50

Hirohito Umeno, Masamitsu Hyodo, Tomoyuki Haji, Hirotaka Hara et al. "A summary of the Clinical Practice Guideline for the Diagnosis and Management of Voice Disorders, 2018 in Japan", *Auris Nasus Larynx*, 2020

Publication

<1%

51

Janet Baker, David I. Ben-Tovim, Andrew Butcher, Adrian Esterman, Kristin McLaughlin. "Development of a modified diagnostic classification system for voice disorders with inter-rater reliability study", *Logopedics Phoniatrics Vocology*, 2009

Publication

<1%

52

Submitted to The Hong Kong Polytechnic University

Student Paper

<1%

53

Geun-Hyo Kim, Yeon-Yoo Lee, In-Ho Bae, Hee-June Park, Soon-Bok Kwon. "Application of the New Version of the Acoustic Voice Quality Index with Korean Speakers", *Communication Sciences & Disorders*, 2018

Publication

<1%

54

T. Jayakumar, Jesnu Jose Benoy, H. Mohamed Yasin. "Effect of Age and Gender on Acoustic Voice Quality Index Across Lifespan: A Cross-sectional Study in Indian Population", *Journal of Voice*, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On