



Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/colli



Nurses and midwives perceptions of missed nursing care – A South Australian study



Claire Verrall, Dip App Science (Nursing), RN, BN, MN^{a,*},
Elizabeth Abery, BHSc (Honours)^b, Clare Harvey, PhD, RN, MA^c,
Julie Henderson, PhD, BA (Hons), Grad Dip (Information
Studies)^a, Eileen Willis, PhD, BEd, MEd (Sociology)^b,
Patti Hamilton, RN, PhD, MS (Nursing), BS (Nursing)^d,
Luisa Toffoli, PhD, RN, MN^e, Ian Blackman, EdD, RN, RPN, Grad
Dip (Health Couns)^a

^a School of Nursing and Midwifery, Faculty of Medicine, Nursing and Health Sciences, Flinders University, Adelaide, South Australia, Australia

^b School of Health Sciences, Faculty of Medicine, Nursing and Health Sciences, Flinders University, Adelaide, South Australia, Australia

^c Eastern Institute of Technology, Napier, Hawke's Bay, New Zealand

^d Texas Woman's University, Denton, TX, United States

^e School of Nursing, University of South Australia, Adelaide, South Australia, Australia

Received 8 November 2013; received in revised form 14 July 2014; accepted 4 September 2014

KEYWORDS

Nursing and
midwifery;
Missed care;
Rationed care;
Staffing;
Skill mix

Summary

Background: Budgetary restrictions and shorter hospital admission times have increased demands upon nursing time leading to nurses missing or rationing care. Previous research studies involving perceptions of missed care have predominantly occurred outside of Australia. This paper reports findings from the first South Australian study to explore missed nursing care.

Aim: To determine and explore nurses' perceptions of reasons for missed care within the South Australian context and across a variety of healthcare settings.

Method: The survey was a collaborative venture between the Flinders University of South Australia, After Hours Nurse Staffing Work Intensity and Quality of Care project team and the Australian Nursing and Midwifery Federation, SA Branch.

Electronic invitations using *Survey Monkey* were sent to randomly selected nurses and midwives and available online for two months. Three hundred and fifty-four nurses and midwives responded. This paper reports qualitative data from answers to the open questions.

* Corresponding author at: School of Nursing and Midwifery, Faculty of Medicine, Nursing and Health Sciences, Flinders University of South Australia, GPO Box 2100, Adelaide, South Australia 5001, Australia. Tel.: +61 8 8201 3869; fax: +61 8 8276 1602.

E-mail addresses: verrall@flinders.edu.au, claire.verrall@flinders.edu.au (C. Verrall), elizabeth.abery@flinders.edu.au (E. Abery).

Findings: Three main reasons for missed care were determined as: competing demands that reduce time for patient care; ineffective methods for determining staffing levels; and skill mix including inadequate staff numbers. These broad issues represented participants' perceptions of missed care.

Conclusion: Issues around staffing levels, skill mix and the ability to predict workload play a major role in the delivery of care. This study identified the increasing work demands on nurses/midwives. Solutions to the rationing of care need further exploration.

© 2014 Australian College of Nursing Ltd. Published by Elsevier Ltd.

Introduction

Australia as with many western countries has seen an increase in the acuity of patients admitted to hospital and this, compounded by shortened lengths of stay intensifies nurses' workload which in turn has a significant impact on how they manage their time and prioritise patient care (Willis, 2009). Current issues within the Australian health-care system such as the size, composition and age of the nursing workforce provide the Australian nurse with a variety of challenges (Preston, 2009). Australia faces an ageing nursing workforce alongside of increased demand for nursing services arising from an ageing population (O'Brien-Pallas, Duffield, & Alknis, 2004). Staff shortages at the macro-level have been associated with calls for greater flexibility in staffing health services (Productivity Commission, 2005). Jacob, McKenna, and D'Amore (2013) argue for a manipulation of skill mix as a means of addressing staffing issues but also as a response to budgetary restraints contributing to increased employment of enrolled nurses and unregulated health professionals, e.g. nurse assistants to provide nursing care. Restructuring and budgetary constraints, irregular staffing levels and skill mix, set the scene for potential missed nursing care (Henderson et al., 2013).

Background

The notion of missed nursing care was first explored by Beatrice Kalisch in 2006. Kalisch and colleagues refer to missed care as "any aspect of required patient care that is omitted (either in part or in whole) or delayed" and acknowledged that it is a response to "multiple demands and inadequate resources" (Kalisch, Landstrom, & Hinshaw, 2009, p. 1509). Missed nursing care has been linked to negative patient outcomes (Schubert, Clarke, Glass, Schaffert-Witvliet, & De Geest, 2009) and attributed to a variety of causes from the work environment, to patient care demands and staffing issues (Aitkin, Clarke, & Sloane, 2002; Kalisch, Doumit, Lee, & Zein, 2013; Kalisch et al., 2009; Needleman, Buerhaus, Mattke, Stewart, & Zelevinsky, 2002; Papastavrou, Panayioti, & Georgios, 2013). Current research suggests that when a nurse's work load increases, there is less time to care for individual patients (Schubert, Glass, Clarke, Schaffert-Witvliet, & De Geest, 2008).

Kalisch in a qualitative study in 2006 identified a range of core nursing tasks that were routinely omitted. These tasks included discharge planning and patient education, emotional support, hygiene and mouth care, documentation of fluid intake and output, ambulation, feeding and general nursing surveillance of the patient. The nurses in her study

identified inadequate staffing levels and skill mix, unexpected workload increases, too few or lack of resources, poor handover and inadequate teamwork and orientation to the ward as key determinants of missed care. Her study led to the development of the MISSCARE survey instrument to formally measure common elements of missed care and the rationales behind them (Kalisch & Williams, 2009). Subsequently, Kalisch and her colleagues associated missed care with three primary antecedents in relation to patient care: (1) the availability of labour resources; (2) access to the material resources, and (3) relationship and communication factors (Kalisch et al., 2009; Kalisch & Williams, 2009). In more recent years, Kalisch has explored more specific aspects of the nursing work environment and its impact on missed care. Findings from these studies have identified a range of factors that contribute to and impact upon missed care. In 2009, Kalisch et al. examined the impact of nursing teamwork on missed care, arguing that it was not simply the number of nurses rostered, but the skill mix of nursing staff that impacted on perceptions of whether care was missed. The study also found that in line with their roles and responsibilities, Registered Nurses (RNs) were more likely than nurse assistants to report missed care and to associate this with an unexpected rise in patient volume or acuity, rates of admission and discharge and access to material resources. A later qualitative study by Kalisch, Gosselin, and Choi (2012) compared perceived differences in work environments between units with high and low levels of identified missed care. This study found that units reporting low levels of missed care had adequate and flexible staffing; effective communication and leadership; strong team focus; and shared accountability for monitoring and assessing work (Kalisch et al., 2012).

While Kalisch's work has been instrumental in developing and refining the concept of missed nursing care further studies have been undertaken in other contexts. Papastavrou et al. (2013) conducted a systematic literature review exploring rationales for missed care that support Kalisch's findings. Evidence collated from this review highlights a growing interest in missed care and attests to the global quest to improve patient quality and safety. When there are insufficient resources, nurses are forced to ration or omit care. It is this that impacts on negative patient outcomes and is a major challenge to quality assurance, risk management, nurse satisfaction and ultimately patient care (Papastavrou et al., 2013). Although much of the research in this area has been conducted outside of Australia, an Australian study conducted by Chaboyer et al. (2008), found that when nursing workload intensified the nursing roles could be "blurred" between acknowledged levels of skill.

These skill sets were identified as nursing procedures such as patient assessment, hygiene, medication administration and other nursing procedures.

The literature is consistent in its findings that missed nursing care does occur and the implications for staff, the patient and institutional management cannot be ignored. The challenges of providing cost effective, quality patient care are real and with an ageing population and associated increases in patient acuity and shortened length of stay are likely to magnify. This study sought to determine and explore nurses' perceptions of reasons for missed care within the South Australian context and across a variety of healthcare settings. This paper presents the qualitative findings from the survey of perceptions of South Australian community and acute care nurses and midwives of the factors which impact their capacity to perform nursing care.

Method

A modified version of Kalisch's MISSCARE survey (Kalisch & Williams, 2009) was used with the author's permission. Modifications to the survey were made to reflect the South Australian context as shift times, and terminology such as categories of employment and categories for education levels varied to that of the USA. Ethics approval was obtained through Flinders University Social and Behavioural Research Ethics Committee.

The survey was disseminated using Survey Monkey with the support of the Australian Nursing and Midwifery Federation (SA Branch) (ANMFSA) and available online for ANMFSA members for two months from November 1st to December 31st 2012. The survey contained demographic questions, questions that explored working conditions, questions concerning missed nursing care (defined as care omitted, postponed or incomplete) and questions concerning perceived reasons for missed or omitted care in the settings that the nurses/midwives practiced. Likert scales arising from the MISSCARE tool were used to estimate data. In addition, open-ended questions offered participants the opportunity to add personal comments concerning nursing care that is missed and their perceptions of the rationale behind this missed care. In keeping with qualitative research methods the inclusion of personal comments provided the opportunity for meaningful data through personal reflection. Further, the use of the qualitative comments is interpretive enabling an investigation of what meaning nurses give to events (Denzin & Lincoln, 2008) in the context of their own experiences and environment (Meyrick, 2006).

Recruitment and sample

Initially the survey was sent via email to 10% (1600) of the ANMFSA membership selected randomly. This sample was later expanded to the wider membership through an advertisement in the ANMFSA emailed newsletter that provided details of the study and a link to the online survey tool. Eligibility to participate required that members be a nurse or midwife currently working in a clinical setting at least once per fortnight.

Data analysis

Thematic analysis was used to interpret the results. This allowed the research team to identify themes within the data, but also to allow for the use of predictive and familiar themes known to the researchers (Pope, Ziebland, & Mays, 2006). These themes were predicted using literature to support their approach when mapping the social process for the study. Included in this approach was the information developed earlier by the researchers for the study by suggesting a blueprint for combining bibliometrics and critical analysis as a way to review scientific work in nursing (Hamilton et al., 2014). In this way both consumers and producers of nursing knowledge can recognise and take account of the social processes involved in development, evaluation and utilisation of new knowledge. Using this model, such key themes familiar to the researchers as nurses were chosen as a starting point. These included delayed care, rostered shifts and staff to patient ratios, patient acuity team collaboration and handover. Using this foundational theme collection the researchers (CH, EA, CH, JH, EW, PH, IB) initially reviewed responses individually to ascertain recurrent concepts, words and sentences from the qualitative data, which identified the initial coding method (Patton, 2002). The researchers then met to discuss, compare and further review themes until agreement was met. The established codes were then developed into categories or secondary codes (Patton, 2002). Three researchers (CV, EA & CH) intensively revised each category to support the validity of the finalised themes.

Findings

As it is difficult to ascertain how many members accessed the email or newsletter a true percentage of invited participants cannot be determined, however, 354 members participated in the survey. Not all participants completed every question with 258 completed in its entirety. The participants ranged in age, gender, years of experience, hours worked, and location and setting, providing a solid cross section of the South Australian nursing and midwifery workforce. Participant profiles of those who responded to the survey were indicative of ANMFSA member profiles. Table 1 provides the demographic profile of participants.

In total, 843 qualitative responses were included in the survey. Qualitative findings of this study indicate that missed care relates to systemic issues in three main areas: competing demands that reduce time for patient care; ineffective methods for determining staffing levels; and skill mix including inadequate staff numbers. The latter finding was particularly prominent within aged care settings. An analysis of the qualitative survey data revealed overall concerns of changing acuity of patients, the ageing population, changes in how nurses provide care and changes in the way nursing care is managed.

Table 2 shows the questions posed that elicited qualitative comment.

Further exploration of the main issues identified from the open-ended survey responses follow.

Table 1 Demographic profile of participants. As some participants did not complete all questions these numbers do not equate to total survey responses.

	<i>n</i> (%)
Gender	
Female	261 (90%)
Male	28 (10%)
Age	
Under 25	6 (2%)
25–34	34 (12%)
35–44	57 (20%)
45–54	108 (37%)
55–64	80 (27%)
65 and above	6 (2%)
Years of experience	
<2 years	38 (13%)
2–5 years	42 (14%)
5–10 years	45 (16%)
More than 10 years	166 (57%)
Number of hours worked	
<30 h per week	95 (33%)
30 h or more per week	195 (67%)
Location	
eMetropolitan	197 (68%)
Rural	93 (32%)
Setting	
Public	218 (75%)
Private	54 (19%)
Agency	18 (6%)

Competing demands

A majority of participants identified competing demands as significantly contributing to missed care. These demands were represented as disruptions to daily routines, alterations in patient acuity during a shift, unplanned admissions and unavailable resources such as certain patient medications and diagnostic equipment. This issue is summarised well by the following comment:

...nursing staff are frustrated by their inability to be everywhere and do everything. Most go home worrying that they haven't done everything they were meant to do, ... the nurses who care the most are stressed by their inability to be the nurses they want to be due to lack of time. We are all just doing the best we can with what we have (Participant 27).

A recurring factor was the lack of properly maintained equipment and resources needed to perform duties, with reference made to budgeting cuts overriding needs:

...a lot of time looking for things or chasing things up, i.e. medication running out, trying to find equipment that is available and working... (Participant 19).

...it seems equipment has been cut but patient acuity has increased (Participant 99).

Table 2 Provides details of the questions allowing qualitative comment and the number of responses elicited for each question (from most prevalent to least).

Questions concerning incidence of missed care	No. of responses
Turning patient every 2 h	95
Ambulation three times a day as ordered	67
Feeding patients while food is still warm	42
Setting up meals for patients who feed themselves	34
Medications administered within 30 min before or after scheduled time	33
Patient bathing/skin care	27
PRN medication requests acted on within 15 min	25
Vital signs assessed as ordered	24
Mouth care	24
Patient discharge planning & education	24
Emotional support to patient and/or family	22
Response to call bell/light initiated within 5 min	21
Attend interdisciplinary care conferences whenever held	21
Assist with toileting needs within 5 min of request	21
Hand washing	21
Monitoring intake/output	21
Full documentation of all necessary data	20
Patient education about illness, tests and diagnostic studies	20
Skin/wound care	17
Bedside glucose monitoring as ordered	16
Focused reassessments according to patient condition	16
Assess effectiveness of medications	16
Patient assessments performed each shift	14
IV/Central line care sive & assessment according to hospital policy	14
<i>Questions concerning rationale for missed care</i>	
Indicate the reasons which contributed to MISSED care in your ward/unit	14
<i>Addition questions</i>	
Is there anything else you would like to tell us about missed care?	107
What suggestions do you have about improving our survey?	69
Total	843
Total number of participants	<i>n</i> = 289

...there were no towels to attend to patient washes/showers as well as no incontinent pads or basic dressing material... (Participant 7).

Frequent additional or unexpected tasks such as phone calls, visitors' requests, stat (immediate) drug administration, as well as impromptu discussions with allied health

about patient care, impacted on the nurse's ability to provide all of the required care in a timely fashion.

Participants suggested that when there are many unexpected interruptions as well as patient admissions, transfers and discharges, nurses may lose track of what they are doing and miss following up certain patients: "...a lot of care is missed when there is too much to be done in a short time frame" (Participant 54).

The enormity of required documentation was a major concern, with the suggestion that more and more data is expected to be documented, removing autonomy from the nursing role:

...some of it unnecessary and old fashioned:, e.g. rounding (Participant 12).

Rounding was introduced by some hospitals in an effort to reduce the number of times basic but essential care was missed, e.g. pressure care and toileting (Meade, Bursell, & Ketelsen, 2006). Additionally, this was viewed as supporting the reduction of falls especially in the elderly. In rounding, staff are required to check their patients each hour and to sign a checklist of essential nursing care. Participants suggested that the very checking that was implemented to enhance care actually detracts from it because staff are required to stop what they are doing to join the required round at the designated time. The rounding policy took precedence over their own professional judgments about how best to spend their time with patients. It can become a complex juggling act as to what becomes 'essential' and what should be dismissed as 'non-essential' when patients are of a higher acuity or require total assistance in all basic activities such as feeding and mobilisation.

Additional comments related to frustrations around documentation being difficult to comprehend or just not done and this impacted upon the quality of further documentation. Examples of missed or incomplete documentation included: incomplete wound charting, Braden scales (used to check pressure points on skin), falls risk assessments, pressure care, property lists and the actual patient handover sheet. There was a suggestion of incongruity between the teaching of documentation skills and the reality of having the time to produce "perfect documentation":

All our time is taken up by caring for patients with very little time left for paperwork. I often work unpaid overtime to complete this (Participant 7).

These factors put all staff under considerable pressure, contribute to missed nursing care and over the longer term may lead to exhaustion and increased sick leave.

Ineffective methods for determining staffing levels

Missed care related to ineffective methods for determining staffing levels, was identified by a number of participants. A significant number of nurses were dissatisfied with the electronic staffing system in use (ExcelCare) and claimed that it did not allow for the changing needs of the patient as well as inherent daily fluctuations in workload due to many unpredictable daily events. ExcelCare is a registered software programme that allows health services to plan and manage care in accordance with the numbers of nurses and

patients on any given shift (Willis, 2009). The prediction of staffing requirements and allocation is done prospectively and data used for the following shift is based on the previous shift data. This creates a situation whereby there is little latitude in staffing numbers to accommodate any clinically unexpected events that may eventuate on any given day. One nurse commented that medication was always late because ExcelCare did not account for:

...complex patients with multiple medications...the hours don't seem to cover the actual care the patients need...these are people we deal with and their care can't be put down on paper...every person is different but they get their care categorised the same as everyone else (Participant 98).

There was a recurring reference that changes in patient acuity; unplanned admissions and unplanned diagnostic procedures were not adequately factored into ExcelCare at an appropriate level to support the need for additional nursing staff. Participants also commented that not all nursing procedures could be captured by ExcelCare, which led to poorer staffing levels and thus missed care.

ExcelCare allows minimal time for admissions that are not already on the ward so multiple expected surgical patients (who may require full admission, ECG, bloods etc.) do not change our ExcelCare requirements enough to allow for adequate staff (Participant 25).

Nurses and midwives indicated that they found the electronic staffing system to be inadequate to meet the needs of changes in patient acuity and that it did not allow for the clinical judgement or time for patient education, which, according to the survey data is a nursing role that is often missed:

...I can imagine the response to the request for an extra staff member to cover other patient's care while you spend twenty minutes talking with a patient/family member (Participant 17).

Issues of missed care were also associated with the inability of electronic staffing software to incorporate the added work required due to agency and relieving staff that may be unfamiliar with the patients, ward environment and processes. This often resulted in junior or inexperienced staff taking on more responsibilities around workload and decision-making, adding to an already demanding workload.

Inadequate skill mix/staffing numbers

The third predominant issue derived from the data identified inadequate skill mix and staffing numbers. This was prominent in responses from participants working in aged care settings. One nurse claimed that residents were "frequently not fed, changed or toileted" due to inadequate staffing numbers along with issues with "new Registered Nurses having limited English as well as limited clinical experience" (Participant 26). In these environments Registered Nurses (RNs) are being back filled with Enrolled Nurses (ENs) and some of the work is outside their scope of practice. While RNs and ENs are both registered with the Australian Health Practitioner Regulation Agency (AHPRA), ENs usually

undertake training less than the three years required for RN training and work alongside RNs generally performing less complex tasks (AIHW, 2013). This in turn places additional stress on existing experienced staff, which results in these nurses having to take on responsibilities that can be identified as working outside their scope of practice. This places additional stress on existing experienced RNs who must take responsibility for all care provided by ENs as well as their own patient allocation; this has potential for missed or delayed care.

Country SA is cutting our staffing levels, running acute beds with nursing home, hostel and A & E services with only 1 RN and minimal staff. . . . We have 60 beds to look after on night duty with one RN, one EN and one carer (Participant 234).

Missed care related to lack of staff was reported in the aged care setting in particular, with suggestion that residents were not being fed a warm meal, had poor oral hygiene and references were made to the omission of hand washing by staff. Inadequate training and lack of skill was identified by participants as a contributing factor to this missed care.

Issues associated with skill mix were commonly reported in rural hospitals because of difficulties recruiting RNs and the limited scope of practice of ENs:

. . . our Enrolled Nurses are not allowed to do medications regardless of whether they are diploma trained or not. This puts pressure on the 1 or 2 Registered Nurses rostered each shift (Participant 91).

Survey findings also noted that nursing tasks are not being performed according to expertise; for example, one nurse stated, "making beds and emptying skips should be done by carers (also known as Health Support Workers), not nurses" (Participant 72). When a RN is off sick, s/he is often replaced with a more junior staff member or "relieving nurse" with inadequate experience or knowledge, requiring supervision and this leads to a situation where care may be missed.

Some participants indicated that when working as a casual or agency nurse, "permanent staff will often give you a heavy work load 'because you don't deal with these people every day'" (Participant 112) so that not only is the patient allocation and health care system contributing to the situation, so too are the nurses themselves. This is not surprising when other participants stated, "we need more staff to take the pressure off" (Participant 220).

. . . junior staff are no longer given the time to learn the specialty area before being thrust into senior roles, giving dangerous cytotoxic drugs etc. It seems it is only getting worse unfortunately. The senior staff are leaving and the juniors are feeling very pressured and anxious about their roles (Participant 172).

Discussion

This study sought to determine and explore nurses' perceptions of reasons for missed care within the South Australian context and across a variety of healthcare settings. The paper presents an analysis of the qualitative responses from 354 South Australian nurses and midwives. Thematic analysis was used to discern responses and identify themes.

Analysis of survey findings revealed three overriding factors as contributors of missed nursing care. These factors have been identified as: competing demands that reduce time for patient care; ineffective methods for determining staffing levels; and inadequate skill mix including insufficient staff numbers. These three factors describe a tension between what nurses perceive as essential care, staffing allocations and the resultant missed care or delayed care the nurses and midwives describe in their daily practice. Results from the South Australian survey support findings by Kalisch and Williams (2009) in relation to significant reasons for missed care. Nurse and midwife participants often mentioned the variety of daily tasks as interruptions to their daily routine such as visitor requests, impromptu meetings with inter-professional colleagues, the enormity of paperwork and an unpredictable workload with increasing intensity; a formula for missed care.

Having reviewed international literature on models of care, the South Australian study points to the need for a more flexible model of delivery that allows nurses to make decisions pertaining to care based on evidence, best practice and professional judgement (Bakker & Mau, 2012; Cohen & Zimmerman, 2010; Fernandez, Tran, Johnson, & Jones, 2014; Hamilton et al., 2014). This decision making however, must be undertaken within the professional boundary of the nurse/midwife who is required to comply with national competency standards as governed by the Nursing and Midwifery Board of Australia. These competency standards are used as a professional framework to assess competence as a component of annual performance review and registration. Understanding scope of practice and negotiating roles within available skill mix, supports patient safety and promotes effective team environments (Schulter, Seaton, & Chaboyer, 2011).

Skill mix in nursing refers to the level of experience and skill a nurse has and needs to be considered in allocating patients based on acuity. Skill mix is frequently used in conjunction with cost effectiveness and quality of care and can be broadly categorised as "the mix of posts in the establishment; the mix of employees in a post; the combination of skills available at a specific time; or the combinations of activities that comprise each role, rather than the combination of different job titles" (Buchan, Ball, & May, 2001, p. 233). While skill mix is determined by a number of influences, the most prevalent of these is financial where less skilled nurses are used to cut costs (Duffield et al., 2005). Kalisch et al. (2009) identify skill mix as a factor in missed care with RNs more likely to identify nursing care as being missed.

The South Australian State Government recently commissioned the consultancy firms of Deloitte Touche Tohmatsu and KPMG to undertake a budget performance review for the two major urban regions. The findings of these two reviews suggest that hospitals in South Australia have more than adequate nursing and midwifery staff in comparison to peer hospitals in other states (Deloitte Touche Tohmatsu, 2012; KPMG, 2012). The ANMFSA (2012) in response, note that the centralisation of acute care services in South Australia makes averaging of staffing levels across the state problematic and note ongoing issues with limited rostering of senior nursing staff after hours. This survey supports these claims by highlighting nurse and midwives perceptions of missed

or rationed care, with both staffing methods and skill mix identified as significant contributors to missed care.

Limitations of this study

The South Australian study replicated one that had already been carried out in the US. Whilst the survey was modified to meet local conditions, it did not address all the concerns of the participants, as indicated, "your survey does not adequately reflect the situations we're facing in order for me to do the survey justice" (Participant 164). The survey was originally designed for acute care settings. Our sample included participants from community and aged care settings who may experience different barriers to those experienced in acute care. The inclusion of open questions provided scope for identifying the range of issues faced. The issues are complex and extensive, and this needs to be reflected in any follow up study.

Implications for practice

Australia is facing a rapidly changing healthcare landscape where an ageing population and increase in chronic disease means that people are living longer and often require more complex care. The health system and its workers need to evolve in response to these changing demands. The findings presented in this paper suggest nursing staff compromise nursing care by omitting essential tasks as they do not have the resources they need to meet care needs. Lack of nursing time is exacerbated by time spent chasing missing medications and equipment, staffing methods that do not account for changes in patient acuity and skillmix resulting in staffing undertaking tasks outside of their scope of practice and/or level of experience. Better nursing care from this perspective may depend upon prioritising care appropriately, and reviewing the capabilities of those providing care, in light of rationing of staff in response to escalating health delivery costs. Staff should have the appropriate skills and equipment to be adapt to daily changes in workload and patient acuity, and the knowledge to appropriately prioritise and re prioritise care when required.

Furthermore, the study highlights a need to develop workload models that are responsive to changes in patient acuity. South Australia, under the 2013 Enterprise bargaining agreement, has replaced ExcelCare with staffing levels determined on the basis of established hours per patient per day ratios for different patient areas, with a requirement that the skill mix of staff meet minimum standards for the hospital which are set at a 70:30 level except for smaller country sites. Mandated staff ratios have been found in other contexts, to lead to increasing employment of RNs and lower mortality rates (Aiken et al., 2010; Sochalski, Konetzka, Zhu, & Volpp, 2008). Further research is required to see how the new staffing models effect capacity to deliver care.

Finally this research highlights the difficulties faced by rural hospitals and aged care which is not captured adequately by the current tool. Work on missed care has largely been undertaken in acute settings and tool reflects this. The factors impacting capacity to deliver all required nursing care in rural and aged care settings which provide long term

care for older patients and employ less RNs is an area which requires further exploration.

Conclusion

This study has identified the reasons that nurses give for missing nursing care. The study highlights the discontinuity between the cost of care budgeted for and the actual care required by patients and others. Many staff identified competing demands and unpredictability of workload which occurs despite use of ExcelCare to determine staffing in public sector hospitals in South Australia as contributing to missed care. Skill mix and increasing use of ENs and care assistants were also implicated primarily in rural and aged care settings. The findings imply that hospital budgeting is an important factor in determining the adequacy of staffing, supplies, and skill mix. Further studies on missed care within the Australian context are needed to determine if similar findings occur in jurisdictions with alternate staffing methods.

Conflict of interest

None declared.

Acknowledgements

Flinders University Faculty of Health Sciences, Seeding Grant and Robert Wood Johnson Foundation Grant provided funding for this study.

The authors wish to thank the Australian Nursing and Midwifery Federation (SA Branch) executive and participating members for their support. Thanks are also extended to the Wilson and Wood Foundations and Midwestern State University, Wichita Falls, Texas, and the Robert Wood Johnson Foundation's INQRI Program.

References

- Aiken, L., Sloane, D., Cimiotti, J., Clarke, S., Flynn, L., Seago, J. A., et al. (2010). *Implications of the California Nurse Staffing Mandate for other states. Health Services Research, 45*(4), 904–921.
- Australian Institute of Health & Welfare (AIHW). (2013). *Nursing workforce definitions*. Government of Australia. Retrieved from <http://www.aihw.gov.au/nursing-workforce-definitions/>
- Australian Nursing & Midwifery Federation (SA Branch) [ANMFSA]. (2012). *Response to the Hospital budget and remediation report*. Adelaide: ANMFSA.
- Aitkin, L. H., Clarke, S. P., & Sloane, D. M. (2002). *Hospital staffing, organization, and quality of care: Cross-national findings. International Journal for Quality in Health Care, 14*(1), 5–13.
- Bakker, D., & Mau, J. (2012). The nursing model of care: Don't forget the patient perspective. *Nursing Management, 43*(10), 8. <http://dx.doi.org/10.1097/01.NUMA.0000419188.26804.8c>
- Buchan, J., Ball, J., & May, F. (2001). *If changing skill-mix is the answer, what is the question? Journal of Health Service Research Policy, 6*(4), 233–238.
- Chaboyer, W., Wallis, M., Duffield, C., Courtney, M., Seaton, P., Holzhauser, K., et al. (2008). *A comparison of activities undertaken by enrolled and registered nurses on medical wards in Australia: An observational study. International Journal of Nursing Studies, 45*, 1274–1284.

- Cohen, L., & Zimmerman, S. (2010). Evidence behind The Green House and similar models of nursing home care. *Aging Health, 6*(6), 717.
- Deloitte Touche Tohmatsu. (2012). *South Australian health budget performance and remediation review: Northern Adelaide local health network*. South Australia Health. Retrieved from <http://www.sahealth.gov.au/wps/wcm/connect/840e96004c1ca4b49337b7e40887aa/NALHN+Budget+Performance+and+Remediation+Review-OBRI-250712.pdf?MOD=AJPERES&CAHEID=840e96004c1ca4b49337b7e40887aa>
- Denzin, N. K., & Lincoln, Y. S. (Eds.). (2008). *Collecting and interpreting qualitative materials*. Thousand Oaks, CA: SAGE Publications.
- Duffield, C., Forbes, J., Fallon, A., Roche, M., Wise, W., & Merrick, E. A. (2005). Nursing skill mix and nursing time: The roles of registered nurses and clinical nurse specialists. *Australian Journal of Advanced Nursing, 23*(2), 14–21.
- Fernandez, R., Tran, D., Johnson, M., & Jones, S. (2014). Interdisciplinary communication in general medical and surgical wards using two different models of nursing care delivery. *Journal of Nursing Management, 18*, 265–274.
- Hamilton, P., Willis, E., Henderson, J., Harvey, C., Toffoli, L., Abery, E., et al. (2014). Mapping social processes at work in nursing knowledge development. *Nursing & Health Sciences*, <http://dx.doi.org/10.1111/nhs.12106>
- Henderson, J., Blackman, I., Hamilton, P., Willis, E., Toffoli, L., Verrall, C., et al. (2013). *After hours nurse staffing, work intensity and quality of care-missed care study: South Australia*. Final report to the Australian Nursing and Midwifery Federation South Australian Branch.
- Jacob, E., McKenna, L., & D'Amore, A. (2013). The changing skill mix in nursing: Considerations for and against different levels of nurse. *Journal of Nursing Management*, <http://dx.doi.org/10.1111/jonm.12162>
- Kalisch, B., Doumit, M., Lee, K. H., & Zein, J. E. (2013). Missed nursing care, level of staffing, and job satisfaction: Lebanon vs. the United States. *Journal of Nursing Administration, 43*(5), 274–279. <http://dx.doi.org/10.1097/NNA.0b013e31828eebaa>
- Kalisch, B., Gosselin, K., & Choi, S. H. (2012). A comparison of patient care units with high versus low levels of missed nursing care. *Health Care Management Review, 37*(4), 320–328. <http://dx.doi.org/10.1097/HMR.0b013e3182497272e>
- Kalisch, B., Landstrom, G. L., & Hindshaw, A. S. (2009). Missed nursing care: A concept analysis. *Journal of Advanced Nursing, 65*(7), 1509–1517.
- Kalisch, B., & Williams, R. (2009). Development and psychometric testing of a tool to measure missed nursing care. *The Journal of Nursing Administration, 39*(5), 211–219.
- KMPG. (2012). *Southern Adelaide local health network: Hospital budget performance and remediation review*. Adelaide: Final report. South Australian Health. Retrieved from <http://www.sahealth.gov.au/wps/wcm/connect/784f92004c1a3439262b6e408a887aa/SALN+Budget+Performance+and+Remediation+Review-OBRI-250712.pdf?MOD=AJPERES&CAHEID=784f92004c1a3439262b6e408a887aa>
- Meade, C. M., Bursell, A. L., & Ketelsen, L. (2006). Effects of nursing rounds on patients' call light use, satisfaction, and safety. *American Journal of Nursing, 106*(9), 58–70.
- Meyrick, J. (2006). What is good qualitative research? A first step towards a comprehensive approach to judging rigour/quality. *Journal of Health Psychology, 11*(5), 799–808.
- Needleman, J., Buerhaus, P. I., Mattke, S., Stewart, M., & Zelevinsky, K. (2002). Nurse-staffing levels and the quality of care in hospitals. *New England Journal of Medicine, 346*(22), 1715–1712.
- O'Brien-Pallas, L., Duffield, C., & Alknis, C. (2004). Who will be there to nurse? Retention of nurses nearing retirement. *Journal of Nursing Administration, 34*(6), 298–302.
- Papastavrou, E., Panayiota, A., & Georgios, E. (2013). Rationing of nursing care and nurse-patient outcomes: A systematic review of quantitative studies. *International Journal of Health Planning and Management*, <http://dx.doi.org/10.1002/hpm.21860>
- Patton, M. Q. (2002). *Qualitative Research & Evaluation Methods* (3rd ed.). London: Sage Publications.
- Pope, C., Ziebland, S., & Mays, N. (2006). Analysing qualitative data. In C. Pope, & N. Mays (Eds.), *Qualitative research in health care*. Massachusetts: Blackwell Publishing Inc.
- Preston, B. (2009). The Australian nurse and midwifery workforce: Issues, developments and the future. *Collegian, 16*(1), 25–34.
- Productivity Commission. (2005). *Australia's health workforce*. Canberra: Research report. Commonwealth of Australia.
- Schubert, M., Clarke, S., Glass, T., Schaffert-Witvliet, B., & De Geest, S. (2009). Identifying thresholds for relationships between impacts of rationing of nursing care and nurse- and patient-reported outcomes on Swiss hospitals: A correlational study. *Journal of Nursing Studies, 46*, 884–893.
- Schubert, M., Glass, T., Clarke, S., Schaffert-Witvliet, B., & De Geest, S. (2008). Validation of the Basel extent of rationing of nursing care instrument. *Nursing Research, 56*(6), 416–424.
- Schulter, J., Seaton, P., & Chaboyer, W. (2011). Understanding nursing scope of practice: A qualitative study. *International Journal of Nursing Studies, 48*, 1211–1222.
- Sochalski, J., Konetzka, R. T., Zhu, J., & Volpp, K. (2008). Will mandated minimum Nurse Staffing Ratios lead to better patient outcomes? *Medical Care, 46*(6), 606–613.
- Willis, E. M. (2009). *Purgatorial time in hospitals*. Berlin, Germany: LAP Lambert Academic Publishing.