



Subject Index to Volume 54

- ACFAS. *see* American College of Foot and Ankle Surgeons
- Achilles tendinitis
infectious, after local injection of human placental extracts (case report), 1193–1196
- Achilles' tendon
acute, using allograft tendon weaving technique, augmented repair of, 1004–1009
detachment and reattachment, using standard midline approach to insertional enthesophytes, safety of, 214–219
rupture of
acute, early weightbearing on biomechanical properties of plantar flexor muscle–tendon complex in treatment of, 220–226
chronic, reconstructed with Achilles tendon allograft and xenograft combination (case report), 1146–1150
fluoroquinolone-mediated (case report), 494–496
negative pressure wound therapy in management of late deep infections after open reconstruction of, 2–6
neglected, outcome of one-stage tensile stress surgical technique and early postoperative rehabilitation in treatment of, 153–159
using diffusion tensor imaging and tendon fiber tracking, assessment of postoperative tendon quality in patients with, 782–786
surgical suture, gait analysis before and after (case report), 247–250
- Achilles tendon-splitting approach
central, operative treatment of Haglund syndrome with, 1053–1056
- Acute compartment syndrome
of foot, due to infection after local hydrocortisone injection (case report), 692–696
- Adjuvant therapy
surgical outcomes after excision of pigmented villonodular synovitis localized to the ankle and hindfoot without, 160–163
- Allograft
Achilles tendon, and xenograft combination, chronic Achilles tendon rupture reconstructed with (case report), 1146–1150
demineralized subchondral bone, role in treatment of shoulder lesions of talus (case report), 717–722
tendon weaving technique, augmented repair of acute Achilles tendon rupture using, 1004–1009
tricortical wedges, and wedge locking plates, in mid-calcaneal length after Evans calcaneal osteotomy, comparison of, 900–904
- All-suture anchor
use for re-creation of anterior talofibular ligament (case report), 126–129
- American Academy of Orthopedic Surgeons Foot and Ankle Questionnaire
Korean version, validity, reliability, and responsiveness of, 46–50
- American College of Foot and Ankle Surgeons
clinical consensus statement in perioperative prophylactic antibiotic use in clean elective foot surgery, 273–279
- Amputation
lower extremity, “jumping stump” phenomenon, 517–518
lower extremity, risk factors associated with elevated ankle brachial indices and radiographic arterial calcification, 473–477
partial hallux, influence of hallux rigidus on reamputation in patients with diabetes mellitus after, 1076–1080
partial metatarsal, radiation therapy for recurrent heterotopic ossification prophylaxis after, 345–349
- Angle stable device
percutaneous, early minimally invasive percutaneous fixation of displaced intra-articular calcaneal fractures with, 51–56
- Ankle
anterolateral impingement syndrome of, effect of obesity on arthroscopic treatment of, 13–16
arthrodesis
biomechanical comparison of internal fixation techniques for, 188–191
functional and clinical assessment of two techniques of, 399–405
asymptomatic lateral, to determine anatomic variants and incidental peroneal tendon pathologic features, MRI of, 164–172
branching patterns of superficial peroneal nerve on implications for ankle arthroscopy and for anterolateral surgical approaches to, 332–337
dislocation of, pathogenesis and treatment strategies for pilon fractures with, 815–820
and foot
arthrodesis, National Survey of Ambulatory Surgery and National Hospital Discharge Survey in, 1037–1041
Blastomyces tenosynovitis of (case report), 1183–1187
giant cell tumors of, high recurrence rates after surgical treatment in (case report), 1141–1145
surgery, and injuries requiring immobilization, risk, prevention, and diagnosis of venous thromboembolism disease in, 497–507
surgery, elective, pain management for, a systematic review, 625–635
surgery, in diabetic patients who have undergone solid organ transplantation, outcomes of, 577–581
surgery, modified suture bridge technique for application with bone anchors in, 944–948
surgery, unplanned readmission following (editorial), 287

- fractures of
 irreducible, dislocation due to tibialis anterior subluxation (case report), 268–272
 isolated, 22-modifier in, 192–197
 low-energy open, in elderly, 203–206
 and modality of hospital transport at single level 1 trauma center, 826–829
 posterior malleolar, radiographic assessment of, 365–369
 repair of acute deltoid ligament complex rupture associated with, 198–202
 supination external rotation, manual stress ankle radiography to predict deep deltoid ligament integrity in, 531–535
 uncertain stability, weightbearing radiographs facilitate functional treatment of, 1042–1046
 Weber B, use of flexible implant and bioabsorbable anchor for deltoid rupture repair in bimalleolar equivalent, 513–516
 Weber C, retrospective audit of patients with, 454–457
 and hindfoot, without adjuvant therapy, surgical outcomes after excision of pigmented villonodular synovitis localized to, 160–163
 injury, Bosworth-Pilon fracture an unusual (case report), 751–753
 instability
 chronic, in young adults, epidemiology and risk factors for, 183–187
 chronic, ligament reattachment technique for high-demand athletes with, 7–12
 irreducible fracture-dislocation of, associated with interposition of tibialis posterior tendon in syndesmosis (case report), 962–966
 mortise, posterior tibial tendon entrapment within an intact (case report), 116–119
 osteoarthritis
 computed tomography atlas for grading, 207–213
 current concepts in management of, a systematic review, 932–939
 periarticular osteoid osteoma of, treated with arthroscopy, 89–93
 Ankle arthroplasty
 total
 in preoperative and postoperative range of motion, comparison of, 809–814
 quantification of subtalar posterior facet involvement during intramedullary guidance of, 805–808
 registry data trends of using, 1213–1142(Letter/Reply)
- Ankle brachial index
 elevated, and radiographic arterial calcification, lower extremity amputation risk factors associated with, 473–477
- Ankle cartilage allograft replacement
 evidence-based rationale for, a systematic review, 940–943
- Ankle deformity
 and foot, progressive, associated with Charcot-Marie-Tooth disease, minimally invasive early operative treatment of (case report), 701–708
- Ankle fusion
 and foot fusion, effect of osteogenic progenitor cell concentration on incidence of, 888–891
 with trabecular metal spacer and anterior fusion plate (case report), 490–493
 use of spinal cage for creating stable constructs in (case report), 254–257
- Ankle joint
 distraction arthroplasty with arthroscopic microfracture in patient with rheumatoid arthritis of, 280–284
 high, after fibular graft harvesting, open dislocation of (case report), 1158–1161
- Ankle syndesmosis
 osteomyelitis after TightRope® fixation of (case report), 130–134
- Anterolateral impingement syndrome
 of ankle, effect of obesity on arthroscopic treatment of, 13–16
- Anterolateral incision
 single distal, minimally invasive plate osteosynthesis of distal tibia and fibular fractures through, 1081–1084
- Antibiotic prophylaxis
 perioperative, use in clean elective foot surgery, ACFAS clinical consensus statement in, 273–279
- Antibiotic treatment
 and surgery, for acute hematogenous calcaneal osteomyelitis of childhood, 840–843
- Arbeitsgemeinschaft für Osteosynthesefragen (AO) 43-C
 tibial plafond fractures, hybrid external fixation for, 1031–1036
- Arterial calcification
 radiographic, lower extremity amputation risk factors associated with elevated ankle brachial indices and, 473–477
- Arthrex TightRope®
 fixation of ankle syndesmosis, osteomyelitis after (case report), 130–134
- Arthrodesis
 of ankle
 biomechanical comparison of internal fixation techniques for, 188–191
 functional and clinical assessment of two techniques of, 399–405
 arthroscopic, of distal tibiofibular syndesmosis, 953–957
 distraction, subtalar joint, to correct calcaneal valgus in pediatric patients with tarsal coalition (case report), 1151–1157
 double versus triple, hindfoot deformity corrected with, 424–427
 of first metatarsophalangeal joint, arthroscopic, for repair of fixed hallux varus deformity, 1127–1131
 of foot and ankle, National Survey of Ambulatory Surgery and National Hospital Discharge Survey in, 1037–1041
 versus hemiarthroplasty, for end-stage hallux rigidus, short-term clinical outcome of, 848–851
 versus hemiarthroplasty, for osteoarthritis of first metatarsophalangeal joint, 1085–1088
 medial column, using an anatomic distal fibular locking plate, 671–676
 medial double, with lateral column sparing and arthrodiastasis, 441–444
 medial double and Lapidus, on arthrodiastasis of calcaneocuboid joint, comparative findings after, 412–416
 primary subtalar joint, for comminuted fractures of calcaneus, 61–65
 of subtalar joint, subsequent, and calcaneal fractures, recreational sports activities after, 1057–1061
 of tarsometatarsal, first, anatomic evaluation of dorsomedial versus plantar plating in, 787–792
 tibiocalcaneal, with porous tantalum spacer and locked intramedullary nail for post-traumatic global avascular necrosis of talus (case report), 1172–1177
 tibiototalcalcaneal
 early results of posterolateral polyaxial angle-stable plate for, 428–432
 using retrograde intramedullary nail fixation in comparison of patients with and without diabetes mellitus, 876–882
- Arthrodiastasis
 of calcaneocuboid joint, comparative findings after medial double and Lapidus arthrodesis on, 412–416
 and lateral column sparing, medial double arthrodesis with, 441–444

- Arthroplasty**
 autogenous capsular interpositional, for painful hallux rigidus, 29–36
 distraction, with arthroscopic microfracture, in patient with rheumatoid arthritis of ankle joint, 280–284
 of hallux interphalangeal joint, compared with nonoperative treatment of recalcitrant hallux ulceration, evaluation of, 541–548
- Arthroscopy**
 of ankle, and for anterolateral surgical approaches to ankle, branching patterns of superficial peroneal nerve on implications for, 332–337
 arthrodesis of distal tibiofibular syndesmosis, 953–957
 first metatarsophalangeal arthrodesis for repair of fixed hallux varus deformity, 1127–1131
 microfracture, in patient with rheumatoid arthritis of ankle joint, distraction arthroplasty with, 280–284
 periarticular osteoid osteoma of ankle treated with, 89–93
 subtalar debridement, and lateral calcaneal osteotomy, for residual pain after calcaneal fracture, simultaneous use of, 37–40
 for treatment of anterolateral impingement syndrome of ankle, effect of obesity on, 13–16
 treatment of osteochondritis dissecans of talus, outcomes are favorable after, 57–60
- Athletes**
 adolescent, medial malleolar stress fracture in (case report), 242–246
 high-demand, with chronic ankle instability, ligament reattachment technique for, 7–12
- Atlas**
 computed tomography, for grading ankle osteoarthritis, 207–213
- Ballroom dancer**
 heel lipoma mimicking plantar fasciitis in (case report), 251–253
- Beam screw**
 medial column, in midfoot Charcot reconstruction, posterior approach for, 433–436
- Bioabsorbable anchor**
 and flexible implant, use for deltoid rupture repair in bimalleolar equivalent Weber B ankle fractures, 513–516
- Biomechanics**
 evaluation of custom foot orthoses for hallux valgus deformity, 852–855
 evaluation of plate versus lag screw only fixation of distal fibula fractures, 896–899
 performance of lateral versus dual locking plates for calcaneal fractures, 830–835
- Bipolar radiofrequency microtenotomy**
 intermediate-term results of partial plantar fascia release with microtenotomy using bipolar radiofrequency microtenotomy, 179–182
- Blastomyces**
 tenosynovitis of foot and ankle (case report), 1183–1187
- Blood loss**
 postoperative, in calcaneal fractures, administration of tranexamic acid reduces, 1106–1110
- Bone anchors**
 in foot and ankle surgery, modified suture bridge technique for application with, 944–948
- Bone coalitions**
 Botulinum toxin type A injection combined with cast immobilization for treating recurrent peroneal spastic flatfoot without (case report), 697–700
- Bone cyst**
 unicameral
 in calcaneus, of mirror image twins (case report), 754–757
 of calcaneus, treatment of, a systematic review, 652–656
- Bone graft**
 for treatment of brachymetatarsia, callus distraction versus single-stage lengthening with, a systematic review, 927–931
- Bosworth-Pilon fracture**
 an unusual ankle injury (case report), 751–753
- Botulinum toxin type A injection**
 combined with cast immobilization, for treating recurrent peroneal spastic flatfoot without bone coalitions (case report), 697–700
- Brachymetatarsia**
 callus distraction versus single-stage lengthening with bone graft for treatment of, a systematic review, 927–931
- Braking function**
 effects of right lower limb orthopedic immobilization on, 554–558
- Bunion correction**
 complications of metatarsal suture techniques for, a systematic review, 230–232
- Bunion deformity**
 evaluation and management of, 102–111
- Calcaneal spurs**
 classification of, and relationship with plantar fasciitis, 594–600
- Calcaneal valgus**
 in pediatric patients with tarsal coalition, subtalar joint distraction arthrodesis to correct (case report), 1151–1157
- Calcaneocuboid joint**
 arthrodiastasis of, comparative findings after medial double and Lapidus arthrodesis on, 412–416
- Calcaneofibular ligament injury**
 isolated (case report), 487–489
- Calcaneotalar ratio**
 in estimation of length of calcaneus, 370–372
- Calcaneus**
 calcaneotalar ratio in estimation of length of, 370–372
 chondroblastoma of, with pathologic fracture and recurrence (case report), 258–267
 fractures of
 administration of tranexamic acid reduces postoperative blood loss in, 1106–1110
 biomechanical performance of lateral versus dual locking plates for, 830–835
 comminuted, primary subtalar joint arthrodesis for, 61–65
 constant fragment, screw placement relative to, 392–394
 displaced intra-articular, functional outcome and patient satisfaction after, 298–305
 displaced intra-articular, percutaneous arthroscopic calcaneal osteosynthesis a minimally invasive technique for, 798–804
 displaced intra-articular, with percutaneous angle stable device, early minimally invasive percutaneous fixation of, 51–56
 intra-articular, in children, minimally invasive (sinus tarsi) approach for open reduction and internal fixation of (case report), 135–139
 of Sanders type II and III, by minimally invasive technique using locking plate, surgical treatment of, 76–81

- with severe soft tissue injury and osteomyelitis, treatment of (case report), 973–977
- simultaneous use of lateral calcaneal osteotomy and subtalar arthroscopic debridement for residual pain after, 37–40
- and subsequent subtalar joint arthrodesis, recreational sports activities after, 1057–1061
- superior peroneal retinacular injuries in, 458–463
- osteotomy
- double, with percutaneous Steinmann pin fixation as part of treatment for flexible flatfoot deformity (case report), 478–482
- Evans, comparison of wedge locking plates and tricortical allograft wedges in mid-calcaneal length after, 900–904
- medial displacement, anatomy of sural nerve with an emphasis on incision for, 341–344
- minimally invasive, using Shannon burr endanger the neurovascular structures, 1062–1066
- percutaneous posterior (case report), 1188–1192
- unicameral bone cyst
- of mirror image twins (case report), 754–757
- treatment of, a systematic review, 652–656
- Callus distraction
- versus single-stage lengthening, with bone graft for treatment of brachymetatarsia, a systematic review, 927–931
- Cancellous bone fixation
- fully threaded versus partially threaded screws to prevent shear in, 1021–1024
- Case reports
- acute compartment syndrome of the foot due to infection after local hydrocortisone injection: a case report, 692–696
- ankle fusion with a trabecular metal spacer and an anterior fusion plate, 490–493
- an unusual ankle injury: the Bosworth-Pilon fracture, 751–753
- Blastomyces* tenosynovitis of the foot and ankle: a case report and review of the literature, 1183–1187
- Botulinum toxin type A injection combined with cast immobilization for treating recurrent peroneal spastic flatfoot without bone coalitions: a case report and review of the literature, 697–700
- calcaneal chondroblastoma with pathologic fracture and recurrence, 258–267
- chronic Achilles tendon rupture reconstructed with Achilles tendon allograft and xenograft combination, 1146–1150
- congenital extraskeletal osteochondroma of the web space displacing the lesser toes: a unique presentation, 726–729
- correction of idiopathic adult hallux varus by tendon transfer, 1197–1201
- coverage of exposed bone of the lateral malleolus with a proximally based lateral malleolar perforator flap, 730–733
- double calcaneal osteotomy with percutaneous Steinmann pin fixation as part of treatment for flexible flatfoot deformity: a review of consecutive cases highlighting our experience with pin fixation, 478–482
- excision of elephantiasis nostras verrucosa lesions in a patient with hereditary lymphedema: case report and review of the literature, 747–750
- fluoroquinolone-mediated Achilles rupture: a case report and review of the literature, 494–496
- gait analysis before and after Achilles tendon surgical suture in a single-subject study: a case report, 247–250
- giant cell tumor of the talus: 19-year follow-up of a patient, 958–961
- giant cell tumors of the foot and ankle bones: high recurrence rates after surgical treatment, 1141–1145
- heel lipoma mimicking plantar fasciitis in a ballroom dancer, 251–253
- infectious Achilles tendinitis after local injection of human placental extracts: a case report, 1193–1196
- interpositional arthroplasty with palmaris longus tendon graft for osteonecrosis of the second metatarsal head: a case report, 237–241
- irreducible ankle fracture-dislocation due to tibialis anterior subluxation: a case report, 268–272
- irreducible fracture-dislocation of the ankle associated with interposition of the tibialis posterior tendon in the syndesmosis: a case report, 962–966
- isolated calcaneofibular ligament injury: a report of two cases, 487–489
- isolated fifth metatarsocuboid coalition: a case report, 734–738
- lateral stress dorsiflexion view: a case series demonstrating clinical utility in midterm hallux limitus, 739–746
- management of idiopathic clubfoot by Ponseti technique in children presenting after one year of age, 967–972
- medial malleolar stress fracture in an adolescent athlete, 242–246
- minimally invasive early operative treatment of progressive foot and ankle deformity associated with Charcot-Marie-Tooth disease, 701–708
- minimally invasive (sinus tarsi) approach for open reduction and internal fixation of intra-articular calcaneus fractures in children: surgical technique and case report of two patients, 135–139
- multifocal bilateral metatarsal tuberculosis: a rare presentation, 112–115
- open dislocation of the high ankle joint after fibular graft harvesting, 1158–1161
- osteomyelitis after TightRope® fixation of the ankle syndesmosis: a case report and review of the literature, 130–134
- partial transmalleolar approach for lateral impingement after total ankle arthroplasty: a case report, 233–236
- percutaneous posterior calcaneal osteotomy, 1188–1192
- pigmented onychomatricoma: a rare pigmented nail unit tumor presenting as longitudinal melanonychia that has potential for misdiagnosis as melanoma, 723–725
- posterior tibial tendon entrapment within an intact ankle mortise: a case report, 116–119
- pulmonary pleomorphic carcinoma metastasis to the midfoot, 483–486
- reconstruction of the pediatric lateral malleolus and physis by free microvascular transfer of the proximal fibular physis, 994–1000
- revision subtalar joint fusion with a porous metal spacer and an intramedullary nail: a case report, 709–712
- role of demineralized allograft subchondral bone in the treatment of shoulder lesions of the talus: clinical results with two-year follow-up, 717–722
- subtalar joint distraction arthrodesis to correct calcaneal valgus in pediatric patients with tarsal coalition: a case series, 1151–1157
- surgical considerations for massive tarsal coalitions in multiple synostosis syndrome: a case report, 1162–1165
- surgical treatment guidelines for digital deformity associated with intrinsic muscle spasticity (intrinsic plus foot) in adults with cerebral palsy, 985–993
- symptomatic osseous abnormalities at the posteromedial tubercle of the talus: magnetic resonance imaging features, 978–984
- talar fractures in children: a possible injury after go-karting accidents, 1206–1212

- talocalcaneal joint middle facet coalition resection with interposition of a juvenile hyaline cartilage graft, 1178–1182
- tenosynovial osteochondromatosis of the flexor hallucis longus tendon treated by tendoscopy, 758–764
- tibiocalcaneal arthrodesis with a porous tantalum spacer and locked intramedullary nail for post-traumatic global avascular necrosis of the talus, 1172–1177
- treatment of calcaneal fracture with severe soft tissue injury and osteomyelitis: a case report, 973–977
- treatment of foot degloving injury with aid of negative pressure wound therapy and dermal regeneration template, 1132–1135
- treatment of longitudinal epiphyseal bracket by excision and polymethylmethacrylate insertion at the preossified disease stage, 1136–1140
- tuberculosis of the cuboid: a case report and review of the literature, 713–716
- unicameral bone cyst in the calcaneus of mirror image twins, 754–757
- unilobed rotational flap for plantar hallux interphalangeal joint ulceration complicated by osteomyelitis, 1166–1171
- use of a definitive cement spacer for simultaneous bony and soft tissue reconstruction of mid- and hindfoot diabetic neuroarthropathy: a case report, 120–125
- use of a locked fibular plate for fixation of a vertical shear medial malleolus fracture: a case report, 1202–1205
- use of an all-suture anchor for re-creation of anterior talofibular ligament: a case report, 126–129
- use of a spinal cage for creating stable constructs in ankle and subtalar fusion, 254–257
- Cast immobilization
- Botulinum toxin type A injection combined with, for treating recurrent peroneal spastic flatfoot without bone coalitions (case report), 697–700
- Cement spacer
- definitive, use for simultaneous bony and soft tissue reconstruction of mid- and hindfoot diabetic neuroarthropathy (case report), 120–125
- Cerebral palsy
- surgical treatment guidelines for digital deformity associated with intrinsic muscle spasticity (intrinsic plus foot) in adults with (case report), 985–993
- Charcot foot
- reconstruction of, posterior approach for medial column beam screw in, 433–436
- Charcot-Marie-Tooth disease
- minimally invasive early operative treatment of progressive foot and ankle deformity associated with (case report), 701–708
- Charcot reconstruction
- comparison of complications for internal and external fixation for, 1072–1075
- Cheilectomy
- for degenerative joint disease of first metatarsophalangeal joint, long-term follow-up of, 1010–1020
- Chevron osteotomy
- distal, role of fixation and postoperative regimens in long-term outcomes of, 356–360
 - versus Mitchell, in hallux valgus surgery, 361–364
 - proximal and distal, for hallux valgus deformity, change in first metatarsal length after, 525–530
- Chondroblastoma
- of calcaneus, with pathologic fracture and recurrence (case report), 258–267
- Clarification
- Two Articles: Similar, Not the Same, 1003
- Clubfoot
- idiopathic, by Ponseti technique in children presenting after one year of age, management of (case report), 967–972
- Column sparing
- lateral, and arthrodiastasis, medial double arthrodesis with, 441–444
- Commentaries
- College Commences Clinical Consensus Statement Development, 151
 - The Importance of Biomechanics, 1215
 - Metallic Hardware Not a Considered Complication, 523
 - The Difference Is in What Is or Is Not Included, 523
 - No Resident Left Behind: Statistical Illiteracy and What Lies Beyond the Abstract, 523–524
- Compartment syndrome
- after fasciotomy, surgical outcomes and quality of life in severe foot trauma and associated, 417–423
- Computed tomography
- atlas, for grading ankle osteoarthritis, 207–213
- Congenital talipes equinovarus
- prognostic value of radiologic appearance of navicular ossification center in, 844–847
- Corticosteroid injection
- and extracorporeal shock wave therapy, as primary therapeutic methods for acute plantar fasciitis, treatment outcomes of, 1047–1052
 - targeting of plantar fascia for, 683–685
- Crossed screw fixation
- versus endosseous fixation device, for Lapidus arthrodesis, 1099–1105
 - with middle cuneiform fixation reduces sagittal mobility in enhanced Lapidus arthrodesis, 437–440
 - modified Lapidus arthrodesis with, 69–75
- Cuboid
- tuberculosis of (case report), 713–716
- Cuneiform fixation
- middle, reduces sagittal mobility in enhanced Lapidus arthrodesis, crossed screw technique with, 437–440
- Debridement
- subtalar arthroscopic, and lateral calcaneal osteotomy, for residual pain after calcaneal fracture, simultaneous use of, 37–40
- Deep deltoid ligament
- integrity, in supination external rotation ankle fracture, manual stress ankle radiography to predict, 531–535
- Degenerative joint disease
- of first metatarsophalangeal joint, long-term follow-up of cheilectomy for, 1010–1020
- Deltoid ligament complex rupture
- acute, associated with ankle fractures, repair of, 198–202
- Dermal regeneration template
- and negative pressure wound therapy, treatment of foot degloving injury with aid of (case report), 1132–1135
- Diabetes mellitus
- after partial hallux amputation, influence of hallux rigidus on reamputation in patients with, 1076–1080
 - conservative surgical treatment of infected ulceration of first metatarsophalangeal joint with osteomyelitis in patients with, 536–540
 - patients who have undergone solid organ transplantation, outcomes of foot and ankle surgery in, 577–581
 - tibiotalocalcaneal arthrodesis using retrograde intramedullary nail fixation in comparison of patients with and without, 876–882

- Diabetic foot
 medical data, bibliometric investigation was to examine, 295–297
 ulcers, plantar lateral column, tibialis anterior tendon lengthening for adjunctive treatment of, 686–691
- Diabetic neuroarthropathy
 mid- and hindfoot, use of definitive cement spacer for simultaneous bony and soft tissue reconstruction of (case report), 120–125
- Diagnostic accuracy
 of clinical tests for Morton's neuroma compared with ultrasonography, 549–553
- Diffusion tensor imaging
 and tendon fiber tracking, assessment of postoperative tendon quality in patients with Achilles tendon rupture using, 782–786
- Digital deformity
 associated with intrinsic muscle spasticity (intrinsic plus foot) in adults with cerebral palsy, surgical treatment guidelines for (case report), 985–993
- Dimeglio idiopathic clubfoot
 classifications of, clinical evaluation of, 582–585
- Dislocation
 of ankle, pathogenesis and treatment strategies for pilon fractures with, 815–820
 of irreducible ankle fracture, due to tibialis anterior subluxation (case report), 268–272
 open, of high ankle joint after fibular graft harvesting (case report), 1158–1161
- Dorsalis pedis artery
 to tarsal navicular, relationship of, 66–68
- Dorsal transversely oriented transposition flap
 for web reconstruction in toe syndactyly surgery, technique of, 1119–1123
- Dorsomedial plating
 versus plantar plating, in first tarsometatarsal arthrodesis, anatomic evaluation of, 787–792
- Double plating
 for management of complex fibular fractures, 288–294
- Editorials
 Mouse Clicks and Key Strokes, 767
 Pooling Results by Means of Meta-Analysis, 521–522
 Unplanned Readmission Following Foot and Ankle Surgery, 287
 White Papers, Position Papers, Clinical Consensus Statements, and Clinical Practice Guidelines: Future Directions for ACFAS, 151–152
 Whither Biomechanics, 1
- Elastic bandage traction technique
 for reduction of distal tibial fractures, 949–952
- Elective foot surgery
 clean, ACFAS clinical consensus statement in perioperative prophylactic antibiotic use in, 273–279
- Elephantiasis nostras verrucosa
 excision in patient with hereditary lymphedema (case report), 747–750
- Enchondroma
 of the foot, 836–839
- Endoscopic release
 modified, of gastrocnemius aponeurosis, 140–142
- Endosseous fixation device
 versus crossed screw fixation, for Lapidus arthrodesis, 1099–1105
- Epidemiology
 of high-heel shoe injuries in U.S. women, 615–619
 and risk factors for chronic ankle instability in young adults, 183–187
- Epiphyseal bracket
 longitudinal, treatment by excision and polymethylmethacrylate insertion at preossified disease stage (case report), 1136–1140
- Equinus deformity
 severe, rigid, use of Ilizarov external fixation without soft tissue release to correct, 821–825
- Evans calcaneal osteotomy
 comparison of wedge locking plates and tricortical allograft wedges in mid-calcaneal length after, 900–904
- External fixation
 Ilizarov, without soft tissue release to correct severe, rigid equinus deformity, use of, 821–825
- External fixator
 limited internal fixation combined with, versus open reduction and internal fixation, for tibial plafond fracture, postoperative complications between, 646–651
- Extracorporeal shock wave therapy
 and corticosteroid injection, as primary therapeutic methods for acute plantar fasciitis, treatment outcomes of, 1047–1052
- Eye tracking
 and gaze patterns, differences between trainees and experts reading plain film bunion radiographs, 382–391
- Fasciotomy
 surgical outcomes and quality of life in severe foot trauma and associated compartment syndrome after, 417–423
- Fibula
 fractures of
 distal, biomechanical evaluation of plate versus lag screw only fixation of, 896–899
 through single distal anterolateral incision, minimally invasive plate osteosynthesis of, 1081–1084
- Fibular graft harvesting
 open dislocation of high ankle joint after (case report), 1158–1161
- Fibular locking plate
 distal, medial column arthrodesis using an anatomic, 671–676
 use for fixation of vertical shear medial malleolus fracture (case report), 1202–1205
- Fibular physis
 proximal, reconstruction of pediatric lateral malleolus and physis by free microvascular transfer of (case report), 994–1000
- Fifth digit derotation arthroplasty
 with absorbable fixation pin, stabilization of, 1116–1118
- Fixation
 cancellous bone, fully threaded versus partially threaded screws to prevent shear in, 1021–1024
 headless screw and locking plate, for talar neck fractures, mechanical comparison of, 905–909
 hybrid external, for Arbeitsgemeinschaft für Osteosynthesefragen (AO) 43-C tibial plafond fractures, 1031–1036
 interference repair, for plantar plate ligament disruption of lesser metatarsophalangeal joints, technique using, 508–512
 internal and external, for Charcot reconstruction, comparison of complications for, 1072–1075
 intramedullary nail, retrograde, in comparison of patients with and without diabetes mellitus, tibiotalar calcaneal arthrodesis using, 876–882
 lagged syndesmotom, our clinical experience, 773–781

- percutaneous, early minimally invasive, of displaced intra-articular calcaneal fractures with percutaneous angle stable device, 51–56
- plate versus lag screw only, of distal fibula fractures, biomechanical evaluation of, 896–899
- and postoperative regimens, role in long-term outcomes of distal chevron osteotomy, 356–360
- TightRope[®], of ankle syndesmosis, osteomyelitis after (case report), 130–134
- of vertical shear medial malleolus fracture, use of locked fibular plate for (case report), 1202–1205
- Fixation pin**
absorbable, stabilization of fifth digit derotation arthroplasty with, 1116–1118
- Flaps**
digital artery, reconstruction of weightbearing forefoot defects with, 41–45
- dorsal transversely oriented transposition, for web reconstruction in toe syndactyly surgery, technique of, 1119–1123
- free perforator, traumatic forefoot reconstructions with, 1025–1030
- unilobed rotational, for plantar hallux interphalangeal joint ulceration complicated by osteomyelitis (case report), 1166–1171
- Flatfoot deformity**
flexible, double calcaneal osteotomy with percutaneous Steinmann pin fixation as part of treatment for (case report), 478–482
- Flexor hallucis longus tendon**
treated by tendoscopy, tenosynovial osteochondromatosis of (case report), 758–764
- Fluoroquinolone**
mediated Achilles rupture (case report), 494–496
- Foot**
279, with 462 multiple interdigital neuromas, retrospective study of, 320–322
- acute compartment syndrome of, due to infection after local hydrocortisone injection (case report), 692–696
- and ankle
arthrodesis, National Survey of Ambulatory Surgery and National Hospital Discharge Survey in, 1037–1041
- Blastomyces* tenosynovitis of (case report), 1183–1187
- giant cell tumors of, high recurrence rates after surgical treatment in (case report), 1141–1145
- surgery, and injuries requiring immobilization, risk, prevention, and diagnosis of venous thromboembolism disease in, 497–507
- surgery, elective, pain management for, a systematic review, 625–635
- surgery, in diabetic patients who have undergone solid organ transplantation, outcomes of, 577–581
- surgery, modified suture bridge technique for application with bone anchors in, 944–948
- surgery, unplanned readmission following (editorial), 287
- development of radiographic union score for determining osteotomy union rates in long bones of, 793–797
- enchondroma of, 836–839
- rigid equinovarus, severe, midterm follow-up of talectomy for, 1093–1098
- Foot deformity**
and ankle, progressive, associated with Charcot-Marie-Tooth disease, minimally invasive early operative treatment of (case report), 701–708
- Foot degloving injury**
with aid of negative pressure wound therapy and dermal regeneration template, treatment of (case report), 1132–1135
- Foot fusion**
and ankle fusion, effect of osteogenic progenitor cell concentration on incidence of, 888–891
- Foot orthoses**
custom, for hallux valgus deformity, biomechanical evaluation of, 852–855
- Foot trauma**
severe, and associated compartment syndrome, after fasciotomy, surgical outcomes and quality of life in, 417–423
- Forefoot**
reconstructions, traumatic, with free perforator flaps, 1025–1030
- Forefoot defects**
weightbearing, with digital artery flaps, reconstruction of, 41–45
- Forefoot surgery**
in elderly compared with younger patient populations, 586–590
- Fracture**
of ankle
irreducible, dislocation due to tibialis anterior subluxation (case report), 268–272
- isolated, 22-modifier in, 192–197
- low-energy open, in elderly, 203–206
- and modality of hospital transport at single level 1 trauma center, 826–829
- posterior malleolar, radiographic assessment of, 365–369
- repair of acute deltoid ligament complex rupture associated with, 198–202
- supination external rotation, manual stress ankle radiography to predict deep deltoid ligament integrity in, 531–535
- uncertain stability, weightbearing radiographs facilitate functional treatment of, 1042–1046
- Weber B, use of flexible implant and bioabsorbable anchor for deltoid rupture repair in bimalleolar equivalent, 513–516
- Weber C, retrospective audit of patients with, 454–457
- of Arbeitsgemeinschaft für Osteosynthesefragen (AO) 43-C tibial plafond, hybrid external fixation for, 1031–1036
- Bosworth-Pilon, an unusual ankle injury (case report), 751–753
- calcaneal
administration of tranexamic acid reduces postoperative blood loss in, 1106–1110
- biomechanical performance of lateral versus dual locking plates for, 830–835
- comminuted, primary subtalar joint arthrodesis for, 61–65
- constant fragment, screw placement relative to, 392–394
- displaced intra-articular, functional outcome and patient satisfaction after, 298–305
- displaced intra-articular, percutaneous arthroscopic calcaneal osteosynthesis a minimally invasive technique for, 798–804
- displaced intra-articular, with percutaneous angle stable device, early minimally invasive percutaneous fixation of, 51–56
- intra-articular, in children, minimally invasive (sinus tarsi) approach for open reduction and internal fixation of (case report), 135–139
- of Sanders type II and III, by minimally invasive technique using locking plate, surgical treatment of, 76–81
- with severe soft tissue injury and osteomyelitis, treatment of (case report), 973–977
- simultaneous use of lateral calcaneal osteotomy and subtalar arthroscopic debridement for residual pain after, 37–40
- and subsequent subtalar joint arthrodesis, recreational sports activities after, 1057–1061

- superior peroneal retinacular injuries in, 458–463
- complex fibular, double plating for management of, 288–294
- of distal tibia and fibular, through single distal anterolateral incision, minimally invasive plate osteosynthesis of, 1081–1084
- distal tibial, elastic bandage traction technique for reduction of, 949–952
- of fibula, distal, biomechanical evaluation of plate versus lag screw only fixation of, 896–899
- of fifth metatarsal base, fracture mechanism and treatment algorithms, anatomic evaluation of, 94–98
- Jones, fifth metatarsal, screw fixation diameter for, 227–229
- of lateral malleolar, specialized fibular locking plate for, 1067–1071
- of malleolar, vertical shear medial, use of locked fibular plate for fixation of (case report), 1202–1205
- pathologic, and recurrence, calcaneal chondroblastoma with (case report), 258–267
- pilon, with ankle dislocation, pathogenesis and treatment strategies for, 815–820
- of posterior malleolar and lateral malleolar, single oblique posterolateral approach for open reduction and internal fixation of, 559–564
- stress, of medial malleolar, in an adolescent athlete (case report), 242–246
- of talar, in children after go-karting accidents (case report), 1206–1212
- of talar neck
 - irreducible Hawkins type III, open reduction of, 677–682
 - mechanical comparison of headless screw fixation and locking plate fixation for, 905–909
 - simple displaced, antegrade-retrograde opposing lag screws for internal fixation of, 23–28
- tibial plafond, postoperative complications between open reduction and internal fixation versus limited internal fixation combined with external fixator for, 646–651
- Free microvascular transfer
 - of proximal fibular physis, reconstruction of pediatric lateral malleolus and physis by (case report), 994–1000
- Fusion plate
 - anterior, and trabecular metal spacer, ankle fusion with (case report), 490–493
- Gait analysis
 - before and after Achilles tendon surgical suture (case report), 247–250
- Gastrocnemius aponeurosis
 - modified endoscopic release of, 140–142
- Gaze patterns
 - and eye tracking, differences between trainees and experts reading plain film bunion radiographs, 382–391
- Giant cell tumor
 - of foot and ankle bones, high recurrence rates after surgical treatment in (case report), 1141–1145
 - of talus (case report), 958–961
- Global avascular necrosis
 - of talus, post-traumatic, tibio calcaneal arthrodesis with porous tantalum spacer and locked intramedullary nail for (case report), 1172–1177
- Go-karting accidents
 - talar fractures in children after (case report), 1206–1212
- Haglund's deformity
 - radiographic evaluation of, 338–340
- Haglund syndrome
 - with central Achilles tendon-splitting approach, operative treatment of, 1053–1056
- Hallux abducto valgus
 - iatrogenic, secondary to tibial sesamoidectomy, 82–88
- Hallux interphalangeal joint
 - arthroplasty, compared with nonoperative treatment of recalcitrant hallux ulceration, evaluation of, 541–548
 - plantar, ulceration complicated by osteomyelitis, unilobed rotational flap for (case report), 1166–1171
- Hallux limitus
 - midterm, lateral stress dorsiflexion view demonstrating clinical utility in (case report), 739–746
- Hallux rigidus
 - end-stage, short-term clinical outcome of hemiarthroplasty versus arthrodesis for, 848–851
 - hemiarthroplasty for, 591–593
 - influence on reamputation in patients with diabetes mellitus after partial hallux amputation, 1076–1080
 - late stage, short-term clinical outcomes after first metatarsal head resurfacing hemiarthroplasty for, 173–178
 - painful, autogenous capsular interpositional arthroplasty surgery for, 29–36
- Hallux valgus
 - corrections using osteodesis procedure, retrospective study of, 406–411
 - deformity
 - biomechanical evaluation of custom foot orthoses for, 852–855
 - change in first metatarsal length after proximal and distal chevron osteotomies for, 525–530
 - fixed, arthroscopic first metatarsophalangeal arthrodesis for repair of, 1127–1131
 - surgery
 - chevron versus Mitchell osteotomy in, 361–364
 - without lateral soft-tissue release, relationship between sesamoid complex and first metatarsal after, 1111–1115
- Hallux varus
 - idiopathic adult, by tendon transfer, correction of (case report), 1197–1201
- Hammer toe deformity
 - intramedullary fixation system for treatment of, 910–916
 - of lateral toes, correction with subtraction osteotomy of proximal phalanx neck, 601–606
- Headless screw fixation
 - and locking plate fixation, for talar neck fractures, mechanical comparison of, 905–909
- Heel lipoma
 - mimicking plantar fasciitis in a ballroom dancer (case report), 251–253
- Hematogenous calcaneal osteomyelitis
 - acute, of childhood, antibiotic treatment and surgery for, 840–843
- Hemiarthroplasty
 - versus arthrodesis
 - for end-stage hallux rigidus, short-term clinical outcome of, 848–851
 - for osteoarthritis of first metatarsophalangeal joint, 1085–1088
 - for hallux rigidus, 591–593
 - hemiarthroplasty resurfacing
 - first metatarsal, for late stage hallux rigidus, short-term clinical outcomes after, 173–178
- Heterotopic ossification prophylaxis
 - recurrent, after partial metatarsal amputation, radiation therapy for, 345–349
- High-heel shoe injury
 - in U.S. women, epidemiology of, 615–619

- Hindfoot
and ankle, without adjuvant therapy, surgical outcomes after excision of pigmented villonodular synovitis localized to, 160–163
- Hindfoot deformity
corrected with double versus triple arthrodesis, 424–427
- Hospital transport
at single level 1 trauma center, ankle fractures and modality of, 826–829
- Hyaline cartilage graft
juvenile, talocalcaneal joint middle facet coalition resection with interposition of (case report), 1178–1182
- Hydrocortisone injection
local, acute compartment syndrome of foot due to infection after (case report), 692–696
- Ilizarov fixation
external, without soft tissue release to correct severe, rigid equinus deformity, use of, 821–825
- Immobilization
risk, prevention, and diagnosis of venous thromboembolism disease in foot and ankle surgery and injuries requiring, 497–507
- Implant
flexible, and bioabsorbable anchor, use for deltoid rupture repair in bimalleolar equivalent Weber B ankle fractures, 513–516
- Implantation
primary, of Salto[®] mobile version and Salto Talaris[™] total ankle prostheses, incidence of revision after, a systematic review, 311–319
- Implant migration
after modular stem fixed-bearing total ankle replacement, evaluation of, 326–331
comparison of two generations of INBONE[®] Total Ankle Replacement System, 892–895
- INBONE[®] Total Ankle Replacement System
comparison of implant migration in two generations of, 892–895
- Infections
Achilles tendinitis after local injection of human placental extracts (case report), 1193–1196
after local hydrocortisone injection, acute compartment syndrome of foot due to (case report), 692–696
of first metatarsophalangeal joint with osteomyelitis in diabetic patients, conservative surgical treatment of, 536–540
late deep, after open reconstruction of Achilles tendon rupture, negative pressure wound therapy in management of, 2–6
- Ingrown toenails
comparison of Winograd procedure and wedge resection plus complete nail plate avulsion in treatment of, 395–398
nail braces as an alternative treatment for, 620–624
- Injury
requiring immobilization, and foot and ankle surgery, risk, prevention, and diagnosis of venous thromboembolism disease in, 497–507
- Insertional enthesophytes
safety of Achilles detachment and reattachment using standard midline approach to, 214–219
- Instability
of ankle
chronic, in young adults, epidemiology and risk factors for, 183–187
chronic, ligament reattachment technique for high-demand athletes with, 7–12
- Intercuneiform hook test
of medial intercuneiform instability after Lapidus arthrodesis, 464–472
- Intercuneiform instability
medial, after Lapidus arthrodesis, intercuneiform hook test of, 464–472
- Interdigital neuromas
multiple, retrospective study of 279 feet with 462, 320–322
- Internal fixation
for ankle arthrodesis, biomechanical comparison of, 188–191
limited, combined with external fixator, for tibial plafond fracture, postoperative complications between, 646–651
and open reduction
of intra-articular calcaneus fractures in children, minimally invasive (sinus tarsi) approach for (case report), 135–139
of posterior malleolar fractures with an associated lateral malleolar fracture, single oblique posterolateral approach for, 559–564
of simple displaced talar neck fractures, antegrade-retrograde opposing lag screws for, 23–28
- Internet
reliable source of information for patients seeking total ankle replacement, 378–381
- Interphalangeal joint
plantar hallux, ulceration complicated by osteomyelitis, unilobed rotational flap for (case report), 1166–1171
- Interpositional arthroplasty
with palmaris longus tendon graft for osteonecrosis of second metatarsal head (case report), 237–241
- Intramedullary fixation system
for treatment of hammer toe deformity, 910–916
- Intramedullary nail
fixation, retrograde, in comparison of patients with and without diabetes mellitus, tibiotalar arthrodesis using, 876–882
locked, and porous tantalum spacer, for post-traumatic global avascular necrosis of talus, tibio-calcaneal arthrodesis with (case report), 1172–1177
and porous metal spacer, revision subtalar joint fusion with (case report), 709–712
- Intrinsic muscle spasticity (intrinsic plus foot)
in adults with cerebral palsy, surgical treatment guidelines for digital deformity associated with (case report), 985–993
- Investigators' corners
Assessing Diagnostic Tests I: You Can't Be Too Sensitive, 519–520
Assessing Diagnostic Tests II: Grading on a Curve, 765–766
Causal Diagrams and Multivariate Analysis III: Confound It!, 145–147
Made to Measure: Designs Tailored to Your Study Needs, 1001–1002
Multiple Choice Answers: What to Do When You Have Too Many Questions, 285–286
Multiple Choice Answers: What to Do When You Have Too Many Questions, 285–286
Strange Data: When the Numbers Just Aren't Normal, 1216–1217
- Irreducible fracture-dislocation
of ankle, associated with interposition of tibialis posterior tendon in syndesmosis (case report), 962–966
- Jones fracture
fifth metatarsal, screw fixation diameter for, 227–229
- Lagged syndesmotic fixation
our clinical experience, 773–781
- Lapidus arthrodesis
comparison of crossed screw fixation with endosseous fixation device for, 1099–1105

- enhanced, crossed screw technique with middle cuneiform fixation reduces sagittal mobility in, 437–440
- intercuneiform hook test of medial intercuneiform instability after, 464–472
- and medial double arthrodesis, on arthrodiastasis of calcaneocuboid joint, comparative findings after, 412–416
- modified, with crossed screw fixation, 69–75
- Lapidus procedure
using proper application of rail external fixation, retrospective study of, 323–325
- Lateral impingement
after total ankle arthroplasty, partial transmalleolar approach for (case report), 233–236
- Lateral malleolar perforator flap
coverage of exposed bone of lateral malleolus with proximally based (case report), 730–733
- Lateral malleolus
pediatric, and physis, by free microvascular transfer of proximal fibular physis, reconstruction of (case report), 994–1000
with proximally based lateral malleolar perforator flap, coverage of exposed bone of (case report), 730–733
- Lateral stress dorsiflexion
view demonstrating clinical utility in midterm hallux limitus (case report), 739–746
- Ligament reattachment technique
for high-demand athletes with chronic ankle instability, 7–12
- Ligament release
in metatarsalgia and Morton's disease, comparison of outcomes between open neurectomy versus percutaneous metatarsal osteotomies and, 373–377
- Lisfranc joint complex
in radiographic and cadaveric comparison, anatomic parameters of, 883–887
- Locking plate fixation
and headless screw fixation, for talar neck fractures, mechanical comparison of, 905–909
- Locking plates
fibular, specialized, for lateral malleolar fractures, 1067–1071
lateral versus dual, for calcaneal fractures, biomechanical performance of, 830–835
surgical treatment of calcaneal fractures of Sanders type II and III by minimally invasive technique using locking plate, 76–81
wedge, and tricortical allograft wedges, in mid-calcaneal length after Evans calcaneal osteotomy, comparison of, 900–904
- Lower extremity
amputation risk factors associated with elevated ankle brachial indices and radiographic arterial calcification, 473–477
- Low-level laser therapy
at 635 nm, for treatment of chronic plantar fasciitis, 768–772
- Low-lying peroneus brevis muscle belly
in patients with peroneal tendon pathologic features, prevalence and role of, 872–875
- Lymphedema
hereditary, excision of elephantiasis nostras verrucosa lesions in patient with (case report), 747–750
- Magnetic resonance imaging
of asymptomatic lateral ankles to determine anatomic variants and incidental peroneal tendon pathologic features, 164–172
in symptomatic osseous abnormalities at posteromedial tubercle of the talus (case report), 978–984
- Malleolar fracture
of ankle, posterior, radiographic assessment of, 365–369
- lateral, specialized fibular locking plate for, 1067–1071
- posterior and lateral, single oblique posterolateral approach for open reduction and internal fixation of, 559–564
- stress, medial, in an adolescent athlete (case report), 242–246
- vertical shear medial, use of locked fibular plate for fixation of (case report), 1202–1205
- Melanoma
pigmented onychomatricoma a rare pigmented nail unit tumor presenting as longitudinal melanonychia potential for misdiagnosis of (case report), 723–725
- Melanonychia
longitudinal, potential for misdiagnosis as melanoma, pigmented onychomatricoma a rare pigmented nail unit tumor presenting as (case report), 723–725
- Metastasis
to midfoot, pulmonary pleomorphic carcinoma (case report), 483–486
- Metatarsal bones
amputation, partial, radiation therapy for recurrent heterotopic ossification prophylaxis after, 345–349
- fifth
base fracture, mechanism and treatment algorithms, anatomic evaluation of, 94–98
Jones fracture, screw fixation diameter for, 227–229
- first
head resurfacing hemiarthroplasty for late stage hallux rigidus, short-term clinical outcomes after, 173–178
length, after proximal and distal chevron osteotomies for hallux valgus deformity, change in, 525–530
and sesamoid complex, after hallux valgus surgery without lateral soft-tissue release, relationship between, 1111–1115
second, head, interpositional arthroplasty with palmaris longus tendon graft for osteonecrosis of (case report), 237–241
tuberculosis, multifocal bilateral (case report), 112–115
- Metatarsalgia
and Morton's disease, comparison of outcomes between open neurectomy versus percutaneous metatarsal osteotomies and ligament release in, 373–377
- Metatarsal osteotomy
percutaneous, versus open neurectomy, and ligament release, in metatarsalgia and Morton's disease, comparison of outcomes between, 373–377
- Metatarsal suture techniques
for bunion correction, complications of, a systematic review, 230–232
- Metatarsocuboid coalition
fifth, isolated (case report), 734–738
- Metatarsophalangeal joint
first
arthrodesis, arthroscopic, for repair of fixed hallux varus deformity, 1127–1131
arthrodesis versus hemiarthroplasty for osteoarthritis of, 1085–1088
long-term follow-up of cheilectomy for degenerative joint disease of, 1010–1020
with osteomyelitis in diabetic patients, conservative surgical treatment of infected ulceration of, 536–540
lesser, technique using interference fixation repair for plantar plate ligament disruption of, 508–512
- Metatarsus adductus
foot type, influence on plantar pressures during walking in adults using pedobarograph, 449–453

- Microtenotomy
using bipolar radiofrequency microtenotomy, intermediate-term results of partial plantar fascia release with, 179–182
- Mid-calcaneal length
after Evans calcaneal osteotomy, comparison of wedge locking plates and tricortical allograft wedges in, 900–904
- Midfoot
pulmonary pleomorphic carcinoma metastasis to (case report), 483–486
- Mineral density
and penetration strength, of subchondral bone plate of talar dome, 17–22
- Minimally invasive surgery
calcaneal osteotomy using Shannon burr endanger the neurovascular structures, 1062–1066
for displaced intra-articular calcaneal fractures, percutaneous arthroscopic calcaneal osteosynthesis a, 798–804
early treatment, of progressive foot and ankle deformity associated with Charcot-Marie-Tooth disease (case report), 701–708
plate osteosynthesis, of distal tibia and fibular fractures through single distal anterolateral incision, 1081–1084
(sinus tarsi), for open reduction and internal fixation of intra-articular calcaneus fractures in children (case report), 135–139
using locking plate, surgical treatment of calcaneal fractures of Sanders type II and III by, 76–81
- Mirror image twins
unicameral bone cyst in calcaneus of (case report), 754–757
- Misdiagnosis
of melanoma, pigmented onychomatricoma a rare pigmented nail unit tumor presenting as longitudinal melanonychia potential for (case report), 723–725
- Mitchell osteotomy
versus chevron, in hallux valgus surgery, 361–364
- Modular stem fixed-bearing
total ankle replacement, evaluating component migration after, 326–331
- Morton's disease
and metatarsalgia, comparison of outcomes between open neurectomy versus percutaneous metatarsal osteotomies and ligament release in, 373–377
- Morton's neuroma
compared with ultrasonography, diagnostic accuracy of clinical tests for, 549–553
- MRI. *see* Magnetic resonance imaging
- Multiple synostosis syndrome
surgical considerations for massive tarsal coalitions in (case report), 1162–1165
- Nail braces
as an alternative treatment for ingrown toenails, 620–624
- Nail plate avulsion
complete, wedge resection plus, and Winograd procedure, in treatment of ingrown toenails, comparison of, 395–398
- National Hospital Discharge Survey
and National Survey of Ambulatory Surgery, in foot and ankle arthrodesis, 1037–1041
- National Survey of Ambulatory Surgery
and National Hospital Discharge Survey, in foot and ankle arthrodesis, 1037–1041
- Navicular ossification center
in congenital talipes equinovarus, prognostic value of radiologic appearance of, 844–847
- Negative pressure wound therapy
and dermal regeneration template, treatment of foot degloving injury with aid of (case report), 1132–1135
in management of late deep infections after open reconstruction of Achilles tendon rupture, 2–6
- Neurectomy
open, versus percutaneous metatarsal osteotomies, and ligament release, in metatarsalgia and Morton's disease, comparison of outcomes between, 373–377
- Neurovascular structures
endanger, minimally invasive calcaneal osteotomy using Shannon burr, 1062–1066
- Obesity
effect on arthroscopic treatment of anterolateral impingement syndrome of ankle, 13–16
- One-stage tensile stress
surgical technique, and early postoperative rehabilitation, in treatment of neglected Achilles tendon rupture, outcome of, 153–159
- Open reduction
of an irreducible Hawkins type III fracture of talar neck, 677–682 and internal fixation
of intra-articular calcaneus fractures in children, minimally invasive (sinus tarsi) approach (case report), 135–139
versus limited internal fixation combined with external fixator, for tibial plafond fracture, postoperative complications between, 646–651
of posterior malleolar fractures with an associated lateral malleolar fracture, single oblique posterolateral approach for, 559–564
- Orthopedic immobilization
right lower limb, effects on braking function, 554–558
- Osseous abnormality
symptomatic, at posteromedial tubercle of the talus, magnetic resonance imaging in (case report), 978–984
- Ostectomy
calcaneal, lateral, and subtalar arthroscopic debridement, for residual pain after calcaneal fracture, simultaneous use of, 37–40
- Osteoarthritis
ankle
computed tomography atlas for grading, 207–213
current concepts in management of, a systematic review, 932–939
of first metatarsophalangeal joint, arthrodesis versus hemiarthroplasty for, 1085–1088
- Osteochondritis dissecans
of talus, outcomes are favorable after arthroscopic treatment of, 57–60
- Osteochondroma
congenital extraskeletal, of web space displacing lesser toes (case report), 726–729
- Osteodesis procedure
retrospective study of 63 hallux valgus corrections using, 406–411
- Osteogenic progenitor cell
concentration, effect on incidence of foot and ankle fusion, 888–891
- Osteoid osteoma
periarticular, of ankle, treated with arthroscopy, 89–93
- Osteomyelitis
after TightRope® fixation of ankle syndesmosis (case report), 130–134
in diabetic patients, conservative surgical treatment of infected ulceration of first metatarsophalangeal joint with, 536–540

- and severe soft tissue injury, treatment of calcaneal fracture with (case report), 973–977
- unilobed rotational flap for plantar hallux interphalangeal joint ulceration complicated by (case report), 1166–1171
- Osteonecrosis
- of second metatarsal head, interpositional arthroplasty with palmaris longus tendon graft for (case report), 237–241
- Osteotomy
- calcaneal
 - double, with percutaneous Steinmann pin fixation as part of treatment for flexible flatfoot deformity (case report), 478–482
 - Evans, comparison of wedge locking plates and tricortical allograft wedges in mid-calcaneal length after, 900–904
 - medial displacement, anatomy of sural nerve with an emphasis on incision for, 341–344
 - minimally invasive, using Shannon burr endanger the neurovascular structures, 1062–1066
 - percutaneous posterior (case report), 1188–1192
 - chevron
 - distal, role of fixation and postoperative regimens in long-term outcomes of, 356–360
 - versus Mitchell, in hallux valgus surgery, 361–364
 - proximal and distal, for hallux valgus deformity, change in first metatarsal length after, 525–530
 - metatarsal, percutaneous, versus open neurectomy, and ligament release, in metatarsalgia and Morton's disease, comparison of outcomes between, 373–377
 - scarf, restoring sesamoid position in, 1089–1092
 - subtraction, of proximal phalanx neck, correction of hammer toe deformity of lateral toes with, 601–606
 - union rates, in long bones of the foot, development of radiographic union score for determining, 793–797
- Outcomes
- after operative fusion of tarsal joints, a systematic review, 636–645
 - are favorable after arthroscopic treatment of osteochondritis dissecans of talus, 57–60
 - clinical, evidence-based rationale for ankle cartilage allograft replacement, a systematic review, 940–943
 - comparison between open neurectomy versus percutaneous metatarsal osteotomies and ligament release in metatarsalgia and Morton's disease, 373–377
 - of corticosteroid injection and extracorporeal shock wave therapy as primary therapeutic methods for acute plantar fasciitis, 1047–1052
 - of foot and ankle surgery in diabetic patients who have undergone solid organ transplantation, 577–581
 - functional, and patient satisfaction, after displaced intra-articular calcaneal fractures, 298–305
 - long-term, of distal chevron osteotomy, role of fixation and postoperative regimens in, 356–360
 - of one-stage tensile stress surgical technique and early postoperative rehabilitation in treatment of neglected Achilles tendon rupture, 153–159
 - short-term clinical
 - after first metatarsal head resurfacing hemiarthroplasty for late stage hallux rigidus, 173–178
 - of hemiarthroplasty versus arthrodesis for end-stage hallux rigidus, 848–851
 - surgical
 - after excision of pigmented villonodular synovitis localized to the ankle and hindfoot without adjuvant therapy, 160–163
 - and quality of life, in severe foot trauma and associated compartment syndrome after fasciotomy, 417–423
- Pain
- residual, after calcaneal fracture, simultaneous use of lateral calcaneal osteotomy and subtalar arthroscopic debridement for, 37–40
- Pain management
- for elective foot and ankle surgery, a systematic review, 625–635
- Palmaris longus tendon graft
- for osteonecrosis of second metatarsal head, interpositional arthroplasty with (case report), 237–241
- Patient satisfaction
- and functional outcome, after displaced intra-articular calcaneal fractures, 298–305
- Pedobarograph
- influence of metatarsus adductus foot type on plantar pressures during walking in adults using, 449–453
- Percutaneous arthroscopic calcaneal osteosynthesis
- a minimally invasive technique for displaced intra-articular calcaneal fractures, 798–804
- Perioperative period
- prophylactic antibiotic, use in clean elective foot surgery, ACFAS clinical consensus statement in, 273–279
- Peroneal retinacular injury
- superior, in calcaneal fractures, 458–463
- Peroneal spastic flatfoot
- recurrent, without bone coalitions, Botulinum toxin type A injection combined with cast immobilization for treatment of (case report), 697–700
- Peroneal tendon
- incidental, pathologic features, and anatomic variants, MRI of asymptomatic lateral ankles to determine, 164–172
 - pathologic features, prevalence and role of low-lying peroneus brevis muscle belly in patients with, 872–875
- Phalanx neck
- proximal, correction of hammer toe deformity of lateral toes with subtraction osteotomy of, 601–606
- Pigmented nail unit tumor
- presenting as longitudinal melanonychia, potential for misdiagnosis as melanoma, pigmented onychomatricoma a rare (case report), 723–725
- Pigmented onychomatricoma
- a rare pigmented nail unit tumor presenting as longitudinal melanonychia potential for misdiagnosis as melanoma (case report), 723–725
- Pigmented villonodular synovitis
- localized to the ankle and hindfoot without adjuvant therapy, surgical outcomes after excision of, 160–163
- Pilon fracture
- with ankle dislocation, pathogenesis and treatment strategies for, 815–820
- Pirani idiopathic clubfoot
- classifications of, clinical evaluation of, 582–585
- Placental extracts
- human, infectious Achilles tendinitis after local injection of (case report), 1193–1196
- Plain film bunion radiographs
- differences in eye tracking and gaze patterns between trainees and experts reading, 382–391
- Plantar calcaneonavicular (spring) ligament tear
- anatomic reconstruction technique for, 1124–1126

- Plantar fascia
 assessment of, plantar fascia coronal length a new parameter for, 445–448
 for corticosteroid injection, targeting of, 683–685
- Plantar fascia coronal length
 a new parameter for plantar fascia assessment, 445–448
- Plantar fascia release
 partial, with microtenotomy using bipolar radiofrequency microtenotomy, intermediate-term results of, 179–182
- Plantar fasciitis
 acute, treatment outcomes of corticosteroid injection and extracorporeal shock wave therapy as primary therapeutic methods for, 1047–1052
 in a ballroom dancer, heel lipoma mimicking (case report), 251–253
 chronic
 low-level laser therapy at 635 nm for treatment of, 768–772
 treatment with percutaneous latticed plantar fasciotomy, 856–859
 classification of calcaneal spurs and relationship with, 594–600
- Plantar fasciotomy
 percutaneous latticed, treatment of chronic plantar fasciitis with, 856–859
- Plantar flexor muscle–tendon complex
 in treatment of acute Achilles tendon rupture, early weightbearing on biomechanical properties of, 220–226
- Plantar plate
 versus dorsomedial plating, in first tarsometatarsal arthrodesis, anatomic evaluation of, 787–792
 ligament disruption, of lesser metatarsophalangeal joints, technique using interference fixation repair for, 508–512
- Plantar pressure
 during walking in adults using pedobarograph, influence of metatarsus adductus foot type on, 449–453
- Plate osteosynthesis
 minimally invasive, of distal tibia and fibular fractures through single distal anterolateral incision, 1081–1084
- Podiatric medicine
 and surgery, residency training experience in, 607–614
 and surgery, selecting residents in, 565–576
- Polyaxial angle–stable plate
 posterolateral, for tibiototalcalcaneal arthrodesis, early results of, 428–432
- Polymethylmethacrylate
 insertion, and longitudinal epiphyseal bracket by excision, at preossified disease stage, treatment of (case report), 1136–1140
- Ponseti technique
 in children presenting after one year of age, management of idiopathic clubfoot by (case report), 967–972
- Porous metal spacer
 and intramedullary nail, revision subtalar joint fusion with (case report), 709–712
- Porous tantalum spacer
 and locked intramedullary nail, for post-traumatic global avascular necrosis of talus, tibio-calcaneal arthrodesis with (case report), 1172–1177
- Postoperative period
 blood loss in calcaneal fractures, administration of tranexamic acid reduces, 1106–1110
 early postoperative, and one-stage tensile stress surgical technique, in treatment of neglected Achilles tendon rupture, outcome of, 153–159
- Postoperative regimens
 and fixation, role in long-term outcomes of distal chevron osteotomy, 356–360
- Preossified disease stage
 treatment of longitudinal epiphyseal bracket by excision and polymethylmethacrylate insertion at (case report), 1136–1140
- Pulmonary pleomorphic carcinoma
 metastasis to midfoot (case report), 483–486
- Quality of life
 and surgical outcomes, in severe foot trauma and associated compartment syndrome after fasciotomy, 417–423
- Radiation therapy
 for recurrent heterotopic ossification prophylaxis after partial metatarsal amputation, 345–349
- Radiographic union score
 for determining osteotomy union rates in long bones of the foot, development of, 793–797
- Radiography
 appearance of navicular ossification center in congenital talipes equinovarus, prognostic value of, 844–847
 assessment of posterior malleolar ankle fractures, 365–369 and cadaveric comparison, anatomic parameters of Lisfranc joint complex in, 883–887
 evaluation of Haglund's deformity, 338–340
 parameters, descriptive quantitative analysis of rearfoot alignment, 860–871
 weightbearing, facilitate functional treatment of ankle fractures of uncertain stability, 1042–1046
- Rail external fixation
 retrospective study of Lapidus procedure using proper application of, 323–325
- Range of motion
 preoperative and postoperative, comparison of two total ankle replacement systems in, 809–814
- Readmission
 unplanned, following foot and ankle surgery (editorial), 287
- Reamputation
 of hallux rigidus, influence in patients with diabetes mellitus after partial hallux amputation, 1076–1080
- Rearfoot
 alignment, radiographic parameters, descriptive quantitative analysis of, 860–871
- Recalcitrant hallux
 ulceration, evaluation of hallux interphalangeal joint arthroplasty compared with nonoperative treatment of, 541–548
- Reconstruction
 with Achilles tendon allograft and xenograft combination, chronic Achilles tendon rupture (case report), 1146–1150
 anatomic technique, for plantar calcaneonavicular (spring) ligament tear, 1124–1126
 of Charcot, comparison of complications for internal and external fixation for, 1072–1075
 of midfoot Charcot, posterior approach for medial column beam screw in, 433–436
 open, of Achilles tendon rupture, negative pressure wound therapy in management of late deep infections after, 2–6
 simultaneous bony and soft tissue, of mid- and hindfoot diabetic neuroarthropathy, use of definitive cement spacer for (case report), 120–125
 of traumatic forefoot, with free perforator flaps, 1025–1030
- Recreational sports activities
 after calcaneal fractures and subsequent subtalar joint arthrodesis, 1057–1061

- Rehabilitation
 early postoperative, and one-stage tensile stress surgical technique, in treatment of neglected Achilles tendon rupture, outcome of, 153–159
- Reoperations
 after tarsal coalition resection, 306–310
- Residency training
 experience in podiatric medicine and surgery, 607–614
- Residents
 in podiatric medicine and surgery, selection of, 565–576
- Rheumatoid arthritis
 of ankle joint, distraction arthroplasty with arthroscopic microfracture in patient with, 280–284
- Rigid equinovarus feet
 severe, midterm follow-up of talectomy for, 1093–1098
- Risk factors
 and epidemiology, for chronic ankle instability in young adults, 183–187
 lower extremity amputation associated with elevated ankle brachial indices and radiographic arterial calcification, 473–477
- Rupture
 of Achilles' tendon
 acute, early weightbearing on biomechanical properties of plantar flexor muscle–tendon complex in treatment of, 220–226
 chronic, reconstructed with Achilles tendon allograft and xenograft combination (case report), 1146–1150
 fluoroquinolone-mediated (case report), 494–496
 negative pressure wound therapy in management of late deep infections after open reconstruction of, 2–6
 neglected, outcome of one-stage tensile stress surgical technique and early postoperative rehabilitation in treatment of, 153–159
 using diffusion tensor imaging and tendon fiber tracking, assessment of postoperative tendon quality in patients with, 782–786
 deltoid, repair in bimalleolar equivalent Weber B ankle fractures, use of flexible implant and bioabsorbable anchor for, 513–516
 plantar plate ligament, of lesser metatarsophalangeal joints, technique using interference fixation repair for, 508–512
 of tibiofibular syndesmotic, bioabsorbable versus metallic screw fixation for, 657–662
- Sagittal mobility
 in enhanced Lapidus arthrodesis, crossed screw technique with middle cuneiform fixation reduces, 437–440
- Salto[®]
 mobile version, and Salto Talaris[™], total ankle prostheses, incidence of revision after primary implantation of, a systematic review, 311–319
- Salto Talaris[™]
 and Salto[®] mobile version, total ankle prostheses, incidence of revision after primary implantation of, a systematic review, 311–319
- Scarf osteotomy
 restoring sesamoid position in, 1089–1092
- Screw
 antegrade-retrograde opposing lag, for internal fixation of simple displaced talar neck fractures, 23–28
 fully threaded versus partially threaded, to prevent shear in cancellous bone fixation, 1021–1024
 insertion, cannulated, at different positions and directions, effects on subtalar joint stress distribution after, 920–926
 placement relative to calcaneal fracture constant fragment, 392–394
 plate versus lag, fixation of distal fibula fractures, biomechanical evaluation of, 896–899
- Screw fixation
 absorbable versus metallic, for distal tibiofibular syndesmosis injuries, 663–670
 bioabsorbable versus metallic, for tibiofibular syndesmotic ruptures, 657–662
 diameter for fifth metatarsal Jones fracture, 227–229
- Sesamoid
 restoring position of, in scarf osteotomy, 1089–1092
- Sesamoid complex
 and first metatarsal, after hallux valgus surgery without lateral soft-tissue release, relationship between, 1111–1115
- Shannon burr
 endanger the neurovascular structures, minimally invasive calcaneal osteotomy using, 1062–1066
- Shear strength
 in cancellous bone fixation, fully threaded versus partially threaded screws to prevent, 1021–1024
- Shoulder lesions
 of talus, role of demineralized allograft subchondral bone in treatment of (case report), 717–722
- Single oblique posterolateral approach
 for open reduction and internal fixation of posterior malleolar fractures with an associated lateral malleolar fracture, 559–564
- Single-stage lengthening
 versus callus distraction, with bone graft for treatment of brachymetatarsia, a systematic review, 927–931
- Sinus tarsi approach
 minimally invasive, for open reduction and internal fixation of intra-articular calcaneus fractures in children (case report), 135–139
- Soft tissue injury
 severe, and osteomyelitis, treatment of calcaneal fracture with (case report), 973–977
- Soft tissue release
 to correct severe, rigid equinus deformity, use of Ilizarov external fixation without, 821–825
 lateral, relationship between sesamoid complex and first metatarsal after hallux valgus surgery without, 1111–1115
- Solid organ transplantation
 outcomes of foot and ankle surgery in diabetic patients who have undergone, 577–581
- Spinal cage
 use for creating stable constructs in ankle and subtalar fusion (case report), 254–257
- Steinmann pin fixation
 percutaneous, as part of treatment for flexible flatfoot deformity, double calcaneal osteotomy with (case report), 478–482
- Stress ankle radiography
 manual, to predict deep deltoid ligament integrity in supination external rotation ankle fracture, 531–535
- Subchondral bone
 demineralized allograft, role in treatment of shoulder lesions of talus (case report), 717–722
- Subchondral bone plate
 of talar dome, mineral density and penetration strength of, 17–22
- Subtalar fusion
 use of spinal cage for creating stable constructs in (case report), 254–257
- Subtalar joint
 arthrodesis
 primary, for comminuted fractures of calcaneus, 61–65

- subsequent, and calcaneal fractures, recreational sports activities after, 1057–1061
- distraction arthrodesis to correct calcaneal valgus in pediatric patients with tarsal coalition (case report), 1151–1157
- fusion, with porous metal spacer and intramedullary nail, revision of (case report), 709–712
- posterior facet involvement, during intramedullary guidance of total ankle arthroplasty, quantification of, 805–808
- stress distribution after cannulated screw insertion at different positions and directions, effects on, 920–926
- Subtle Lisfranc subluxation
 - results of operative and nonoperative treatment of, 350–355
- Subtraction osteotomy
 - of proximal phalanx neck, correction of hammer toe deformity of lateral toes with, 601–606
- Superficial peroneal nerve
 - branching patterns on implications for ankle arthroscopy and for anterolateral surgical approaches to ankle, 332–337
- Supination external rotation
 - ankle fracture, manual stress ankle radiography to predict deep deltoid ligament integrity in, 531–535
- Sural nerve
 - with an emphasis on incision for medial displacement calcaneal osteotomy, anatomy of, 341–344
- Surgery
 - and antibiotic treatment, for acute hematogenous calcaneal osteomyelitis of childhood, 840–843
 - conservative treatment, of infected ulceration of first metatarsophalangeal joint with osteomyelitis in diabetic patients, 536–540
 - considerations for massive tarsal coalitions in multiple synostosis syndrome (case report), 1162–1165
 - foot and ankle
 - in diabetic patients who have undergone solid organ transplantation, outcomes of, 577–581
 - elective, pain management for, a systematic review, 625–635
 - modified suture bridge technique for application with bone anchors in, 944–948
 - forefoot, in elderly compared with younger patient populations, 586–590
 - of hallux valgus
 - chevron versus Mitchell osteotomy in, 361–364
 - without lateral soft-tissue release, relationship between sesamoid complex and first metatarsal after, 1111–1115
 - and podiatric medicine
 - residency training experience in, 607–614
 - selecting residents in, 565–576
 - of toe syndactyly, technique of dorsal transversely oriented transposition flap for web reconstruction in, 1119–1123
 - treatment guidelines for digital deformity associated with intrinsic muscle spasticity (intrinsic plus foot) in adults with cerebral palsy (case report), 985–993
 - treatment in giant cell tumors of foot and ankle bones, high recurrence rates after (case report), 1141–1145
- Suture
 - surgical, of Achilles tendon, gait analysis before and after (case report), 247–250
- Suture bridge technique
 - modified, for application with bone anchors in foot and ankle surgery, 944–948
- Suture button technique
 - assessment of anatomic risk during syndesmotomic stabilization with, 917–919
- Syndesmosis
 - irreducible fracture-dislocation of ankle associated with interposition of tibialis posterior tendon in (case report), 962–966
 - tibiofibular, distal, arthroscopic arthrodesis of, 953–957
- Syndesmotomic stabilization
 - with suture button technique, assessment of anatomic risk during, 917–919
- Talar dome
 - mineral density and penetration strength of subchondral bone plate of, 17–22
- Talectomy
 - for severe rigid equinovarus feet, midterm follow-up of, 1093–1098
- Talocalcaneal joint middle facet coalition
 - resection with interposition of juvenile hyaline cartilage graft (case report), 1178–1182
- Talofibular ligament
 - anterior, use of an all-suture anchor for re-creation of (case report), 126–129
- Talus
 - fractures of, in children after go-karting accidents (case report), 1206–1212
 - giant cell tumor of (case report), 958–961
 - neck fractures of
 - irreducible Hawkins type III, open reduction of, 677–682
 - mechanical comparison of headless screw fixation and locking plate fixation for, 905–909
 - neck simple displaced fractures, antegrade-retrograde opposing lag screws for internal fixation of, 23–28
 - osteocondritis dissecans of, outcomes are favorable after arthroscopic treatment of, 57–60
 - posteromedial tubercle, magnetic resonance imaging in symptomatic osseous abnormalities at (case report), 978–984
 - role of demineralized allograft subchondral bone in treatment of shoulder lesions of (case report), 717–722
 - tibiocalcaneal arthrodesis with porous tantalum spacer and locked intramedullary nail for post-traumatic global avascular necrosis of (case report), 1172–1177
- Tarsal coalition
 - massive, in multiple synostosis syndrome, surgical considerations for (case report), 1162–1165
 - subtalar joint distraction arthrodesis to correct calcaneal valgus in pediatric patients with (case report), 1151–1157
- Tarsal coalition resection
 - reoperations after, 306–310
- Tarsal joints
 - outcome after operative fusion of, a systematic review, 636–645
- Tarsal navicular
 - relationship of dorsalis pedis artery to, 66–68
- Tarsometatarsal joint
 - arthrodesis of, first, anatomic evaluation of dorsomedial versus plantar plating in, 787–792
- Tendinopathy
 - chronic, simple technique for repair of, 143–144
- Tendon fiber tracking

- and diffusion tensor imaging, assessment of postoperative tendon quality in patients with Achilles tendon rupture using, 782–786
- Tendon quality
postoperative, in patients with Achilles tendon rupture using diffusion tensor imaging and tendon fiber tracking, assessment of, 782–786
- Tendon transfer
correction of idiopathic adult hallux varus by (case report), 1197–1201
- Tendon weaving technique
allograft, augmented repair of acute Achilles tendon rupture using, 1004–1009
- Tendoscopy
tenosynovial osteochondromatosis of flexor hallucis longus tendon treated by (case report), 758–764
- Tenosynovial osteochondromatosis
of flexor hallucis longus tendon treated by tendoscopy (case report), 758–764
- Tenosynovitis
Blastomyces, of foot and ankle (case report), 1183–1187
- Thompson test
intraoperative value of, 99–101
- Tibia
fractures of
distal, elastic bandage traction technique for reduction of, 949–952
distal, through single distal anterolateral incision, minimally invasive plate osteosynthesis of, 1081–1084
plafond fractures, Arbeitsgemeinschaft für Osteosynthesefragen (AO) 43-C, hybrid external fixation for, 1031–1036
- Tibialis anterior subluxation
irreducible ankle fracture-dislocation due to (case report), 268–272
- Tibialis anterior tendon lengthening
for adjunctive treatment of plantar lateral column diabetic foot ulcers, 686–691
- Tibial plafond
fracture of, postoperative complications between open reduction and internal fixation versus limited internal fixation combined with external fixator for, 646–651
- Tibial sesamoidectomy
iatrogenic hallux abducto valgus secondary to, 82–88
- Tibial tendon
posterior, in syndesmosis, irreducible fracture-dislocation of ankle associated with interposition of (case report), 962–966
- Tibial tendon entrapment
posterior, within an intact ankle mortise (case report), 116–119
- Tibiocalcaneal arthrodesis
with porous tantalum spacer and locked intramedullary nail for post-traumatic global avascular necrosis of talus (case report), 1172–1177
- Tibiofibular joint
syndesmotic rupture, bioabsorbable versus metallic screw fixation for, 657–662
- Tibiofibular syndesmosis
distal
arthroscopic arthrodesis of, 953–957
injury, absorbable screws versus metallic screws for, 663–670
- Tibiotalar calcaneal arthrodesis
early results of posterolateral polyaxial angle-stable plate for, 428–432
using retrograde intramedullary nail fixation in comparison of patients with and without diabetes mellitus, 876–882
- Toe
lateral, hammer toe deformity of, correction with subtraction osteotomy of proximal phalanx neck, 601–606
lesser, congenital extraskeletal osteochondroma of web space displacing (case report), 726–729
- Toe syndactyly
surgery, technique of dorsal transversely oriented transposition flap for web reconstruction in, 1119–1123
- Total ankle arthroplasty
Internet a reliable source of information for patients seeking, 378–381
partial transmalleolar approach for lateral impingement after (case report), 233–236
in preoperative and postoperative range of motion, comparison of, 809–814
quantification of subtalar posterior facet involvement during intramedullary guidance of, 805–808
registry data trends of using, 1213–1142(Letter/Reply)
- Total ankle prosthesis
Salto[®] mobile version and Salto Talaris[™], incidence of revision after primary implantation of, a systematic review, 311–319
- Total ankle replacement
evaluating component migration after modular stem fixed-bearing, 326–331
- Trabecular metal spacer
and anterior fusion plate, ankle fusion with (case report), 490–493
- Tranexamic acid
reduces postoperative blood loss in calcaneal fractures, administration of, 1106–1110
- Transmalleolar approach
partial, for lateral impingement after total ankle arthroplasty (case report), 233–236
- Trauma center
single level 1, ankle fractures and modality of hospital transport at, 826–829
- Tuberculosis
of the cuboid (case report), 713–716
multifocal bilateral metatarsal (case report), 112–115
- Ulceration
of first metatarsophalangeal joint with osteomyelitis in diabetic patients, conservative surgical treatment of, 536–540
of plantar hallux interphalangeal joint, complicated by osteomyelitis, unilobed rotational flap for (case report), 1166–1171
of plantar lateral column diabetic foot, tibialis anterior tendon lengthening for adjunctive treatment of, 686–691
recalcitrant hallux, evaluation of hallux interphalangeal joint arthroplasty compared with nonoperative treatment of, 541–548
- Ultrasonography
diagnostic accuracy of clinical tests for Morton's neuroma compared with, 549–553
- Venous thromboembolism disease
in foot and ankle surgery and injuries requiring immobilization, risk, prevention, and diagnosis of, 497–507
- Walking
adults, influence of metatarsus adductus foot type on plantar pressures using pedobarograph in, 449–453
- Web reconstruction
in toe syndactyly surgery, technique of dorsal transversely oriented transposition flap for, 1119–1123

- Web space
displacing lesser toes, congenital extraskeletal osteochondroma of (case report), 726–729
- Wedge resection
plus complete nail plate avulsion, and Winograd procedure, in treatment of ingrown toenails, comparison of, 395–398
- Weightbearing
early, on biomechanical properties of plantar flexor muscle–tendon complex in treatment of acute Achilles tendon rupture, 220–226
forefoot defects with digital artery flaps, reconstruction of, 41–45
radiographs facilitate functional treatment of ankle fractures of uncertain stability, 1042–1046
- Winograd procedure
and wedge resection plus complete nail plate avulsion, in treatment of ingrown toenails, comparison of, 395–398
- Xenograft
Achilles tendon, and allograft combination, chronic Achilles tendon rupture reconstructed with (case report), 1146–1150