



Index to Volume 34

Keywords

- Abdominal aortic aneurysm, acute, multidisciplinary team for, 1519–1523
- Abdominal blunt trauma, usefulness of resuscitative endovascular balloon occlusion of aorta in detecting source of a hemorrhage due to, a case report, 2057.e1–2057.e3
- Abdominal compressions, interposed, combination with chest compressions, in swine model of ventricular fibrillation, 968–974
- Abdominal pain
acute
gadolinium-based contrast agent anaphylaxis induced, a case report, 1737.e1–1737.e2
idiopathic ovarian vein thrombosis, rare cause of, a case report, 935.e1–935.e2
discharged from ED, variation in advanced imaging for pediatric patients with, 2320–2325
in ED patients, aortoduodenal fistula as cause of, a case report, 2468.e1–2468.e4
in pregnancy, omental infarction, unusual cause of, a case report, 1184.e3–1184.e4
self-assessment of hunger among ED patients with, lack of association with disease severity, 104–106
who are at low risk for acute appendicitis, prospective validation of biomarker panel to identify pediatric ED patients with, 1373–1382
- Abdominal vascular graft, complications, emergency medicine approach to, 2014–2017
- Above knee amputation, bilateral, uncommon complication of hyperosmolar hyperglycemic state, a case report, 341.e1–341.e2
- Abscess
epidural, *Streptococcus pneumoniae*, rare pathogen in, a case report, 1919.e3–1919.e4
peritonsillar, using bedside emergency ultrasound to change diagnosis of, a case report, 1186.e5–1186.e7
ventriculoseptal, and intracardiac fistulization, infected heart, a case report, 678.e1–678.e3
- Acidemia, severe, on arrival not predictive of neurologic outcomes in post-cardiac arrest patients, 425–428
- Acidosis, profound, and rhabdomyolysis, due to dietary supplement use, survival after, a case report, 2259.e1–2259.e3
- ACS, Acute coronary syndrome (ACS). *see* Acute coronary syndrome (ACS)
- Activated charcoal
may not be necessary in all oral overdoses of medication (letter), 319–321
use, prehospital, in antipsychotic overdose (letter), 316–317
- Acuity of illness, socioeconomic status, and prehospital delay, on presentation (letter), 1712–1714
- Acupuncture, vs intravenous morphine, in management of acute pain in ED, 2112–2116
- Acute abdomen, spontaneous perforation of pyometra presenting as, rare condition with considerable mortality, a case report, 761.e3–761.e5
- Acute compartment syndrome, orthopedic literature 2015, 1663–1667
- Acute coronary syndrome (ACS)
apical hypertrophic cardiomyopathy mimicking, a case report, 1322.e1–1322.e2
atypical, late diagnosis of Wellens syndrome in patient presenting with, case report, 338.e3–338.e5
electrocardiogram in, high-risk electrocardiographic presentations lacking anatomically oriented ST-segment elevation, 611–617
Kounis hypersensitivity-associated, and akotsubo cardiomyopathy, myocardial injury through serum troponin I and echocardiography in anaphylaxis (letter), 650–651
meaning of elevated troponin I levels, not always, 145–148
- Acute generalized exanthematous pustulosis, uncommon cause of fever and rash, case report, 681.e1–681.e2
- Acute hemorrhagic edema of infancy, and common mimics, a case report, 936.e3–936.e6
- Acute mesenteric ischemia
baseline disease is more important predictor of intestinal necrosis than CT findings in patients with, 2261–2265
superior mesenteric artery dissection does not necessarily mean (letter), 2457
- Acute myeloid leukemia, in ED, nontraumatic splenic rupture as initial manifestation of, a case report, 1731.e1–1731.e3
- Adolescent, Wallenberg's lateral medullary syndrome in, a case report, 2254.e1–2254.e2
- Adrenomedullin, bioactive, plasma, as prognostic biomarker in acute heart failure, 257–262
- Adropin, levels of, and target organ damage, secondary to high blood pressure in ED, 2061–2064
- Adult
in ED, *Plesiomonas shigelloides* meningitis in, a case report, 1329.e1–1329.e2
ED patients, use of inhaled nitrous oxide for analgesia in, 269–273
with in-hospital cardiac arrest, meta-analysis of outcomes of 2005 and 2010 cardiopulmonary resuscitation guidelines for, 1133–1139
older
in ED, effect of surgical consult in treatment of abdominal pain in, 1524–1527
ED as primary source of hospital admission for, 943–947
underserved, cognitive health and risk of ED revisit in, 1973–1976
optimal wrist positioning for radial arterial cannulation in, 2372–2378
and pediatric academic EDs, impact of scribes on patient throughput in, 1982–1985
presenting to ED with meningitis, validation of bacterial meningitis score in, 1265–1267

- prognostic factors of *Streptococcus pneumoniae* infection in, 202–206
 under resuscitation performed by paramedics, comparison of NIO and EZ-IO intraosseous access devices in, randomized crossover manikin trial (letter), 1166–1167
 retropharyngeal air and pneumomediastinum, rare complication of influenza A and asthma in, a case report, 338.e1–338.e2
 swine model, of ventricular fibrillation, effects of proximal and distal routes of intraosseous epinephrine administration on short-term resuscitative outcome measures in, randomized controlled study, 49–53
 undergoing venipuncture, prospective, randomized, double-blind controlled trial comparing vapocoolant spray vs placebo spray in, 798–804
 unilateral absence of pulmonary artery concomitant with chronic pulmonary thromboembolism in, a case report, 1916.e3–1916.e5
 working, US private sector, paid sick leave is associated with fewer ED visits among, 784–789
 young
 with acute scrotal pain, clinical findings in, 1931–1933
 identifying optimal hand placement site for chest compression by measuring hand width and sternal length in, 407–411
- Adverse events
 and errors, in ED, medical record reviewers can reliably identify, 1043–1048
 little known, of lidocaine, Hoigne syndrome, a case report, 679.e3–679.e4
 in pulmonary emboli patients diagnosed with contrast CT, relation of contrast nephropathy to, 1247–1250
 in pulmonary embolism patients
 contrast-induced nephropathy related to (letter), 2042–2043
 contrast-induced nephropathy related to (reply), 2043–2044
 unusual, of acute urinary retention, STEMI, a case report, 347.e1–347.e2
- Age
 effects on case fatality rates of injury patients by mechanism, 515–520
 and gender group, bystander cardiopulmonary resuscitation training experience and self-efficacy of, 1331–1337
 -related differences, diseases associated with electrolyte imbalance in ED, 1923–1926
- Agitation, patients, in ED, chemical agents for sedation of, 2426–2431
- Air pollutants, influence on appearance of acute myocardial infarction in region with humid continental climate (letter), 1679–1682
- Air pollution, and weather conditions, effects on incidence of acute MI, 449–454
- Airtraq
 ETVIEW SL, and Macintosh laryngoscopes, comparison for face-to-face tracheal intubation, randomized crossover manikin trial (letter), 1893–1894
 and Miller laryngoscopes, for intubation in patients with inline manual neck stabilization, comparison of (letter), 1721
- AirTraq Avant[®], vs Macintosh laryngoscope, for intubation during simulated resuscitation by novice physicians, comparison of (letter), 1708–1709
- Airway, circulated and non-circulated, visual recognition of anatomical structures in, 1236–1240
- Airway management
 advanced, prehospital, role on outcomes for out-of-hospital cardiac arrest patients, 2101–2106
 comparison of video laryngoscopy and Macintosh direct laryngoscopy for, 1886
 difficult
 comparison of emergency physician and anesthesiologist attitudes toward (letter), 754–755
 in severely injured patients, I-gel as alternative airway tool for, a case report, 340.e1–340.e4
 for dyspnea with mediastinal mass, emergent radiation therapy as definite, a case report, 2258.e5–2258.e6
- Airway obstruction
 complete, assessment of LifeVac on human cadaver with (letter), 1673–1674
 upper, in pediatric patients, paramedic assessment and treatment of, exploratory analysis by Children's Safety Initiative-Emergency Medical Services, 599–601
- Albumin
 to fibrinogen ratio, and severity of coronary artery disease in patients with STEMI, relationship between, 1037–1042
 ischemia-modified, in cardiovascular disease (letter), 1699
 serum level, is associated with recurrence of acute ischemic stroke, 1812–1816
- Alcohol, blood levels, and plasma lactate (letter), 1725–1726
- Alcohol consumption, acute, and discharge against medical advice, of injured patients in ED, association between, 464–468
- Alcohol ingestion, accidental, in 7-month-old baby, fulminant hepatic failure requiring liver transplantation due to, a case report, 2466.e1–2466.e2
- Alcohol intake, quantitative analysis of high plasma lactate concentration in ED patients after, 825–829
- Alcohol intoxication, role in psychiatrists' recension of emergency physicians' involuntary admissions (letter), 2226–2227
- Alvarado score, C-reactive protein, and ultrasound, predictive value in acute appendicitis, 189–192
- Alzheimer disease, in-hospital mortality due to pulmonary embolism in patients with, sex effect (letter), 665–666
- Ambulatory care, -sensitive condition conditions, association of mental health disorders and Medicaid with ED admissions for, 820–824
- Amylase
 and lipase
 normal, acute pancreatitis with, a case report, 940.e5–940.e7, 2254.e3–2254.e6
- Analgesia
 in adult ED patients, use of inhaled nitrous oxide for, 269–273
 for adult patients with femoral shaft fractures after injury, intravenous morphine titration as rapid and efficient, 2107–2111
 calcaneal fracture, ED ultrasound-guided posterior tibial nerve blocks for, a case report, 1183.e1–1183.e3
 in patients with painful conditions, ED triage pain protocol reduces time to receiving, 2362–2366
 prescription, in acute appendicitis, effects of pain severity and CT imaging on, 36–39
 and procedural sedation, ketamine-propofol combination (ketofol) vs propofol for, 558–569
- Anaphylactic shock, biphasic, contrast-induced, combined use of ECMO and hemodialysis in case of, a case report, 1919.e1–1919.e2
- Anaphylaxis
 appropriateness of epinephrine use in ED patients with, 174–179
 evaluation of myocardial injury through serum troponin I and echocardiography in, 140–144
 firefighters able to perform intraosseous access and start fluid resuscitation in patient with (letter), 1707–1708
 myocardial injury through serum troponin I and echocardiography in, Takotsubo cardiomyopathy and Kounis hypersensitivity-associated ACS (letter), 650–651
 -triggered Takotsubo syndrome, prevalence of (letter), 748
- ANCA. *see* Anticytoplasm neutrophil antibody (ANCA)
- Anesthesia, hand, in healthy volunteers, forearm ultrasound-guided nerve blocks vs landmark-based wrist blocks for, 730–734
- Anesthesiologist, comparing with emergency physician, attitudes toward difficult airway management (letter), 754–755
- Angiography
 coronary, emergency, in out-of-hospital cardiac arrest patients without STEMI, a case report, 118.e1–118.e3
- CT
 in acute stroke (revisiting the 4Ps of imaging), 282–287

- of chest, saddle embolism diagnosed by point-of-care transthoracic echocardiography before, a case report, 2467.e1–2467.e2
- coronary, diagnostic performance of smartphone reading in patients with acute chest pain at ED, 1794–1798
- epipericardial fat necrosis clinically mimicking pulmonary embolism, a case report, 2056.e5–2056.e6
- pulmonary, CT, risk of pulmonary embolism after prior negative, 1968–1972
- Anion gap**
- metabolic acidosis
 - normal, in salicylate overdose (letter), 2457–2458
 - salicylate toxicity in absence of, a case report, 1328.e1–1328.e3
- Ankle fracture, postsplinting x-rays of, 1625–1626
- Ankle injury, orthopedic literature 2015, 1663–1667
- Antibiotics**
- administration in patients with sepsis in ED, pharmacist impact on time to, 2117–2121
 - dosing adjustment, impact of emergency medicine pharmacist on, 980–984
 - in ED, procalcitonin-guided therapy for initiation of, 1286–1293
 - empiric therapy, in ED, health care–associated acute pyelonephritis associated with inappropriate, 1415–1420
 - failure in cellulitis trials, 1645–1652
 - and fluids, in ED, triage sepsis alert and sepsis protocol lower times to, 1–9
 - intravenous, in ED, impact of prospective verification of, 2392–2396
 - outpatient, failure among patients hospitalized for acute bacterial skin infections, 957–962
 - provider selection aids appropriate treatment for outpatient UTI, ED pharmacist monitoring of, 1600–1603
 - retrieval, and discharge instruction delivery/modality (letter), 1910
 - retrieval, seventy-two-hour, from ED, randomized controlled trial of discharge instructional modality, 999–1005
 - seventy-two-hour retrieval, from ED, reporting the strength of association (letter), 1909–1910
 - treating appendicitis with, 609–610
- Anticoagulants**
- oral
 - discontinued in patient with cardiac valve prosthesis after traumatic intracranial hemorrhage, a case report, 1736.e5–1736.e7
 - non-vitamin K antagonist, efficacy on Lemierre syndrome, a case report, 2059.e3–2059.e5
 - reversal, emergent, pediatric, using 4-factor PCC, a case report, 1182.e1–1182.e2
- Anticytoplasm neutrophil antibody (ANCA), –positive vasculitis with diffuse alveolar hemorrhage related to spider bite, a case report, 339.e3–339.e4
- Antioxidant, and oxidant levels, investigation in patients with acute stroke in emergency service, 2379–2383
- Antivenom, F(ab')₂, new, for treatment of crotaline envenomation in children, 2003–2006
- Anuria, and acute kidney injury, an uncommon case of bilateral synchronous ureteric calculi, a case report, 2256.e1–2256.e2
- Anxiety**
- in infants and young children's
 - effects of nursery songs, reporting mistake of effect measure of association (letter), 663
 - before and during head CT, randomized single-blinded clinical trial on effects of nursery songs for (letter), 663
- Aortic coarctation, complicated with spinal epidural hemorrhage, paraplegia caused by aortic coarctation, a case report, 680.e1–680.e2
- Aortic dissection**
- acute
 - association with troponin leak, coronary anatomy, and prognosis, electrocardiography changes in, 1431–1436
 - spontaneous herniation mimicking, a case report, 2053.e1–2053.e2
 - ascending, emergency physician focused cardiac ultrasound improves diagnosis of, 486–492
 - manifesting as dysphagia and hoarseness, Ortner's syndrome, a case report, 1185.e1–1185.e3
 - type B, acute, after endovascular therapy, hypoalbuminemia predicts clinical outcome in patients with, 1369–1372
- Aortic graft, in hollow viscus, a case report, 678.e5–678.e6
- Aortic syndrome, acute, missed, limiting scan range of cardiac CT and chance of, 2007–2010
- Aortoduodenal fistula, as cause of abdominal pain in ED patients, a case report, 2468.e1–2468.e4
- Apical hypertrophic cardiomyopathy, mimicking acute coronary syndrome, a case report, 1322.e1–1322.e2
- Appendectomy, effect of time delays for, as observed on CT in patients with noncomplicated appendicitis, 167–169
- Appendicitis**
- acute
 - Bayesian comparative model of CT scan and ultrasonography in assessment of, 2070–2073
 - in children, evaluation of proadrenomedullin as diagnostic or prognostic biomarker of, 2298–2305
 - effects of pain severity and CT imaging on analgesia prescription in, 36–39
 - predictive value of C-reactive protein, ultrasound and Alvarado score in, 189–192
 - prospective validation of biomarker panel to identify pediatric ED patients with abdominal pain who are at low risk for, 1373–1382
 - spot urinary 5-hydroxyindoleacetic acid is not ideal diagnostic test for, 1750–1753
 - addition of biomarker panel to a clinical score to identify patients at low risk for, 2266–2271
 - barium, a case report, 1732.e1–1732.e2
 - noncomplicated, effect of time delays for appendectomy as observed on CT in patients with, 167–169
 - perforated
 - with concurrent ovarian hyperstimulation syndrome, following induction ovulation with HMG and HCG, a case report, 2051.e5–2051.e6
 - in pediatric patient, omental torsion mimicking, emergency bedside sonography, a case report, 684.e3–684.e4
 - stump, diagnostic challenges with, a case report, 1188.e–1188.e2
 - suspected, usefulness of new and traditional serum biomarkers in children with, 871–876
 - treating with antibiotics, 609–610
- Artemisinin, protects mice against burn sepsis through inhibiting NLRP3 inflammasome activation, 772–777
- Arterial base deficit, is still useful prognostic marker in trauma, 626–635
- Arterial occlusion, lower extremity, bilateral, acute, without risk factors or embolic source, a case report, 762.e3–762.e4
- Arthroscopy, of irreducible posterolateral knee dislocation, a case report, 2256.e3–2256.e4
- Asphyxia, emergency endotracheal intubation under fluoroscopy guidance for patients with, 2177–2181
- Asthma**
- attacks, moderate to severe, effect of nebulized magnesium sulfate in treatment of, randomized clinical trial, 883–886
 - bronchial, beneficial action of magnesium sulfate in (letter), 1693–1694
 - care in ED, characteristics of heroin-dependent patients seeking, 895–898
 - and influenza A, in adult, retropharyngeal air and pneumomediastinum, rare complication of, a case report, 338.e1–338.e2
- Atrial fibrillation**
- "false," unexpected mismatch between chamber volume and electrocardiogram signal in, a case report, 1740.e1–1740.e2

- hospital-level variation and predictors of admission after ED visits for, 2094–2100
- rapid ventricular response, or tachycardia, three faces of (letter), 931–932
- rare complication of scorpion venom, a case report, 937.e3–937.e4
- recent-onset, safety and efficacy of pharmacological cardioversion of, 1486–1490
- risk alleles, and response, association to acute rate control therapy, 735–740
- stroke prevention for patients with (letter), 1685–1687
- Atrial myxoma, complications of, a case report, 2465.e1–2465.e2
- Atrial septal defect, can mimic pulmonary thromboembolism after major surgery, a case report, 121.e3–121.e6
- Atrial standstill, and thromboembolic complications (letter), 932
- Autophagy, improved neurologic outcome after cardiopulmonary resuscitation in rats, activation of, 1511–1518
- Avalanche victims, potassium as a potential predictive biomarker of brain hypoxia in (letter), 1315–1316
- Avascular necrosis, of lunate bone, Kienbock disease, a case report, 1185.e5–1185.e6
- Axillary artery laceration, after anterior shoulder dislocation presenting as hypovolemic shock, a case report, 1739.e1–1739.e4
- Azithromycin, Stevens-Johnson syndrome after treatment with, uncommon culprit, a case report, 676.e1–676.e3
- Backboard, effect on compression depth during cardiac arrest in ED, simulation study, 274–277
- Bacteremia, in female patients with acute pyelonephritis, diagnostic performance of initial serum lactate for predicting, 1359–1363
- Bacterial meningitis
- acute lethargy in young woman due to latent disseminated cancer mimicking, diagnostic pitfall, a case report, 2050.e5–2050.e7
 - and CSF/blood glucose ratio, without pleocytosis in immunocompromised host (letter), 922
 - without pleocytosis (letter), 2443
- Bacterial meningitis score, in adults presenting to ED with meningitis, validation of, 1265–1267
- Bacterial strains
- in blood cultures
 - of septic patients, procalcitonin levels predict identification of (letter), 2453–2454
 - of septic patients, procalcitonin levels predict to identify, 2150–2153
- Barium, appendicitis, a case report, 1732.e1–1732.e2
- Basic life support (BLS), standard, and feedback CPR devices, by nurses, randomized trial of chest compressions effectiveness comparing, 381–385
- Bed width, effect on quality of compressions in simulated resuscitation, randomized crossover manikin study, 2272–2276
- Behavioral emergency, emergency medical services annotated literature in review 2016, 2193–2199
- Behcet disease, spontaneous bilateral pneumothorax unusual complication of, a case report, 1320.e1–1320.e2
- Beta agonist–induced lactic acidosis, evidence-based approach to critical question, 666–668
- Beta-blockers, long-term use in chronic heart failure (letter), 931
- Bevacizumab, treatment, presenting to ED as chest pain and headache, posterior reversible encephalopathy syndrome after, a case report, 1916.e1–1916.e2
- Bibliometric analysis, of top 100 most-cited clinical studies on ultrasound in ED, 1210–1216
- Bicycle crash
- in California
 - bicyclists found at fault for (letter), 1699–1701
 - trends in (letter), 1318–1319
- Bicyclists, found at fault for bicycle crashes in California (letter), 1699–1701
- Biochemistry analyzer
- and blood gas, in patients with moderate to severe hyperkalemia, agreement of serum potassium measured by, 794–797
 - and blood gas analyzer, in patients with hyponatremia, eunatremia, and hypernatremia, comparison of serum sodium levels measured by, 1473–1479
- Bioelectrical impedance analysis, NT-proBNP, IVC ultrasound, for diagnosis of acute decompensated CHF, 1817–1822
- Biohazard suit, blind intubation through supraglottic airway laryngopharyngeal tube with (letter), 2040
- Biological test
- routine
 - in self-poisoning patients, results from observational prospective multicenter study (letter), 2036–2037
 - in self-poisoning patients, results from observational prospective multicenter study (reply), 2037
- Biomarker
- cardiac, elevated, are not associated with mortality in low-risk cardiac patients with severe sepsis (letter), 1899–1900
 - diagnostic or prognostic, of acute appendicitis in children, evaluation of proadrenomedullin as, 2298–2305
 - inflammatory, predicting prognosis in patients with acute dyspnea, 370–374
 - panel, addition to clinical score to identify patients at low risk for appendicitis, 2266–2271
 - panel to identify pediatric ED patients with abdominal pain who are at low risk for acute appendicitis, prospective validation of, 1373–1382
 - prognostic, in acute heart failure, plasma bioactive adrenomedullin as, 257–262
 - in sepsis (letter), 924–925
 - serum, new and traditional, usefulness in children with suspected appendicitis, 871–876
- Bispectral index, monitoring in subarachnoid hemorrhage–associated out-of-hospital cardiac arrest, a case report, 934.e1–934.e3
- Bite, spider, ANCA–positive vasculitis with diffuse alveolar hemorrhage related to, a case report, 339.e3–339.e4
- Bland-Altman analysis, comparison of 2 available methods for measuring intracompartmental pressure, 1765–1771
- Bleeding, sites, in elderly trauma patients who required massive transfusion, comparison with younger patients, 123–127
- Blind intubation, using intubating supraglottic airway devices, effect of head position on success rate of, 1193–1197
- B-line score, rapid, clinical application with lung ultrasonography in differentiating between pulmonary infection and pulmonary infection with acute left ventricular heart failure, 278–281
- Blood cultures
- of septic patients, procalcitonin levels predict identification of bacterial strains in (letter), 2453–2454
 - of septic patients, procalcitonin levels predict to identify bacterial strains in, 2150–2153
- Blood gas analysis, radial pseudoaneurysm after puncture for, a case report, 757.e1–757.e2
- Blood gas analyzer
- and biochemistry, in patients with moderate to severe hyperkalemia, agreement of serum potassium measured by, 794–797
 - and biochemistry autoanalyzer, in patients with hyponatremia, eunatremia, and hypernatremia, comparison of serum sodium levels measured by, 1473–1479
 - in ED, reliability of electrolyte and metabolite results measured by, 419–424
- Blood glucose ratio, CSF, and bacterial meningitis, without pleocytosis in potentially immunocompromised host (letter), 922
- Blood pressure
- high, in ED, adropin levels and target organ damage secondary to, 2061–2064

- higher prehospital, prolongs door to needle thrombolysis times, 1268–1272
- predischARGE, effect on follow-up outcomes in patients with severe hypertension in ED, 834–839
- Blood products transfusion, risk of transfusion-related acute lung injury after (letter), 1674–1675
- Blood urea nitrogen/creatinine ratio, –based hydration therapy, 3 months after acute ischemic stroke, favorable outcome of, 2414–2418 BLS. *see* Basic life support (BLS)
- Blunt abdominal trauma, 27 years after splenectomy, hemoperitoneum after, a case report, 1325.e3–1325.e4
- Blunt chest trauma, severe, sernal fractures and delayed cardiac tamponade due to, a case report, 758.e1–758.e3
- BNP. *see* Brain natriuretic peptide (BNP)
- Body mass index
associated with inappropriate tidal volumes in adults intubated in ED (letter), 1682–1683
impact on patient survival after therapeutic hypothermia after resuscitation, 722–725
- Body weight, novice rescuer, relationship with chest compression depth, during CPR, 2411–2413
- Bone Injection Gun, and Jamshidi intraosseous access devices, by paramedics with and without chemical-biological-radiological-nuclear personal protective equipment, comparison of (letter), 1307–1308
- Boxer fractures, ultrasound-guided ulnar nerve block for (letter), 1726–1727
- Brachial plexus blockade, hemidiaphragmatic paresis associated with, a case report, 1733.e5–1733.e7
- Bradycardia
caused by intravenous nicardipine in elderly patient with acute ischemic infarct, a case report, 761.e1–761.e2
and hypotension after synthetic cannabinoid use, a case report, 2055.e1–2055.e2
- Brain hypoxia, in avalanche victims, potassium as a potential predictive biomarker of (letter), 1315–1316
- Brain infarction, multiple, large, with sudden cardiac arrest patient, after therapeutic hypothermia, unexpected survival of, a case report, 2462.e1–2462.e3
- Brain ischemia, and hyperbaric oxygen therapy, efficacy in acute CO intoxication, dual-energy CT shows, a case report, 1327.e3–1327.e4
- Brain natriuretic peptide (BNP)
- B-type, normal, intermediate- to high-risk pulmonary embolism with, a case report, 2463.e1–2463.e3
consistent in using 14 different units instead of ng/L for (letter), 750–751
- Brain stem encephalopathy, hypertensive, mimicking central pontine myelinolysis, potential pitfall, a case report, 1918.e5–1918.e7
- Brest score, development of clinical prediction score for congestive heart failure diagnosis in emergency care setting, 2277–2283
- BRIPPED scan, for evaluation of ED patients with shortness of breath, 386–391
- Bronchial asthma, beneficial action of magnesium sulfate in (letter), 1693–1694
- Brucella*, –induced sublingual abscess causes acute shortness of breath, a case report, 1737.e3–1737.e4
- Brugada electrocardiogram pattern, induced by cannabis, a case report, 1738.e1–1738.e4
- Brugada syndrome
massive pulmonary embolism mimicking ECG pattern of, a case report, 933.e1–933.e2
warm-up, associated with Takotsubo cardiomyopathy, a case report, 2051.e1–2051.e3
- Burner and stinger syndrome, pediatric, magnetic resonance imaging for, 1442–1445
- Burns
sepsis, through inhibiting NLRP3 inflammasome activation, artemisinin protects mice against, 772–777
severe, effects of insulin combined with ethyl pyruvate on inflammatory response and oxidative stress in multiple-organ dysfunction syndrome rats with, 2154–2158
- 2,3-Butanedione monoxime, facilitates successful resuscitation in dose-dependent fashion in pig model of cardiac arrest, 1053–1058
- Bystanders
CPR, two-thirds receive no, 2011–2013
emergency assistance, decreases time to defibrillation in public venue, randomized controlled trial, 590–593
- Calcaneocuboid articulations (Chopart joints) dislocation, isolated dorsal, from low-energy mechanism, a case report, 1733.e1–1733.e4
- Cancer
disseminated, latent, mimicking bacterial meningitis, acute lethargy in young woman due to, diagnostic pitfall, a case report, 2050.e5–2050.e7
ED pilot intervention to facilitate outpatient acute care for patients with, 1934–1938
- Cannabinoids
synthetic
bradycardia and hypotension after, a case report, 2055.e1–2055.e2
hyperemesis, resulting in rhabdomyolysis and acute renal failure, a case report, 765.e1–765.e2
hyperthermia and severe rhabdomyolysis from, a case report, 121.e1–121.e2
hyperthermia and severe rhabdomyolysis from (letter), 1690–1691
- Cannabis, Brugada electrocardiogram pattern induced by, a case report, 1738.e1–1738.e4
- CAP. *see* Community-acquired pneumonia (CAP)
- Capnography
and cerebral oximetry with blood volume index, in intubated and hyperventilated patients, 1102–1107
use to assess quality of pediatric ventilation with 3 different airway modalities, 69–74
- Carbon dioxide
end-tidal
and occult injury, in trauma patients, 2146–2149
predicts sepsis and severe sepsis, prehospital screening tool utilizing, 813–819
- Carbon monoxide
factory, human, a case report, 2251.e3–2251.e4
intoxication, acute, dual-energy CT shows brain ischemia and hyperbaric oxygen therapy efficacy in, a case report, 1327.e3–1327.e4
poisoning
ability of firefighters to recognize (letter), 1710
acute, pentraxin 3, ischemia-modified albumin, and myeloperoxidase in predicting cardiac damage in, 1927–1930
in acute and chronic periods, cardiac effects of (letter), 1303
COHb levels predict long-term development of acute myocardial infarction in, 840–844
deaths in United States, 1999 to 2012 (letter), 904
myth busting in, 295–297
patients with headache, treatment in, 2140–2145
reversible left bundle-branch block due to, a case report, 342.e1–342.e3
severe neurologic impairment and uncommon MRI findings after, a case report, 119.e1–119.e2
- Carboxyhemoglobin (COHb)
extreme elevation, survival following (letter), 1168–1169
levels, predict long-term development of acute myocardial infarction in carbon monoxide poisoning, 840–844
- Cardiac arrest
asphyxial, rat model, protective effects of cyclosporine A and hypothermia on neuronal mitochondria in, 1080–1085
avalanche-induced, serum potassium concentration predicts brain hypoxia on CT after, 856–860

- blood potassium after avalanche-induced (letter), 1317–1318
- comparative effectiveness of standard CPR vs active compression-decompression CPR with CardioPump for treatment of, 542–547
- due to pulmonary embolism, thrombolysis during continuous chest compression in patient with, prolonged CPR–induced spinal cord injury, a case report, 681.e3–681.e5
- in ED, effect of backboard on compression depth during, simulation study, 274–277
- effects of Shen-Fu injection on coagulation-fibrinolysis disorders in porcine model of, 469–476
- effects of tibial intraosseous and intravenous administration of vasopressin on kinetics and survivability in, 429–432
- emergency medical services annotated literature in review 2016, 2193–2199
- failure of extracorporeal life support due to refractory ionized hypocalcemia during, a case report, 2251.e1–2251.e2
- and green tea (editorial), 576–577
- hypoxic and non-hypoxic, with non-cardiac etiology, prognostic value of gray matter to white matter ratio in, 1583–1588
- in-hospital
- initial PETCO₂ predicts outcomes of, 2367–2371
 - meta-analysis of outcomes of 2005 and 2010 cardiopulmonary resuscitation guidelines for adults with, 1133–1139
- most dangerous complication of intranasal cocaine abuse in young man, a case report, 1731.e5–1731.e7
- out-of-hospital
- assessment of oxidative stress after, 1561–1566
 - drowning-associated, relationship between drowning location and outcome after, 1799–1803
 - incidence and clinical features of intracranial hemorrhage causing, 2326–2330
 - interaction of diabetes mellitus and cardiac diseases on survival outcomes in, 702–707
 - patients, favorable neurological outcomes by early epinephrine administration within 19 minutes after emergency medical service call for, 2284–2290
 - patients, laryngeal tubes or masks used for (letter), 749–750
 - patients without STEMI, emergency coronary angiography in, a case report, 118.e1–118.e3
 - in patient with congenital long QT syndrome, extracorporeal CPR and therapeutic hypothermia for, a case report, 1321.e1–1321.e3
 - prehospital endotracheal intubation and survival after, results from the Korean nationwide registry, 128–132
 - prehospital therapeutic hypothermia after, 2209–2216
 - proper adjustment for sicker patients with (letter), 329
 - public-access automated external defibrillators and defibrillation for (letter), 2041–2042
 - resuscitated, with normal postresuscitation 12-lead ECG, a case report, 346.e1–346.e3
 - role of prehospital advanced airway management on outcomes for patients with, 2101–2106
 - sodium bicarbonate dilemma in patients with (letter), 1314–1315
 - studies must adjust for sicker patients properly (letter), 328–329
 - subarachnoid hemorrhage-associated, bispectral index monitoring in, a case report, 934.e1–934.e3
 - successful use of targeted temperature management in pregnancy after, a case report, 122.e3–122.e4
 - thrombolysis of patients with (letter), 916–917
- physicians are able to recognize ineffective (agonal) breathing as element of (letter), 1165
- pig model, 2,3-butanedione monoxime facilitates successful resuscitation in dose-dependent fashion in, 1053–1058
- post
- severe acidemia on arrival not predictive of neurologic outcomes in patients with, 425–428
 - who undergo therapeutic hypothermia, early neurologic examination is not reliable for prognostication in patients with, 975–979
- in prehospital situations, GCS used as prognostic factor in unconscious patients following (letter), 1178–1179
- rats, induced by ventricular fibrillation, benefits of respective and combined use of green tea polyphenols and ERK inhibitor on survival and neurologic outcomes in, 570–575
- resulting from myocardial infarction and outcome, blood lactate concentration after (letter), 1311–1313
- serum potassium cutoff for in-hospital triage after avalanche-induced (letter), 1317
- and severe hypothermia, successfully treated without external mechanical circulatory support, a case report, 1913.e5–1913.e6
- sudden, patient, with large multiple brain infarction, after therapeutic hypothermia, unexpected survival of, a case report, 2462.e1–2462.e3
- survivor treated with targeted temperature management, prevalence and risk factors for central diabetes insipidus in, 1400–1405
- traumatic, 15 minutes resuscitation duration appropriate before termination of, 505–509
- ventricular tachycardia, during central line placement, a case report, 114.e3–114.e4
- Cardiac disease, and diabetes mellitus, on survival outcomes in out-of-hospital cardiac arrest, interaction of, 702–707
- Cardiac effects, of carbon monoxide poisoning in acute and chronic periods (letter), 1303
- Cardiac injury, in acute carbon monoxide poisoning, pentraxin 3, ischemia-modified albumin, and myeloperoxidase in predicting, 1927–1930
- Cardiac stress testing, in patients with acute chest pain and negative troponins, clinical score to obviate the need for, 1421–1426
- Cardiac tamponade, delayed, and sternal fractures, due to severe blunt chest trauma, a case report, 758.e1–758.e3
- Cardiac valve prosthesis, after traumatic intracranial hemorrhage, discontinuing oral anti-coagulants in patient with, a case report, 1736.e5–1736.e7
- Cardiologist, direct involvement, in ED triage of cardiology patients (letter), 325–326
- Cardiopulmonary resuscitation (CPR)
- 2005 and 2010 guidelines, for adults with in-hospital cardiac arrest, meta-analysis of outcomes of, 1133–1139
 - bystander, training experience and self-efficacy of age and gender group, 1331–1337
 - bystander, two-thirds receive no, 2011–2013
 - in ED of urban academic hospital, outcome of patients receiving, 1595–1599
 - extended manikin, impact of feedback device on chest compression quality during, randomized crossover study, 1754–1760
 - extracorporeal, and therapeutic hypothermia, for out-of-hospital cardiac arrest in patient with long QT syndrome, a case report, 1321.e1–1321.e3
 - 3 feedback devices, and standard BLS by nurses, randomized trial of chest compressions effectiveness comparing, 381–385
 - hands-only, very brief training for laypeople in, effect of real-time feedback, 993–998
 - healthcare professionals' claims of having seen patient returning to normal activity after (letter), 925–927
 - hemorrhagic shock caused by intramuscular hematoma of pectoralis major after, a case report, 682.e5–682.e6
 - high-quality, rescuer factors predict, CPR–manikin-based study of health care providers, 20–24
 - induced spinal cord injury, prolonged, thrombolysis during continuous chest compression inpatient with cardiac arrest due to pulmonary embolism, a case report, 681.e3–681.e5
 - intraosseous access devices as method of vascular access during (letter), 321–322
 - intravascular access method to choose during (letter), 1160
 - McGrath-MAC vs Macintosh laryngoscopes for child tracheal intubation during (letter), 2453
 - mitigating hyperventilation during, 643–646
 - muscular fitness as mediator of quality, 1845–1849

- need for abdominal only, in treatment of hemorrhagic shock and trauma arrests (letter), 1156–1157
- neonatal, safe and effective, current guideline recommendations for, 1658–1660
- next step for, 97–99
- nurses use mechanical chest compression devices during (letter), 2044–2045
- phosphodiesterase-5 inhibition improves macrocirculation and microcirculation during, 162–166
- prolonged, after electric shock, survival without sequelae after, a case report, 679.e1–679.e2
- in rats, activation of autophagy improved neurologic outcome after, 1511–1518
- relationship between chest compression depth and novice rescuer body weight during, 2411–2413
- simulated, mechanical chest compression with LifeLine ARM device during (letter), 917
- standard, vs active compression-decompression CPR, with CardioPump for treatment of cardiac arrest, comparative effectiveness of, 542–547
- suprasternal notch as landmark of chest compression depth in, 433–436
- and teaching does influence retention of their knowledge, children can teach their parents, 1159–1160
- in ventricular fibrillation model of rats, edaravone improves survival and neurological outcomes after, 1944–1949
- videolaryngoscopy or direct laryngoscopy for child tracheal intubation during (letter), 2453
- CardioPump, treatment of cardiac arrest, comparative effectiveness of standard CPR vs active compression-decompression CPR with, 542–547
- Cardiovascular arrest, cocaine-induced, in rat model, intravenous lipid emulsion in resuscitation of, 1452–1454
- Cardiovascular disease
- ischemia modified albumin in (letter), 1699
 - outcomes in tertiary care centers in Japan (letter), 109–111
- Cardiovascular responses, to energy drinks in healthy population, 1205–1209
- Cardioversion, pharmacological, of recent-onset atrial fibrillation, safety and efficacy of, 1486–1490
- C5a receptor, on polymorphonuclear neutrophil and C5a2intra/C5a2 ratio level for early sepsis in ED, prognostic significance of, 2084–2089
- Carotid blowout, and cerebral gas embolism related to bidirectional carotid-esophageal fistula, serious complication of esophageal cancer under radiotherapy, a case report, 683.e5–683.e6
- Carotid endarterectomy, cerebral hyperperfusion syndrome, rare postoperative complication of, a case report, 2054.e1–2054.e2
- Carotid-esophageal fistula, bidirectional, carotid blowout and cerebral gas embolism related to, serious complication of esophageal cancer under radiotherapy, a case report, 683.e5–683.e6
- Case fatality rate, age effects on, of injury patients by mechanism, 515–520
- Catheter, -directed option for pulmonary embolism treatment (letter), 1691–1693
- Catheterization
- peripheral venous, influence of peripheral venous catheter size and resident postgraduate years on success of (letter), 1716–1718
 - short, ultrasound-guided placement, prospective evaluation in internal jugular veins of difficult venous access patients, 578–581
 - venous, topical ethyl chloride to reduce pain associated with, randomized crossover trial, 845–850
- Cauda equina syndrome, thoracic spinal cord compression masquerading as, a case report, 756.e3–756.e5
- Caudal traction, of ipsilateral arm, influence on ultrasound image for supraclavicular central venous catheterization, 851–855
- CBRN-PPE. *see* Chemical, biological, radiations, nuclear, personal protective equipment (CBRN-PPE)
- Celiac sprue, presenting as pseudo MI, a case report, 1187.e3–1187.e5
- Cellulitis
- failure of antibiotics trials in, 1645–1652
 - intravenous home infusion therapy instituted from 24-hour clinical decision unit for patients with, 1273–1275
- Central line placement, ventricular tachycardia cardiac arrest during, a case report, 114.e3–114.e4
- Central venous access, ultrasound-guided, which probe is preferred for viewing subclavian vein using supraclavicular approach?, 1761–1764
- Central venous catheterization, supraclavicular, influence of caudal traction of ipsilateral arm on ultrasound image for, 851–855
- Cerebellitis, acute, and multi-organ damage, methadone overdose causing, a case report, 343.e1–343.e3
- Cerebral blood volume index, with cerebral oximetry, and capnography, in intubated and hyperventilated patients, 1102–1107
- Cerebral edema, induced by hyperammonemia, a case report, 2461.e3–2461.e5
- Cerebral function, in elderly patients with pulmonary embolism, spectrum of disturbances in, 651
- Cerebral hyperperfusion syndrome, rare postoperative complication of carotid endarterectomy, a case report, 2054.e1–2054.e2
- Cerebral oximetry, with blood volume index and capnography in intubated and hyperventilated patients, 1102–1107
- Cerebral vasospasm, hemodynamic challenges in traumatic subarachnoid hemorrhage complicated by (letter), 904–906
- Cerebrospinal fluid (CSF)
- blood glucose ratio and bacterial meningitis without pleocytosis in potentially immunocompromised host (letter), 922
 - cellularity, normal, in pneumococcal meningitis, wider implications of, 657
 - Streptococcus pneumoniae* meningitis without pleocytosis of, a case report, 941.e1–941.e2
- Cervical cancer, screening, and HPV vaccination in ED, model for implementation of (letter), 648
- Cervical spine
- clearance, orthopedic literature 2015, 1663–1667
 - immobilization, may be of value following firearm injury to head and neck, 726–729
- Chamber volume, and electrocardiogram signal, in “false” atrial fibrillation, unexpected mismatch between, a case report, 1740.e1–1740.e2
- Chemical, biological, radiations, nuclear, personal protective equipment (CBRN-PPE)
- comparison of two chest compression techniques when using, randomized crossover manikin trial (letter), 913–915
 - comparison of two intravascular access techniques when using, randomized crossover manikin trial (letter), 1170–1172
- Chemical agents, for sedation of agitated patients in ED, 2426–2431
- Chest compressions
- active, hands-on defibrillation during, 2172–2176
 - combination with interposed abdominal compressions, in swine model of ventricular fibrillation, 968–974
 - comparison of two techniques when using CBRN-PPE randomized crossover manikin trial (letter), 913–915
 - continuous
 - ability of paramedics to perform endotracheal intubation during, randomized cadaver study comparing Pentax AWS and Macintosh laryngoscopes, 1835–1839
 - in patient with cardiac arrest due to pulmonary embolism, thrombolysis during, prolonged CPR-induced spinal cord injury, a case report, 681.e3–681.e5 - depth
 - during cardiac arrest in ED, effect of backboard on, simulation study, 274–277
 - in CPR, suprasternal notch as landmark of, 433–436
 - on mattress, flexible pressure sensor could correctly measure, 899–902
 - obtained using PocketCPR as educational tool according to smartphone attachment site, assessment of (letter), 2243–2246
 - relationship with novice rescuer body weight during CPR, 2411–2413

- effectiveness, comparing 3 feedback CPR devices and standard BLS by nurses, randomized trial of, 381–385
- interruption times, comparison across 2 automated devices, randomized, crossover simulation study, 57–62
- in manikin, utility of gum-elastic bougie for tracheal intubation during, randomized crossover trial, 54–56
- by measuring hand width and sternal length in young adults, identifying optimal hand placement site for, 407–411
- mechanical, with LifeLine ARM device during simulated CPR (letter), 917
- mechanical devices, nurses use, during CPR (letter), 2044–2045
- needle guides for venous catheter insertion during, crossover simulation trial, 989–992
- performed by nurses, tracheal intubation with Macintosh laryngoscope with and without (letter), 2448–2449
- quality, during extended manikin CPR, impact of feedback device on, randomized crossover study, 1754–1760
- in simulated resuscitation, effect of bed width on quality of, randomized crossover manikin study, 2272–2276
- Chest pain**
- acute
- at ED, diagnostic performance of smartphone reading of the coronary CT angiography in patients with, 1794–1798
 - and negative troponins, clinical score to obviate the need for cardiac stress testing in patients with, 1421–1426
 - rupture of noncoronary sinus of Valsalva into right atrium, unusual cause of, a case report, 2052.e1–2052.e3
 - and coronary artery disease, copeptin and high-sensitivity cardiac troponin to exclude severe coronary stenosis in patients with, 493–498
 - geriatric patients, in ED observation unit, prospective evaluation of outcomes among, 207–211
 - and headache, posterior reversible encephalopathy syndrome after bevacizumab treatment presenting to ED as, a case report, 1916.e1–1916.e2
 - high-frequency QRS analysis superior to conventional ST-segment analysis of women with, 437–442
 - increased observation services in Medicare beneficiaries with, 16–19
 - low-risk patients, presenting to ED chest pain unit, barriers to outpatient stress testing follow-up for, 790–793
- Chest radiography, comparing posteroanterior with lateral and anteroposterior, in initial detection of parapneumonic effusions, 2402–2407
- Chest wall thickness, and needle thoracostomy success rates, determination at second and fifth intercostal spaces, cadaver-based study, 2310–2314
- CHF. *see* Chronic heart failure (CHF)
- Child abuse, Lipschütz ulcer cause of misdiagnosis when suspecting, a case report, 1326.e1–1326.e2
- Childbirth, in EMS practice (letter), 1888
- Children. *see* Pediatrics
- Cholecystitis, gangrenous, acute, use of dynamic CT attenuation value for diagnosis of, 2306–2309
- Chronic diseases, multimorbid, sex differences in frequent ED use among those with, 2127–2131
- Chronic eosinophilic pneumonia (Carrington disease), diagnosis of, a case report, 1734.e5–1734.e6
- Chronic heart failure (CHF), acute decompensated, NT-proBNP, IVC ultrasound, and bioelectrical impedance analysis for diagnosis of, 1817–1822
- C1 inhibitor**
- in sepsis and clinical outcome (letter), 1179–1180
 - serial change in patients with sepsis, 594–598
 - serial change in patients with sepsis (erratum), 1741–1742
- Circulatory shock, intraosseous route, new route to life in patients with (letter), 922–923
- Cirrhosis
- in ED, patients with, early predictors of infection and mortality, 25–29
 - serotonin syndrome associated with therapeutic metaxalone in patient with, a case report, 346.e5–346.e6
- Clavicular fracture, orthopedic literature 2015, 1663–1667
- Clenbuterol, causing NSTEMI in teenage female desiring to lose weight, a case report, 1739.e5–1739.e7
- Clinical decision support, impact on adherence to OAR, 412–418
- Clinical decision unit, 24-hour, for patients with cellulitis, intravenous home infusion therapy instituted from, 1273–1275
- Clonidine, centrally acting antihypertensive, combined to nitroglycerin, venous dilator, to handle severe pulmonary edema, a case report, 676.e5–676.e7
- Closed reduction, of anterior shoulder dislocation in super obese, a case report, 1181.e1–1181.e2
- Clotiapine, poisoning, pediatric, clinical manifestations and toxicokinetics, a case report, 2469.e1–2469.e2
- C-MAC, and Macintosh laryngoscopes, for face-to-face intubation in manikin, randomized crossover trial comparing (letter), 920–922
- Coagulation–fibrinolysis disorder, in porcine model of cardiac arrest, effects of Shen-Fu injection on, 469–476
- Cocaine
- induced cardiovascular arrest, in rat model, intravenous lipid emulsion in resuscitation of, 1452–1454
 - intranasal, abuse in young man, cardiac arrest most dangerous complication of, a case report, 1731.e5–1731.e7
- Coccidioidomycosis, presentations in ED (letter), 906–910
- Cognitive health, and risk of ED revisit in underserved older adults, 1973–1976
- COHb. *see* Carboxyhemoglobin (COHb)
- Collapse, and syncope, in acute pulmonary embolism, 1251–1257
- Colles fractures, in ED, ultrasound-guided supracondylar radial nerve block for (letter), 1718–1720
- Colorectal cancer, perforated, with liver metastasis, catastrophic pneumoperitoneum in patient with, a case report, 344.e3–344.e5
- Coma blister, in nontraumatic rhabdomyolysis, a case report, 1324.e1–1324.e2
- Comatose, survivors after hanging, neurologic outcome of, 1467–1472
- Communication, clinician–patient, patient treatment in ED hallways and patient perception of (letter), 1163–1164
- Community-acquired pneumonia (CAP), patients, validation of NEWS-L in, 536–541
- Complications
- en-route, prediction during interfacility transport by outcome predictive scores in ED, 877–882
 - thromboembolic, and atrial standstill (letter), 932
- Computed tomography (CT)
- angiogram, of chest, saddle embolism diagnosed by point-of-care thoracic echocardiography before, a case report, 2467.e1–2467.e2
 - angiography
 - in acute stroke (revisiting the 4Ps of imaging), 282–287
 - epipericardial fat necrosis clinically mimicking pulmonary embolism, a case report, 2056.e5–2056.e6
 - after avalanche-induced cardiac arrest, serum potassium concentration predicts brain hypoxia on, 856–860
 - and bedside point-of-care ultrasound, in elbow injuries, comparison of, 2186–2190
 - cardiac, and chance of missed acute aortic syndrome, limiting scan range of, 2007–2010
 - contrast, relation of contrast nephropathy to adverse events in pulmonary emboli patients diagnosed with, 1247–1250
 - coronary, angiography, diagnostic performance of smartphone reading in patients with acute chest pain at ED, 1794–1798
 - dual-energy, shows brain ischemia and hyperbaric oxygen therapy efficacy in acute CO intoxication, a case report, 1327.e3–1327.e4
 - dynamic, attenuation value, use for diagnosis of acute gangrenous cholecystitis, 2306–2309

- findings, in patients with acute mesenteric ischemia, baseline disease is more important predictor of intestinal necrosis than, 2261–2265
- head
and near-infrared spectroscopy, interpretations of ED patients with minor head injury, comparison of, 1364–1368
- randomized single-blinded clinical trial on effects of nursery songs for infants and young children's anxiety before and during (letter), 663
- interpretations in multiply injured patients, comparison of emergency physicians and on-call radiologists, 2331–2335
- measurement, optic nerve sheath diameter/eye diameter, quantitative evaluation of intracranial pressure by, 2336–2342
- noncontrast abdomen/pelvis, in evaluation of older adults (letter), 2230–2232
- normal initial, and minor head injury, outcomes of warfarinized patients with, 75–78
- and pain severity, effects on analgesia prescription in acute appendicitis, 36–39
- in patients with uncomplicated appendicitis, effect of time delays for appendectomy as observed on, 167–169
- pulmonary angiogram, risk of pulmonary embolism after prior negative, 1968–1972
- settings for optimal detection of wooden foreign bodies (letter), 2237–2238
- for trauma, focused assessment with (letter), 1721–1723
- and ultrasonography, in assessment of acute appendicitis, Bayesian comparative model of, 2070–2073
- whole-body, who will benefit from (letter), 1907–1908
- Concierge, impact on ED revisits and outpatient follow-up (letter), 2445–2446
- Consciousness, clear, even with glucose level of 5 mg/dL (0.2 mmol/L), patient with, a case report, 941.e3–941.e4
- Contraception, emergency, survey of emergency physicians regarding (letter), 1176–1178
- Contrast staining, after acute ischemic stroke from hemorrhagic transformation during emergency evaluation, differentiation of, a case report, 2255.e5–2255.e6
- Convergence spasm, very rare paroxysmal symptom in multiple sclerosis, a case report, 117.e5–117.e6
- Cooling, brain, rapid and selective method, using vortex tube, 887–894
- Copeptin, and high-sensitivity cardiac troponin to exclude severe coronary stenosis in patients with chest pain and coronary artery disease, 493–498
- Copper deficiency, presenting as progressive peripheral neuropathy, a case report, 340.e5–340.e6
- Coronary anatomy, troponin leak, and prognosis, electrocardiography changes in acute aortic dissection—association with, 1431–1436
- Coronary artery disease
and chest pain, copeptin and high-sensitivity cardiac troponin to exclude severe coronary stenosis in patients with, 493–498
and SCUBE1, relation between (letter), 647–648
severity, and fibrinogen to albumin ratio, in patients with STEMI, relationship between, 1037–1042
- Coronary artery dissection, spontaneous, with ST elevation in aVR and posterior leads, a case report, 2470.e5–2470.e7
- Coronary occlusion, acute, diagnosis in patients with non-STEMI by point-of-care echocardiography with speckle tracking, a case report, 1914.e3–1914.e6
- Coronary stenosis, severe, in patients with chest pain and coronary artery disease, copeptin and high-sensitivity cardiac troponin to exclude, 493–498
- Corrected flow time, noninvasive ultrasound measure to detect preload reduction by nitroglycerin, 1859–1862
- Corticosteroids, inhaled, increase blood neutrophil count by decreasing expression of neutrophil adhesion molecules Mac-1 and L-selectin, 1977–1981
- Cost-effectiveness analysis, of point-of-care testing for dehydration in pediatric ED, 1573–1575
- Costs
of decontaminating ED after finding bed bug, results from a single academic medical center (letter), 649
of ED episodes of care in United States, 357–365
hospital, and prehospital, of emergency care for frequent ED patients, 459–463
of living, weighing numerous variables involved (letter), 1168
savings associated with transfer of trauma patients within accountable care organization, 455–458
and tackling causes, of ED presentation for American football injuries, 1198–1204
- CPR, Cardiopulmonary resuscitation (CPR). *see* Cardiopulmonary resuscitation (CPR)
- C-reactive protein, ultrasound and Alvarado score, predictive value in acute appendicitis, 189–192
- Cricothyroid membrane, comparison of ultrasound vs anatomic landmark to locate (letter), 1696
- Cricothyroidotomy
comparison of ultrasonography and surface landmarks in detecting localization for, 254–256
surgical, emergency, postobstructive pulmonary edema that developed immediately after, a case report, 936.e1–936.e2
- Critical access hospital, ED to quaternary medical center, 1427–1430
- Critically ill, 2 children, delayed sequence intubation with ketamine in, a case report, 1190.e1–1190.e2
- Crotalinae* envenomation, simulated, variability of limb measurements performed by emergency medicine nurses in (letter), 1683–1685
- Crotaline envenomation, in children, new F(ab')₂ antivenom for treatment of, 2003–2006
- Cryothyroidotomy, anatomic landmark *versus* ultrasound approach to (letter), 1696–1697
- CSF. *see* Cerebrospinal fluid (CSF)
- CT. *see* Computed tomography (CT)
- Cunningham technique
for dislocated shoulder reduction (letter), 1890–1891
redux, Sool's method of dislocated shoulder reduction (letter), 1891
- Cyclosporine A, and hypothermia, protective effects on neuronal mitochondria in rat asphyxial cardiac arrest model, 1080–1085
- CYP2C19, drug-drug and drug-gene interactions in ED patients, 245–249
- Cysticercosis
optic nerve, retroorbital, a case report, 2461.e1–2461.e2
role of *Taenia asiatica* in human cases of (letter), 2030
- Dabigatran etexilate, use, in patient with acute kidney injury, successful hemostasis and reversal of highly elevated PT/INR after, a case report, 758.e5–758.e6
- Dagnosis, late, Wellens syndrome in patient presenting with atypical ACS, a case report, 338.e3–338.e5
- Dapsone, overdose, acute, incidence and patterns of hemolytic anemia in, 366–369
- Death
carbon monoxide poisoning, in United States, 1999 to 2012 (letter), 904
opioid, shows startling findings, analysis of prescription patterns in (letter), 748–749
prescription-related, medical examiner, and prescription drug monitoring program data, description of, 510–514
- Debridement, hydrosurgical, in ED, effective prevention of posttraumatic tattoo using, a case report, 759.e3–759.e4
- Decision making
shared
in ED, 1668–1672
to reduce unnecessary radiography in extremity trauma, physician and patient gestalt lead to, 349–356
to reduce unnecessary radiography in extremity trauma, physician and patient gestalt lead to (erratum), 348
- Deep vein thrombosis, upper extremity, diagnostic limitations of imaging, Paget-Schroetter syndrome, a case report, 683.e1–683.e3

Defibrillation

- hands-on, during active chest compressions, 2172–2176
- and public-access automated external defibrillators, for out-of-hospital cardiac arrest (letter), 2041–2042
- in public venue, emergency dispatcher assistance decreases time to, randomized controlled trial, 590–593

Defibrillator

- automated external
 - paramedic students' knowledge and attitude regarding (letter), 1887–1888
 - public-access, and defibrillation, for out-of-hospital cardiac arrest (letter), 2041–2042
- cardiac, implantable, electrical shocks, ED evaluation and management of, 1140–1147

Dehydration

- in children, sonographic aorta/IVC cross-sectional area index for evaluation of, 1840–1844
- in pediatric ED, cost-effectiveness of point-of-care testing for, 1573–1575

Delirium, assessment, quick and easy, for nonphysician research personnel, 1031–1036

Desmopressin, effectiveness in renal colic pain management, 1535–1541

De Winter syndrome, impending anterior MI (letter), 2450–2451

Dexketoprofen, IV, ketoprofen gel improves low back pain in addition to, 1458–1461

Diabetes insipidus, central, in cardiac arrest survivor treated with targeted temperature management, prevalence and risk factors for, 1400–1405

Diabetes mellitus, and cardiac diseases, on survival outcomes in out-of-hospital cardiac arrest, interaction of, 702–707

Diabetic ketoacidosis, looking through electrocardiogram window, 263–265

Diagnosis

- accuracy, of finger fractures, bedside ultrasonography for, 809–812
- of acute coronary occlusion in patients with NSTEMI by point-of-care echocardiography with speckle tracking, a case report, 1914.e3–1914.e6
- of acute decompensated CHF, NT-proBNP, IVC ultrasound, and bioelectrical impedance analysis for, 1817–1822
- of acute gangrenous cholecystitis, use of dynamic CT attenuation value for, 2306–2309
- of ascending aortic dissection, emergency physician focused cardiac ultrasound improves, 486–492
- bedside, in ED, impending paradoxical embolus, a case report, 1917.e3–1917.e5
- challenges with stump appendicitis, a case report, 1188.e–1188.e2
- of chronic eosinophilic pneumonia (Carrington disease), a case report, 1734.e5–1734.e6
- congestive heart failure, in emergency care setting, Brest score, development of clinical prediction score for, 2277–2283
- differential
 - of acute pericarditis, role of combined ECG criteria in (letter), 1309
 - between septic arthritis and transient synovitis in ED, guideline for, 1631–1636
- emergent, and management of TASER penetrating ocular injury, a case report, 1740.e3–1740.e5
- intraabdominal, prediction model to identify patients without concerning, 1354–1358
- of irreducible posterolateral knee dislocation, a case report, 2256.e3–2256.e4
- limitations of imaging upper extremity deep vein thrombosis, Paget-Schroetter syndrome, a case report, 683.e1–683.e3
- and management, of lateral subtalar fracture-dislocation with maintenance of talonavicular joint, a case report, 2055.e3–2055.e5
- and management, of metacarpal fractures, effectiveness of bedside point-of-care ultrasonography in, contribution for management of metacarpal fractures (letter), 674

marker, potential, for peptic ulcer perforation, neutrophil-to-lymphocyte ratio, 403–406

of metacarpal fractures, role of ultrasonography in, 1868–1871

or prognostic biomarker, of acute appendicitis in children, evaluation of proadrenomedullin as, 2298–2305

performance of smartphone reading of coronary CT angiography in patients with acute chest pain at ED, 1794–1798

of peritonsillar abscess, using bedside emergency ultrasound to change, a case report, 1186.e5–1186.e7

of phalanx fractures, accuracy of bedside ultrasonography for (letter), 1698 pitfall

acute lethargy in young woman due to latent disseminated cancer mimicking bacterial meningitis, a case report, 2050.e5–2050.e7

and characteristics, of spontaneous visceral artery dissection in ED, 1092–1096

poststernotomy mediastinitis in ED, 618–622

prognosis, and therapeutic guidance of sepsis, clinical value of soluble uPAR in, 375–380

and prognostic tool at every step in pathway of care for acute heart failure, lung ultrasound, 656–657

of Takotsubo syndrome, in ED, pivotal role of pre-admission electrocardiogram for (letter), 903–904

test, for acute appendicitis, spot urinary 5-hydroxyindoleacetic acid is not ideal, 1750–1753

tool, unusual, endobronchial ultrasound, for pulmonary embolism, a case report, 684.e1–684.e2

and treatment of uncontrolled chronic hypertension, barriers to emergency physician (letter), 2241–2242

Diaphragmatic excursion

acute dyspnea by, practicality sustainable in ED (letter), 2441–2442

measurement in emergency patients with acute dyspnea, 1653–1657

DIC. *see* Disseminated intravascular coagulation (DIC)

Dietary supplement, survival after profound acidosis and rhabdomyolysis due to use of, a case report, 2259.e1–2259.e3

Dieulafoy lesion, little known sleeping giant of gastrointestinal bleeds, a case report, 2464.e3–2464.e5

Digoxin, intravenous lipid emulsion prolongs survival in rats intoxicated with, 1112–1116

Discharge against medical advice, and acute alcohol consumption, of injured patients in ED, association between, 464–468

Disease control guidelines, for ED patients with sexually transmitted diseases, compliance with centers for (letter), 1727–1729

Dislocation

anterior shoulder

effectiveness of newly developed reduction method of, 1406–1410

presenting as hypovolemic shock, axillary artery laceration after, a case report, 1739.e1–1739.e4

in super obese, closed reduction of, a case report, 1181.e1–1181.e2

atraumatic bilateral posterior shoulder, rare case of sleep deprivation, a case report, 1322.e3–1322.e4

knee, irreducible posterolateral, diagnosis, arthroscopic view, and treatment of, a case report, 2256.e3–2256.e4

shoulder

reduction, Cunningham technique for (letter), 1890–1891

reduction, Sool's method of, Cunningham technique redux (letter), 1891

of talonavicular and calcaneocuboid articulations (Chopart joints), isolated dorsal, from low-energy mechanism, a case report, 1733.e1–1733.e4

Disseminated intravascular coagulation (DIC), recombinant human soluble thrombomodulin and short-term mortality of infection patients with, 1876–1882

Distal radius fracture

reduction using looped-stockinette technique (letter), 2038–2040

supracondylar ultrasound-guided nerve block for reduction of, novel technique in ED, 912–913

- supracondylar ultrasound-guided nerve block for reduction of, novel technique in ED (letter), 1172
- Disulfide, thiol, homeostasis, prognostic importance in patients with acute pulmonary thromboembolism, 2315–2319
- Dizziness, in EDs, underuse of information about timing and triggers in diagnosing (letter), 1687–1689
- Doctors, junior, trained to use intraosseous access (letter), 107
- Dog bites, of extremities, serious, negative pressure wound therapy for, prospective randomized trial, 1006–1010
- Domestic violence, and sexual assault, prevalence of strangulation in survivors of, 1281–1285
- Doxapram, treatment of ethanol poisoning and associated hypoventilation with, a case report, 2253.e1–2253.e2
- Drowning
 - assessing efficacy of rescue equipment in lifeguard resuscitation efforts for, 480–485
 - relationship between location and outcome, after drowning-associated out-of-hospital cardiac arrest, 1799–1803
 - timeline, new systematic model of drowning process (letter), 2224–2226
- Drug abuse, prescription, prescribing controlled medications to patients who die of, 30–35
- Dysphagia, and hoarseness, aortic dissection manifesting as, Ortnier's syndrome, a case report, 1185.e1–1185.e3
- Dyspnea
 - acute
 - by diaphragmatic excursion, practicality sustainable in ED (letter), 2441–2442
 - diaphragmatic excursion measurement in emergency patients with, 1653–1657
 - emergency endotracheal intubation under fluoroscopy guidance for patients with, 2177–2181
 - inflammatory biomarkers predicting prognosis in patients with, 370–374
 - in emergency medicine, Keutel syndrome, rare cause of, a case report, 935.e3–935.e5
 - with mediastinal mass, emergent radiation therapy as definite airway management for, a case report, 2258.e5–2258.e6
- Early goal-directed therapy (EGDT)
 - end of, 292–294
 - what have we learned from (letter), 2449–2450
- Earthquake, Yushu, 2010, China, profile and procedures for fractures among 1323 fracture patients from, 2132–2139
- ECG. *see* Electrocardiography (ECG)
- Echocardiography
 - marker, handy, in acute pericarditis, echo probe sign (letter), 1883–1884
 - point-of-care
 - physical examination along with, indisputable right path (letter), 673
 - with speckle tracking, diagnosis of acute coronary occlusion in patients with NSTEMI, a case report, 1914.e3–1914.e6
 - strain, in acute heart failure (letter), 2234–2236
 - and serum troponin I
 - in anaphylaxis, evaluation of myocardial injury through, 140–144
 - in anaphylaxis, myocardial injury through, Takotsubo cardiomyopathy and Kounis hypersensitivity-associated ACS (letter), 650–651
 - transesophageal, for extracorporeal life support vascular cannula placement, emergency physician-performed, 1637–1639
- ECMO, Extracorporeal membrane oxygenation (ECMO). *see* Extracorporeal membrane oxygenation (ECMO)
- ECS. *see* Emergency Coma Scale (ECS)
- ED, Emergency department (ED). *see* Emergency department (ED)
- Edaravone, improves survival and neurological outcomes after CPR in ventricular fibrillation model of rats, 1944–1949
- Edema, upper extremity, caused by sotalol, a case report, 1920.e5–1920.e6
- Editorials, 576–577
- Education, ultrasound-guided intravenous, does not increase utilization, alteration of timing of nursing continuing education units for (letter), 2248–2249
- EGDT. *see* Early-goal directed therapy
- Elbow injury, comparison of bedside point-of-care ultrasound and computed tomography in, 2186–2190
- Elbow trauma, blunt, performance of 4-way ROM test for radiographic injuries after, 235–239
- Elderly. *see* Geriatric patients
- Electrical burn, experimental model, effects of N-acetylcysteine and ethyl pyruvate on ischemia-reperfusion injury in, 1217–1224
- Electric shock, survival without sequelae after prolonged CPR after, a case report, 679.e1–679.e2
- Electrocardiogram
 - in ACS patient, high-risk electrocardiographic presentations lacking anatomically oriented ST-segment elevation, 611–617
 - combined criteria, role in differential diagnosis of acute pericarditis (letter), 1309
 - pre-admission, pivotal role for diagnosis of Takotsubo syndrome in ED (letter), 903–904
 - reading, influence of computerized interpretation of (letter), 2031–2032
 - right- and left-sided, simultaneous recording in acute pulmonary embolism, a case report, 1183.e5–1183.e7
 - right-sided, usage in acute pulmonary embolism, 1437–1441
 - signal, and chamber volume, in “false” atrial fibrillation, unexpected mismatch between, a case report, 1740.e1–1740.e2
 - window, looking at diabetic ketoacidosis through, 263–265
- Electrocardiography (ECG)
 - abnormalities, predict neurogenic pulmonary edema in patients with subarachnoid hemorrhage with, 79–82
 - changes in acute aortic dissection—association with troponin leak, coronary anatomy, and prognosis, 1431–1436
 - characteristics of acute embolism in pulmonary trunk and main pulmonary arteries, 212–217
 - 12-lead, resuscitated out-of-hospital cardiac arrest with normal postresuscitation, a case report, 346.e1–346.e3
 - pattern
 - of Brugada syndrome, massive pulmonary embolism mimicking, a case report, 933.e1–933.e2
 - de Winter, evolution to (letter), 330–332
 - presentations, high-risk, lacking anatomically oriented ST-segment elevation, electrocardiogram in ACS patient, 611–617
 - recognition, of high lateral infarct, South African flag sign, teaching tool for easier (letter), 107–109
- Electrolyte, and metabolite results, measured by blood gas analyzer in ED, reliability of, 419–424
- Electrolyte imbalance, in ED, diseases associated with, age-related differences, 1923–1926
- Electronic cigarette, explosion injuries, a case report, 2252.e1–2252.e3
- Electronic medical record
 - alert, for severe sepsis among ED patients, effect of, 185–188
 - can reliably identify errors and adverse events in ED, 1043–1048
- Embolism
 - gas, cerebral, and carotid blowout, related to bidirectional carotid-esophageal fistula, serious complication of esophageal cancer under radiotherapy, a case report, 683.e5–683.e6
 - source, or risk factors, acute bilateral lower extremity arterial occlusion without, a case report, 762.e3–762.e4
- Emergency care
 - for frequent ED patients, prehospital and hospital costs of, 459–463
 - increase in moped injuries requiring, 2000–2002
 - setting, Brest score, development of clinical prediction score for congestive heart failure diagnosis in, 2277–2283
 - subsequent use, and weight loss surgery, population-based cohort study, 861–865

- Emergency Coma Scale (ECS), evaluation of accuracy of, E-COMET STEP II (letter), 100–101
- Emergency department (ED)
- 1939, sex differences in STEMI activation for patients presenting to, 1939–1943
 - academic, adult and pediatric, impact of scribes on patient throughput in, 1982–1985
 - accuracy of rapid sequence intubation medication dosing in obese patients intubated in, 2423–2425
 - acupuncture vs intravenous morphine in management of acute pain in, 2112–2116
 - acute dyspnea by diaphragmatic excursion, practicality sustainable in (letter), 2441–2442
 - admission as quality measure, critical analysis of unplanned ICU transfer within 48 hours from, 1505–1510
 - admissions, for ambulatory care-sensitive condition conditions, association of mental health disorders and Medicaid with, 820–824
 - adropin levels and target organ damage secondary to high blood pressure in, 2061–2064
 - adult patients, use of inhaled nitrous oxide for analgesia in, 269–273
 - after alcohol intake, quantitative analysis of high plasma lactate concentration in patients in, 825–829
 - among those with multimorbid chronic diseases, sex differences in frequent use, 2127–2131
 - aortoduodenal fistula as cause of abdominal pain in, a case report, 2468.e1–2468.e4
 - assessing patient activation and health literacy in, 93–96
 - assessing severity, immediacy, and ideal setting in patients in, 1276–1280
 - association between acute alcohol consumption and discharge against medical advice of injured patients in, 464–468
 - bedside sonographic measurement of optic nerve sheath diameter as predictor of intracranial pressure, 963–967
 - bibliometric analysis of top 100 most-cited clinical studies on ultrasound in, 1210–1216
 - body mass index associated with inappropriate tidal volumes in adults intubated in (letter), 1682–1683
 - characteristics and diagnostic pitfalls of spontaneous visceral artery dissection in, 1092–1096
 - characteristics and outcomes for acute heart failure in elderly patients presenting to, 2159–2166
 - characteristics of heroin-dependent patients seeking asthma care in, 895–898
 - characterizing children presenting to three freestanding (letter), 2045–2048
 - chemical agents for sedation of agitated patients in, 2426–2431
 - chest pain unit, barriers to outpatient stress testing follow-up for low-risk chest pain patients presenting to, 790–793
 - clinical trials in, some suggestions (letter), 670
 - comparing patients who leave prematurely, before vs after medical evaluation, National Hospital Ambulatory Medical Care Survey analysis, 830–833
 - contrast-enhanced ultrasound as imaging technique for patients with acute flank pain into (letter), 324–325
 - cost of decontaminating, after finding bed bug, results from a single academic medical center (letter), 649
 - critical access hospital, to quaternary medical center, 1427–1430
 - demographic characteristics of individuals admitted to the Hospital for Suicidal Ideation in (letter), 1174–1175
 - diagnosing poststernotomy mediastinitis in, 618–622
 - diagnostic performance of smartphone reading of the coronary CT angiography in patients with acute chest pain at, 1794–1798
 - difficult access patients, randomized controlled trial assessing use of ultrasound for nurse-performed intravenous placement in, 1950–1954
 - discharged home, increased door to admission time is associated with prolonged throughput for patients in, 1783–1787
 - diseases associated with electrolyte imbalance in, age-related differences, 1923–1926
 - effective prevention of posttraumatic tattoo using hydrosurgical debridement in, a case report, 759.e3–759.e4
 - effect of backboard on compression depth during cardiac arrest in, simulation study, 274–277
 - effect of electronic medical record alert for severe sepsis among patients in, 185–188
 - effect of predischARGE blood pressure on follow-up outcomes in patients with severe hypertension in, 834–839
 - effect of surgical consult in treatment of abdominal pain in older adults in, 1524–1527
 - efficacy of transcatheter cardiac pacing in, 2090–2093
 - efficiency, drivers of, statistical and cluster analysis of volume, staffing, and operations, 155–161
 - elderly fall patients, serious conditions for, 1394–1399
 - episodes, costs of care in United States, 357–365
 - evaluation and management of implantable cardiac defibrillator electrical shocks, 1140–1147
 - fewer visits, among US private sector working adults, paid sick leave is associated with, 784–789
 - giant bladder (10,500 mL in volume) in, a case report, 1327.e1–1327.e2
 - guideline for differential diagnosis between septic arthritis and transient synovitis in, 1631–1636
 - hallways, and patient perception of clinician-patient communication, patient treatment in (letter), 1163–1164
 - health care-associated acute pyelonephritis associated with inappropriate empiric antibiotic therapy in, 1415–1420
 - hemophagocytic lymphohistiocytosis in, a case report, 2057.e5–2057.e8
 - in high immunoglobulin E values at admission predict mortality inpatients with sepsis, 1589–1594
 - and homelessness, myths and facts (letter), 307
 - and homelessness, myths and facts (reply), 307–308
 - Icare vs Tono-Pen in (letter), 670–673
 - impact of freestanding on regional emergency medical services system, 1342–1346
 - impact of prospective verification of intravenous antibiotics in, 2392–2396
 - impact of push-dose phenylephrine use on subsequent preload expansion in, 2419–2422
 - impending paradoxical embolus, bedside diagnosis in, a case report, 1917.e3–1917.e5
 - information technology, proactive approach to, case of pediatric procedural sedation (letter), 1901–1902
 - initial presentation in patients with intracranial hematoma, prognostic value of perihematomal edema area at, 1241–1246
 - lack of awareness among emergency physicians of potentially inappropriate medications in elderly patients at (letter), 1888–1890
 - length of stay during mass casualty incident, factors associated with, 1462–1466
 - length of stay in, variation across classifications of clinical condition and patient discharge disposition, 83–87
 - lidocaine as adjuvant to morphine improve pain relief in patients with acute renal colic presenting to, 443–448
 - maximizes patient confidence in emergency physicians, point-of-care ultrasonography at, 657–659
 - May-Thurner syndrome, curious syndrome in, a case report, 1920.e1–1920.e3
 - mechanical ventilator settings and lung protective ventilation in, 1446–1451
 - Medicaid beneficiaries who continue to use, focus on Illinois Medical Home Network, 197–201
 - medical record reviewers can reliably identify errors and adverse events in, 1043–1048
 - with meningitis, validation of bacterial meningitis score in adults presenting to, 1265–1267
 - mesoamerican nephropathy a novel case of kidney failure in US, a case report, 1323.e5–1323.e6

- model for implementation of cervical cancer screening and HPV vaccination in (letter), 648
- neuro-ultrasonography and mental foramen (letter), 2240–2241
- nontraumatic splenic rupture as initial manifestation of acute myeloid leukemia in, a case report, 1731.e1–1731.e3
- novel technique in, supracondylar ultrasound-guided nerve block for reduction of distal radius fractures, 912–913
- novel technique in, supracondylar ultrasound-guided nerve block for reduction of distal radius fractures (letter), 1172
- observation unit, prospective evaluation of outcomes among geriatric chest pain patients in, 207–211
- observation units, venous thromboembolism prophylaxis in (letter), 2238–2240
- opioid prescribing, is not associated with higher patient satisfaction scores (letter), 2032–2034
- optimal dosing of intravenous ketamine for procedural sedation in children in, 1347–1353
- partial segmental thrombosis of corpus cavernosum presenting to, a case report, 1182.e3–1182.e5
- patients older than 50 years, nonsuicidal self-injury and suicide attempts among, comparison of risk factors and ED visit outcomes, 1016–1021
- patients' perspectives on insufficiency of scales to rate their pain in (letter), 2246–2247
- patients with abdominal pain, self-assessment of hunger among, lack of association with disease severity, 104–106
- patients with acute dyspnea, diaphragmatic excursion measurement in, 1653–1657
- patients with anaphylaxis, appropriateness of epinephrine use in, 174–179
- patients with cirrhosis in, early predictors of infection and mortality, 25–29
- patients with confirmed sepsis, presenting hemodynamic phenotypes in, 2291–2297
- patients with minor head injury, comparison of near-infrared spectroscopy and head CT interpretations of, 1364–1368
- patients with sepsis, prognostic role of non-critical lactate levels for in-hospital survival time among, 170–173
- patients with sexually transmitted diseases, compliance with centers for disease control guidelines for (letter), 1727–1729
- patients with shortness of breath, BRIPPED scan for evaluation of, 386–391
- for patients with solid malignancy, application of high-flow nasal cannula in (letter), 2222–2223
- patients with unanticipated hyperkalemia, succinylcholine in (letter), 1706–1707
- patient with unanticipated hyperkalemia, succinylcholine in, a case report, 2054.e3–2054.e5
- pediatric
- with abdominal pain who are at low risk for acute appendicitis, prospective validation of biomarker panel to identify, 1373–1382
 - cost-effectiveness of point-of-care testing for dehydration in, 1573–1575
 - experience, REDCap (letter), 2048–2049
 - treatment of septic hip in, retrospective case series analysis, 602–605
- pharmacist impact on time to antibiotic administration in patients with sepsis in, 2117–2121
- pharmacist monitoring of provider antibiotic selection aids appropriate treatment for outpatient UTI, 1600–1603
- pilot intervention to facilitate outpatient acute care for cancer patients, 1934–1938
- pivotal role of pre-admission electrocardiogram for diagnosis of Takotsubo syndrome in (letter), 903–904
- placement of transvenous pacemakers, outcomes of, 1411–1414
- Plesiomonas shigelloides* meningitis in an adult in, a case report, 1329.e1–1329.e2
- point-of-care ultrasonography for management of shoulder dislocation in, 866–870
- point-of-care ultrasound in pediatric small bowel obstruction, a case report, 2464.e1–2464.e2
- potassium measurement in, interpret with caution (letter), 753
- potassium measurement in, interpret with caution (reply), 753–754
- practice, treatment for maternal supraventricular tachyarrhythmia in pregnant patients in (letter), 1702–1704
- predicting tularemia with clinical, laboratory and demographical findings in, 218–221
- prediction of en-route complications during interfacility transport by outcome predictive scores in, 877–882
- predictive performance of qSOFA for mortality and ICU admission in patients with infection at, 1788–1793
- prehospital and hospital costs of emergency care for frequent patients in, 459–463
- prescribing practices, rescheduling of hydrocodone changed, 2388–2391
- presentation for American football injuries, tackling causes and costs of, 1198–1204
- presentations of coccidioidomycosis in (letter), 906–910
- as primary source of hospital admission for older (but not younger) adults, 943–947
- procalcitonin-guided therapy for initiation of antibiotics in, 1286–1293
- prognostic significance of C5a2 on polymorphonuclear neutrophil and C5a2intra/C5a2 ratio level for early sepsis in, 2084–2089
- propofol administration for procedural sedation in (letter), 1675–1676
- protective lung mechanical ventilation in (letter), 1900
- psychiatric patients, prolonged length of stay in, multivariable predictive model, 133–139
- qualitative patient interviews conducted within, 1169–1170
- reducing use requires interventions at multiple levels (letter), 1175–1176
- reliability of electrolyte and metabolite results measured by blood gas analyzer in, 419–424
- removal of pharyngeal wire bristle foreign body, a case report, 1328.e5–1328.e6
- resource-limited, ultrasound-guided pediatric forearm fracture reductions in, 40–44
- resuscitating tracheostomy patient in, 1148–1155
- revisits
- and outpatient follow-up, impact of concierges on (letter), 2445–2446
 - risk of, and cognitive health, in underserved older adults, 1973–1976
- routine tests and automated urinalysis in patients with suspected urinary tract infection at, 1528–1534
- seventy-two-hour antibiotic retrieval from, reporting the strength of association (letter), 1909–1910
- shared decision making in, 1668–1672
- Streptococcus gallolyticus (bovis)*, rare presentation of meningitis in, a case report, 677.e1
- Swyer-James-Macleod syndrome in, a case report, 1329.e3–1329.e4
- time to disposition, for patients with nontraumatic shock, impact of point-of-care ultrasonography on, 1022–1030
- transitional care clinics for follow-up and primary care linkage for patients discharged from, 1230–1235
- trauma patients, twenty-four-hour packed red blood cell requirement is strongest independent prognostic marker of mortality in, 1121–1124
- trends in hospital closures nationwide and across Medicaid expansion, 2006–2013, 1262–1264
- triage, of cardiology patients, direct cardiologist involvement in (letter), 325–326
- triage pain protocol reduces time to receiving analgesics in patients with painful conditions, 2362–2366
- triage sepsis alert and sepsis protocol lower times to fluids and antibiotics in, 1–9
- ultrasound-guided posterior tibial nerve blocks for calcaneal fracture analgesia, a case report, 1183.e1–1183.e3

- ultrasound-guided supracondylar radial nerve block for Colles fractures in (letter), 1718–1720
- underuse of information about timing and triggers in diagnosing dizziness in (letter), 1687–1689
- unusual presentation of disseminated Mycobacterium tuberculosis in, a case report, 1915.e5–1915.e7
- urban
- dedicated palliative care nurse improves access to palliative care and hospice services in (letter), 2440–2441
 - high-impact HCV testing for injection drug users in, 1108–1111
 - large, HCV infection, in 1945–1965 birth cohort (baby boomers) in, 697–701
 - with rapid HIV screening program, evaluation of hidden HIV infections in, 180–184
- of urban academic hospital, outcome of patients receiving CPR in, 1595–1599
- US
- and ambulatory care clinics with STI from 2001 to 2010, retrospective cross-sectional study of patients treated in, 1808–1811
 - changing epidemiology and management of infectious diseases in, 1059–1065
 - and French, treatment of acute renal colic in, simulated cases and real cases in acute pain management, 1955–1958
 - in US, 2002 to 2013, epidemiological study of surfing injuries presenting to, 1491–1496
 - use of WhatsApp application for orthopedic consultations in (letter), 1305–1307
 - utility of bolus intravenous nicardipine for hypertensive emergencies in, a case report, 2250.e1–2250.e3
 - utility of thyroid testing in (letter), 2029–2030
 - value of point-of-care ultrasound for detecting nail bed injury in, 1850–1854
 - variation in advanced imaging for pediatric patients with abdominal pain discharged from, 2320–2325
 - ventilation in, attention must be paid (letter), 1901
 - video laryngoscope as new standard of care in trauma (letter), 1313–1314
 - visit, for atrial fibrillation, hospital-level variation and predictors of admission after, 2094–2100
- visits
- frailty defined by SHARE-FI and adverse outcomes after (letter), 2443–2445
 - for patients with woman's health-related problems, enrollment with primary care provider does not preclude, 266–268
 - 5-year comparison by homeless and nonhomeless patients, 805–808
 - wait times, relationship between racial disparities illness severity, 10–15
- Emergency department (ED), CYP2C19 drug-drug and drug-gene interactions in patients in, 245–249
- Emergency department (ED) posterior reversible encephalopathy syndrome after bevacizumab treatment presenting as chest pain and headache in, a case report, 1916.e1–1916.e2
- Emergency department (ED) seventy-two-hour antibiotic retrieval from, randomized controlled trial of discharge instructional modality, 999–1005
- Emergency management, of ventricular assist devices, 1294–1301
- Emergency medical service (EMS)
- regional, impact of freestanding ED on, 1342–1346
- Emergency medical services (EMS)
- absence of gasping upon arrival, indicate need for endotracheal intubation (letter), 329–330
 - call for out-of-hospital cardiac arrest patients, favorable neurological outcomes by early epinephrine administration within 19 minutes after, 2284–2290
 - dispatcher, decreases time to defibrillation in public venue, randomized controlled trial, 590–593
 - investigation of oxidant and antioxidant levels in patients with acute stroke in, 2379–2383
 - practice, childbirth in (letter), 1888
 - predicts shorter door-to-balloon time and smaller infarct size, STEMI notification by, 1610–1613
 - psychiatric, development and testing of procedures for violence screening and suicide risk stratification on, 499–504
 - self-termination of ventricular fibrillation during transport by, a case report, 940.e1–940.e3
 - specialist availability in, contributions of response times and use of ad hoc coverage in New York State, 687–693
 - vs private vehicle, outcomes of NSTEMI patients transported by, 531–535
- Emergency medicine
- approach, inborn errors of metabolism (letter), 317–318
 - approach to abdominal vascular graft complications, 2014–2017
 - approach to transplant complications, 2200–2208
 - China's research status in, 15-year survey of literature, 392–397
 - Keutel syndrome, rare cause of dyspnea in, a case report, 935.e3–935.e5
 - milestone levels, interrater agreement of (letter), 1677–1679
 - nurses in simulated Crotalinae envenomation, variability of limb measurements performed by (letter), 1683–1685
- pharmacist
- impact on antibiotic dosing adjustment, 980–984
 - impact on time to thrombolysis in acute ischemic stroke, 1997–1999
 - prehospital, use and implementation of SOPs and checklists in, 2432–2439
 - tranexamic acid is used to treat upper gastrointestinal bleeds in (letter), 1892–1893
- Emergency physicians
- comparing with anesthesiologist, attitudes toward difficult airway management (letter), 754–755
 - comparison with on-call radiologists, CT interpretations in multiply injured patients, 2331–2335
 - diagnosis and treatment of uncontrolled chronic hypertension, barriers to (letter), 2241–2242
 - focused cardiac ultrasound improves diagnosis of ascending aortic dissection, 486–492
 - health and wellness among (letter), 1715–1716
 - involuntary admissions, role of alcohol intoxication in psychiatrists' recension of (letter), 2226–2227
 - lack of awareness among, of potentially inappropriate medications in elderly patients at EDs (letter), 1888–1890
 - performed transesophageal echocardiography for extracorporeal life support vascular cannula placement, 1637–1639
 - point-of-care ultrasonography at ED maximizes patient confidence in, 657–659
 - sonographers, can identify and inject regions of trigeminal nerve foramina in human cadaveric model (letter), 312–314
 - survey regarding emergency contraception (letter), 1176–1178
- Emergency response training, efficacy of simulation application SimMon for (letter), 927–928
- Emergency Severity Index, level 3, shock index, modified and age shock index for prediction of mortality in, 2079–2083
- Emergency transport, of stroke suspects in rural state, 1640–1644
- Emphysema, retropharyngeal, and pneumomediastinum, rare complication of influenza A and asthma in adult, a case report, 338.e1–338.e2
- EMS, Emergency medical services (EMS). *see* Emergency medical service (EMS)
- Endocarditis
- infective, with initial presentation of visual disturbances, a case report, 2052.e5–2052.e7
 - and septic emboli, Hampton's hump in patient with, a case report, 942.e1–942.e3
- Endotracheal intubation
- absence of gasping upon EMS arrival indicate need for (letter), 329–330

- with access to patient through back seat of car, paramedics are able to perform, randomized crossover manikin study (letter), 1161–1163
- blind, via LMA Fastrach, firefighters are able to perform (letter), 2458
- bougie-assisted, overcoming difficulties of (letter), 111–112
- comparison of, performed with 3 devices by paramedics wearing chemical, biological, radiological, and nuclear personal protective equipment (letter), 1902–1903
- during continuous chest compressions, ability of paramedics to perform, randomized cadaver study comparing Pentax AWS and Macintosh laryngoscopes, 1835–1839
- difficult, retrograde intubation more successful than direct laryngoscopic technique in, 2384–2387
- emergency, under fluoroscopy guidance for patients with acute dyspnea or asphyxia, 2177–2181
- during manual inline cervical stabilization performed by nurses (letter), 2456–2457
- need to use elevated sniffing position for, 928
- pediatric, gold rules for (letter), 1711–1712
- performed by nurses, comparison of GlideScope and Macintosh laryngoscopes for (letter), 2041
- prehospital, and survival, after out-of-hospital cardiac arrest, results from the Korean nationwide registry, 128–132
- with tactical fiberoptic imaging systems, 664–665
- using ETTI during simulated resuscitation, comparison of exchange of laryngeal mask airway and lgel for (letter), 106–107
- when using ETView tube, use of semirigid stylet increase efficacy of (letter), 1908–1909
- Endotracheal tube
- placement
 - tracheal ultrasonography and ultrasonographic lung sliding for confirming, speed and reliability, 953–956
 - using ultrasonography, paramedics are able to confirm (letter), 923–924
 - single lumen, VivaSight, comparison with Macintosh laryngoscope, for emergency intubation by experienced paramedics in standardized airway manikin with restricted access, randomized, crossover trial (letter), 929–930
 - VivaSight-SL, tracheal intubation with, by paramedics in cervical immobilized manikin (letter), 309–310
- Endovascular balloon occlusion
- of aorta
 - partial resuscitative, as hemorrhagic shock adjunct for ectopic pregnancy, a case report, 1917.e1–1917.e2
 - resuscitative, usefulness in detecting source of hemorrhage due to abdominal blunt trauma, a case report, 2057.e1–2057.e3
- End-stage renal disease, efficacy of loop diuretics in management of undocumented patients with, 1552–1555
- End-tidal carbon dioxide
- comparison with O₂, achieved by polypropylene and conventional polyvinyl chloride facemask in healthy individuals, 1157–1159
 - and occult injury, in trauma patients, 2146–2149
- Energy drinks
- in healthy population, cardiovascular responses to, 1205–1209
 - and realistic results, concordant with daily clinical practice, study on (letter), 1167
- Epiglottitis, acute, hypertrophic pachymeningitis of the craniocervical junction mimicking, a case report, 2467.e3–2467.e4
- Epinephrine
- early administration, within 19 minutes after EMS call for out-of-hospital cardiac arrest patients, favorable neurological outcomes by, 2284–2290
 - intraosseous administration, effects of proximal and distal routes on short-term resuscitative outcome measures in adult swine model of ventricular fibrillation, randomized controlled study, 49–53
 - use in ED patients with anaphylaxis, appropriateness of, 174–179
- Epipericardial fat necrosis, clinically mimicking pulmonary embolism, CT angiography findings, a case report, 2056.e5–2056.e6
- ERK. *see* Extracellular signal regulated kinase (ERK)
- Eschmann Tracheal Tube Introducer (ETTI), during simulated resuscitation, comparison of exchange of laryngeal mask airway and lgel for tracheal tube using (letter), 106–107
- Esophageal cancer, under radiotherapy, carotid blowout and cerebral gas embolism related to bidirectional carotid-esophageal fistula, serious complication of, a case report, 683.e5–683.e6
- Esophageal coin, suspected, a case report, 680.e3–680.e4
- Esophageal heat transfer device, introduction into therapeutic hypothermia protocol, 741–745
- Ethanol, poisoning, and associated hypoventilation, with doxapram, treatment of, a case report, 2253.e1–2253.e2
- Ethics, and regulatory challenges in advancing prehospital research, focus on sepsis, 623–625
- Ethyl chloride, topical, to reduce pain associated with venous catheterization randomized crossover trial, 845–850
- Ethyl pyruvate
- on inflammatory response and oxidative stress in multiple-organ dysfunction syndrome rats with severe burns, effects of insulin combined with, 2154–2158
 - and N-acetylcysteine
 - effect on ischemia-reperfusion injury in experimental model of ischemic stroke, 1804–1807
 - effects on ischemia-reperfusion injury in experimental electrical burn model, 1217–1224
- ETTI. *see* Eschmann Tracheal Tube Introducer
- ETView SL, Airtraq, and Macintosh laryngoscopes, comparison for face-to-face tracheal intubation, randomized crossover manikin trial (letter), 1893–1894
- ETView tube
- use of semirigid stylet increase efficacy of endotracheal intubation when using (letter), 1908–1909
 - without Macintosh laryngoscope, novice physicians intubate with (letter), 2242–2243
- Eunatremia, comparison of serum sodium levels measured by blood gas analyzer and biochemistry autoanalyzer in patients with, 1473–1479
- Exercise, stretching, ligamentum flavum hematoma due to, a case report, 2058.e3–2058.e6
- Extracellular signal regulated kinase (ERK) inhibitor, and green tea polyphenols, benefits of respective and combined use on survival and neurologic outcomes in cardiac arrest rats induced by ventricular fibrillation, 570–575
- Extracorporeal life support
- failure due to refractory ionized hypocalcemia during cardiac arrest, a case report, 2251.e1–2251.e2
 - vascular cannula placement, emergency physician-performed transesophageal echocardiography for, 1637–1639
- Extracorporeal membrane oxygenation (ECMO)
- and hemodialysis, combined use in case of contrast-induced biphasic anaphylactic shock, a case report, 1919.e1–1919.e2
 - preparation, simultaneous oropharyngeal and surgical, triple setup airway, a case report, 2468.e5–2468.e7
 - and salvage thrombolysis, for massive pulmonary embolism during distal femur fracture surgery, a case report, 1189.e3–1189.e5
- Eye diameter, optic nerve sheath diameter, CT measurement, quantitative evaluation of intracranial pressure by, 2336–2342
- Facemask, polypropylene and conventional polyvinyl chloride, in healthy individuals, comparison of O₂ and EtCO₂ achieved by, 1157–1159
- Facial nerve paralysis, bilateral, in outpatient setting, a case report, 1320.e3–1320.e4
- Facial paralysis, bilateral, in non-Hodgkin lymphoma during remission period, rare but important condition to be investigated, a case report, 2463.e5–2463.e7

- Fall
 in elderly patients, serious conditions for ED, 1394–1399
 mechanical, 582–585
- Febrile neutropenia, provider familiarity with specialty society guidelines for risk stratification and management of patients with (letter), 1704–1705
- Feedback
 device, impact on chest compression quality during extended manikin CPR, randomized crossover study, 1754–1760
 real-time, effect of, very brief training for laypeople in hands-only CPR, 993–998
- Femoral fracture, distal, surgery, salvage thrombolysis and ECMO for massive pulmonary embolism during, a case report, 1189.e3–1189.e5
- Femoral shaft fractures, after injury, intravenous morphine titration as rapid and efficient analgesia for adult patients with, 2107–2111
- Fever
 and flank pain, with renal vein thrombosis, unusual initial presentation of systemic lupus erythematosus, a case report, 937.e1–937.e2
 high, postpartum, and imitators of HELLP syndrome (letter), 2236–2237
 or risk factors, meningococcal sepsis caused by septic arthritis of a lumbar facet joint in patient without, a case report, 762.e1–762.e2
 and rash, acute generalized exanthematous pustulosis, uncommon cause of, a case report, 681.e1–681.e2
 rat bite, caused by kiss, a case report, 1190.e3–1190.e4
- Fiberoptic imaging system, tactical, endotracheal intubation with, 664–665
- Fibrinogen, to albumin ratio, and severity of coronary artery disease in patients with STEMI, relationship between, 1037–1042
- Fibromuscular dysplasia, presenting as hemorrhagic shock due to spontaneous rupture of right gastroepiploic artery aneurysm, a case report, 677.e3–677.e5
- Firefighters
 ability to recognize carbon monoxide poisoning (letter), 1710
 able to perform intraosseous access and start fluid resuscitation in anaphylactic patient (letter), 1707–1708
 able to recognize ventricular fibrillation (letter), 1885–1886
 are able to perform blind endotracheal intubation via LMA Fastrach (letter), 2458
- First aid, caregivers, skill decay or maintenance between tourniquet uses among, exploration in manikin model (letter), 1897–1899
- Flexible pressure sensor, could correctly measure depth of chest compression on mattress, 899–902
- Fluid balance, positive, as major predictor of clinical outcome of patients with sepsis/septic shock after ICU discharge, 2122–2126
- Fluid infusion, rate, affects rapidity of mean arterial pressure, restoration during controlled hemorrhage, 1743–1749
- Fluids, and antibiotics, in ED, triage sepsis alert and sepsis protocol lower times to, 1–9
- Fluorescein, novel technique to diagnose parotid duct injuries at bedside using, 308–309
- Fluoroscopy, guidance for patients with acute dyspnea or asphyxia, emergency endotracheal intubation under, 2177–2181
- Food impaction, esophageal, pediatric, point-of-care ultrasonography in detection of, a case report, 763.e1–763.e3
- Football injury, American, tackling causes and costs of ED presentation for, 1198–1204
- Foot fracture, postsplinting x-rays of, 1625–1626
- Forearm nerve block
 ultrasound-guided, for hand blast injuries, case series and multidisciplinary protocol (letter), 1895–1897
 ultrasound-guided, vs landmark-based wrist blocks for hand anesthesia in healthy volunteers, 730–734
- Foreign body
 impaction, esophageal, effectiveness of glucagon in relieving, 1049–1052
 soft tissue, ultrasound for evaluation of, before and after addition of fluid to surrounding interstitial space in cadaveric model, 1779–1782
 wooden, computed tomography settings for optimal detection of (letter), 2237–2238
- Fracture
 among 1323 fracture patients from 2010 Yushu earthquake, China, profile and procedures for, 2132–2139
 boxer
 anatomical challenges in ultrasound-guided ulnar nerve block for (letter), 1903–1904
 ultrasound-guided ulnar nerve block for (letter), 1726–1727
 calcaneal, analgesia, ED ultrasound-guided posterior tibial nerve blocks for, a case report, 1183.e1–1183.e3
 Colles, in ED, ultrasound-guided supracondylar radial nerve block for (letter), 1718–1720
 distal femur, surgery, salvage thrombolysis and ECMO for massive pulmonary embolism during, a case report, 1189.e3–1189.e5
 distal radius
 reduction using looped-stockinette technique (letter), 2038–2040
 supracondylar ultrasound-guided nerve block for reduction of, novel technique in ED (letter), 1172
 femoral shaft, after injury, intravenous morphine titration as rapid and efficient analgesia for adult patients with, 2107–2111
 finger, accuracy of bedside ultrasonography for diagnosis of, 809–812
 forearm, pediatric, ultrasound-guided, reductions in resource-limited ED, 40–44
 hip, intracapsular and extracapsular, ultrasound-guided nerve blocks for (erratum), 1921
 metacarpal
 effectiveness of bedside point-of-care ultrasonography in diagnosis and management of, contribution for management of metacarpal fractures (letter), 674
 role of ultrasonography in diagnosis of, 1868–1871
 of nondisplaced hand, wrist, ankle, and foot fractures, postsplinting x-rays of, 1625–1626
 patella, accuracy of point-of-care ultrasound as diagnostic tool for, 1576–1578
 phalanx, accuracy of bedside ultrasonography for diagnosis of (letter), 1698
 sternal
 and delayed cardiac tamponade due to severe blunt chest trauma, a case report, 758.e1–758.e3
- Frailty, defined by SHARE-FI and adverse outcomes after ED visit (letter), 2443–2445
- Fresh frozen plasma
 permissive for systemic inflammatory response syndrome, infection, and sepsis in multiple-injured patients, 1480–1485
- Fuchs syndrome, rare and unique variant of Stevens-Johnson Syndrome (letter), 910
- Gadolinium-based contrast agent anaphylaxis, induced acute abdominal pain, a case report, 1737.e1–1737.e2
- Gastroepiploic artery aneurysm, right, spontaneous rupture, fibromuscular dysplasia presenting as hemorrhagic shock due to, a case report, 677.e3–677.e5
- Gastrointestinal bleeding
 Dieulafoy lesion, little known sleeping giant of, a case report, 2464.e3–2464.e5
 upper, tranexamic acid is used to treat, in emergency medicine (letter), 1892–1893
- Gastrointestinal injury, sustained in 2-hit porcine model of septic shock, ulinastatin ameliorates, 1497–1504
- GCS. *see* Glasgow Coma Scale (GCS)
- Gender, and age group, bystander cardiopulmonary resuscitation training experience and self-efficacy of, 1331–1337
- Geriatric patients
 with acute ischemic infarct, bradycardia caused by intravenous nicardipine in, a case report, 761.e1–761.e2

- adults, noncontrast abdomen/pelvis computed tomographic scan in evaluation of (letter), 2230–2232
- chest pain, in ED observation unit, prospective evaluation of outcomes among, 207–211
- at EDs, lack of awareness among emergency physicians of potentially inappropriate medications in (letter), 1888–1890
- fall, serious conditions for ED, 1394–1399
- presenting to ED, characteristics and outcomes for acute heart failure in, 2159–2166
- with pulmonary embolism, spectrum of disturbances in cerebral function in, 651
- trauma, who required massive transfusion, bleeding sites in, comparison with younger patients, 123–127
- Giant bladder, (10,500 mL in volume) in ED, a case report, 1327.e1–1327.e2
- Glasgow Coma Scale (GCS), used as prognostic factor in unconscious patients following cardiac arrest in prehospital situations (letter), 1178–1179
- Glenohumeral dislocation, posterior, sling and swathe modification for immobilization of, 668–670
- GlideScope, comparison with Macintosh laryngoscope, for endotracheal intubation performed by nurses (letter), 2041
- Gloves, chemical protective, using skin protective lotion IB1 as substitute for, 1986–1990
- Glucagon, effectiveness in relieving esophageal foreign body impaction, 1049–1052
- Glucose, level, of 5 mg/dL (0.2 mmol/L), patient with clear consciousness even with, a case report, 941.e3–941.e4
- GONAIIS method, new autoreduction method for anterior shoulder dislocation, a case report, 120.e5–120.e7
- Gray matter to white matter ratio, in hypoxic and non-hypoxic cardiac arrest with non-cardiac etiology, prognostic value of, 1583–1588
- Green tea, and cardiac arrest (editorial), 576–577
- Green tea polyphenols, and ERK inhibitor, benefits of respective and combined use on survival and neurologic outcomes in cardiac arrest rats induced by ventricular fibrillation, 570–575
- Gum elastic bougie
- utility for tracheal intubation during chest compressions in manikin, randomized crossover trial, 54–56
 - utilization to facilitate single lung intubation, 2408–2410
- Hamman syndrome, pneumomediastinum combined with hyperosmolar hyperglycemic state, a case report, 2058.e1–2058.e2
- Hampton's hump, in patient with endocarditis and septic emboli, a case report, 942.e1–942.e3
- Hand dominance, in intravenous drug using patients does not affect peripheral venous access sites identified by ultrasound, 1088–1091
- Hand fracture, nondisplaced, postsplinting x-rays of, 1625–1626
- Hanging, neurologic outcome of comatose survivors after, 1467–1472
- HCG. *see* Human chorionic gonadotropin (HCG)
- HCV. *see* Hepatitis C virus (HCV)
- Headache
- and chest pain, posterior reversible encephalopathy syndrome after bevacizumab treatment presenting to ED as, a case report, 1916.e1–1916.e2
 - treatment in carbon monoxide poisoning patients with, 2140–2145
- Head injury
- minor
 - comparison of near-infrared spectroscopy and head CT interpretations of ED patients with, 1364–1368
 - and normal initial CT scan, outcomes of warfarinized patients with, 75–78
- Head position, effect on success rate of blind intubation using intubating supraglottic airway devices, 1193–1197
- Head trauma
- infantile, accuracy of history of injury obtained from caregiver in, 1863–1867
 - minor, stuttering after, a case report, 685.e3–685.e4
- Health care
- associated acute pyelonephritis, associated with inappropriate empiric antibiotic therapy in ED, 1415–1420
 - waste, prevention of, 1661–1662
- Health centers, for disease control guidelines for ED patients with sexually transmitted diseases, compliance with (letter), 1727–1729
- Health literacy, and patient activation, assessing, in ED, 93–96
- Heart
- infected, ventriculoseptal abscess and intracardiac fistulization, a case report, 678.e1–678.e3
 - metastasis to, rare life-threatening condition, a case report, 1912.e3–1912.e4
 - stopping honey, not just Turkish honey, a case report, 1915.e1–1915.e3
- Heart failure
- acute
 - in elderly patients presenting to ED, characteristics and outcomes for, 2159–2166
 - lung ultrasound, diagnostic and prognostic tool at every step in pathway of care for, 656–657
 - plasma bioactive adrenomedullin as prognostic biomarker in, 257–262
 - point-of-care strain echocardiography in (letter), 2234–2236
 - resolution of sonographic B-lines as measure of pulmonary decongestion in, 1129–1132
 - usefulness of thrombolysis in MI risk index in, 2351–2355
 - chronic, long-term use of beta-blockers in (letter), 931
 - congestive, diagnosis in emergency care setting, Brest score, development of clinical prediction score for, 2277–2283
 - left ventricular, acute, with pulmonary infection, and pulmonary infection, clinical application of rapid B-line score with lung ultrasonography in differentiating between, 278–281
- HELLP syndrome
- late postpartum, over 10 days after delivery, a case report, 2258.e1–2258.e3
 - postpartum high fever and imitators of (letter), 2236–2237
- Hematoma
- intracranial, perihematomal edema diameter as predictive indicator of 30-day mortality of (letter), 1701–1702
 - intramuscular, of pectoralis major after CPR, hemorrhagic shock caused by, a case report, 682.e5–682.e6
 - ligamentum flavum, due to stretching exercise, a case report, 2058.e3–2058.e6
- Hemidiaphragmatic paresis, associated with brachial plexus blockade, a case report, 1733.e5–1733.e7
- Hemodialysis, and ECMO, combined use in case of contrast-induced biphasic anaphylactic shock, a case report, 1919.e1–1919.e2
- Hemodynamics
- challenges in traumatic subarachnoid hemorrhage complicated by cerebral vasospasm (letter), 904–906
 - phenotypes, presenting in ED patients, with confirmed sepsis, 2291–2297
- Hemolytic anemia, incidence and patterns in acute dapson overdose, 366–369
- Hemopericardium, fatal, after intravenous rt-PA for acute ischemic stroke, a case report, 2462.e5–2462.e6
- Hemoperitoneum, after blunt abdominal trauma 27 years after splenectomy, a case report, 1325.e3–1325.e4
- Hemophagocytic lymphohistiocytosis, in ED, a case report, 2057.e5–2057.e8
- Hemorrhage
- alveolar, diffuse, related to spider bite, ANCA–positive vasculitis with, a case report, 339.e3–339.e4
 - controlled, infusion rate of fluid affects rapidity of mean arterial pressure restoration during, 1743–1749
 - due to abdominal blunt trauma, usefulness of resuscitative endovascular balloon occlusion of aorta, in detecting source of, a case report, 2057.e1–2057.e3
 - subarachnoid
 - ECG abnormalities predict neurogenic pulmonary edema in patients with, 79–82

- retroclival, posttraumatic, a case report, 2257.e3–2257.e5
- Hemorrhagic shock
 adjunct, for ectopic pregnancy, partial resuscitative endovascular balloon occlusion of aorta as, a case report, 1917.e1–1917.e2
 caused by intramuscular hematoma of pectoralis major after CPR, a case report, 682.e5–682.e6
 due to leech bite with tranexamic acid, treatment of patient in, a case report, 2253.e3–2253.e4
 due to spontaneous rupture of right gastroepiploic artery aneurysm, fibromuscular dysplasia presenting as, a case report, 677.e3–677.e5
 pyruvate alleviates lipid peroxidation and multiple-organ dysfunction in rats with, 525–530
 pyruvate alleviates lipid peroxidation and multiple-organ dysfunction in rats with (erratum), 1330
 and trauma arrests, need for abdominal only CPR in treatment of (letter), 1156–1157
- Hemorrhagic transformation, during emergency evaluation, differentiating contrast staining after acute ischemic stroke from, a case report, 2255.e5–2255.e6
- Hemostasis, successful, and reversal of highly elevated PT/INR after dabigatran etexilate use in patient with acute kidney injury, a case report, 758.e5–758.e6
- Hemothorax, traumatic, successful transcatheter lumbar arterial embolization of, a case report, 1188.e3–1188.e5
- Hepatic dysfunction, lactate clearance and mortality in septic patients with, 1011–1015
- Hepatic failure, fulminant, requiring liver transplantation due to accidental alcohol ingestion in 7-month-old baby, a case report, 2466.e1–2466.e2
- Hepatitis C virus (HCV), testing, high-impact, for injection drug users in urban ED, 1108–1111
- Hepatitis C virus (HCV) infection, in 1945–1965 birth cohort (baby boomers) in large urban ED, 697–701
- Hernia, pulmonary, posttraumatic, in child, a case report, 759.e3–759.e4
- Herniation, spontaneous, mimicking acute aortic dissection, a case report, 2053.e1–2053.e2
- Heroin, -dependent patients, seeking asthma care, characteristics in ED, 895–898
- HGM. *see* Human menopausal gonadotropin (HGM)
- Hiccups, persistent, followed by acute respiratory failure, a case report, 1181.e3–1181.e4
- Hill-Sachs lesion, engaging, large, traumatic posterior shoulder dislocation with, splinting technique, a case report, 682.e1–682.e3
- Hip fracture
 intracapsular and extracapsular, ultrasound-guided nerve blocks for, 586–589
 orthopedic literature 2015, 1663–1667
- HIV. *see* Human immunodeficiency virus (HIV)
- HIV-1. *see* Human immunodeficiency virus-1 (HIV-1)
- Hoarseness, and dysphagia, aortic dissection manifesting as, Ortner's syndrome, a case report, 1185.e1–1185.e3
- Hoigne syndrome, little-known adverse effect of lidocaine, a case report, 679.e3–679.e4
- Home infusion therapy, intravenous, instituted from 24-hour clinical decision unit for patients with cellulitis, 1273–1275
- Homelessness
 and ED use
 myths and facts (letter), 307
 myths and facts (reply), 307–308
 and nonhomeless patients, 5-year comparison of ED visits by, 805–808
- Hospital admission
 in ED, as quality measure, critical analysis of unplanned ICU transfer within 48 hours from, 1505–1510
 after ED visits for atrial fibrillation, hospital-level variation and predictors of, 2094–2100
 inpatient, prior authorization for child and adolescent psychiatric patients deemed to be in need of (letter), 915–916
 intensive care unit, routine, necessary for small subdural hemorrhages, 521–524
 for older (but not younger) adults, ED as primary source of, 943–947
 predict mortality in ED patients with sepsis, high immunoglobulin E values at, 1589–1594
 for suicidal ideation in ED, demographic characteristics of individuals (letter), 1174–1175
 time, increased door to, is associated with prolonged throughput for ED patients discharged home, 1783–1787
- Hospital discharge
 from ED, transitional care clinics for follow-up and primary care linkage for patients, 1230–1235
 from ED, variation in advanced imaging for pediatric patients with abdominal pain, 2320–2325
 ICU, positive fluid balance as major predictor of clinical outcome of patients with sepsis/septic shock after, 2122–2126
 increased door to admission time is associated with prolonged throughput for ED patients, 1783–1787
 instruction delivery/modality and antibiotic retrieval (letter), 1910
- HPV. *see* Human papillomavirus (HPV)
- Human chorionic gonadotropin (HCG), and HMG, concurrent ovarian hyperstimulation syndrome with perforated appendicitis following induction ovulation with, a case report, 2051.e5–2051.e6
- Human immunodeficiency virus (HIV), screening program, rapid, evaluation of hidden HIV infections in urban ED with, 180–184
- Human immunodeficiency virus-1 (HIV-1), acute, seroconversion, in pediatric patient, rhabdomyolysis as presenting feature of, a case report, 760.e3–760.e5
- Human immunodeficiency virus (HIV) infection, hidden, evaluation in urban ED with rapid HIV screening program, 180–184
- Human menopausal gonadotropin (HGM), and HCG, concurrent ovarian hyperstimulation syndrome with perforated appendicitis following induction ovulation with, a case report, 2051.e5–2051.e6
- Human papillomavirus (HPV), vaccination, and cervical cancer screening, in ED, model for implementation of (letter), 648
- Humerus fracture, Holstein Lewis, a case report, 345.e5–345.e6
- Humid continental climate, influence of air pollutants on appearance of acute myocardial infarction in region with (letter), 1679–1682
- Hunger, self-assessment among ED patients with abdominal pain, lack of association with disease severity, 104–106
- Hydration therapy, blood urea nitrogen/creatinine-based, 3 months after acute ischemic stroke, favorable outcome of, 2414–2418
- Hydrocodone, rescheduling of, changed ED prescribing practices, 2388–2391
- Hydrothorax, presenting, predicts failure of needle aspiration in primary spontaneous pneumothorax, 1075–1079
- 5-hydroxyindoleacetic acid, urinary, spot, is not ideal diagnostic test for acute appendicitis, 1750–1753
- Hyperammonemia
 cerebral edema induced by, a case report, 2461.e3–2461.e5
 clinical, therapeutic, and educational aspects (letter), 1894
 isolated, due to urea cycle disorder, rare cause of postpartum coma (letter), 1894–1895
 isolated, due to urea cycle disorder rare cause of postpartum coma, a case report, 1324.e3–1324.e4
 pediatric occult methanol intoxication with, a case report, 343.e5–343.e6
- Hyperbaric oxygen therapy, and brain ischemia, efficacy in acute CO intoxication, dual-energy CT shows, a case report, 1327.e3–1327.e4
- Hypercalcemia, -induced pancreatitis presenting with acute ST-elevations mimicking myocardial infarction, a case report, 1187.e1–1187.e2
- Hyperemesis, synthetic cannabinoid, resulting in rhabdomyolysis and acute renal failure, a case report, 765.e1–765.e2
- Hyperglycemia
 and hyperosmolality, comprehensive interpretation on clinical outcomes among ischemic stroke patients, 2343–2350
 seriously, continuous renal replacement therapy, new treatment of, a case report, 2469.e3–2469.e4

- Hyperkalemia**
 moderate to severe, agreement of serum potassium measured by blood gas and biochemistry analyzer in patients with, 794–797
 succinylcholine, and rocuronium (letter), 1705–1706
 unanticipated, succinylcholine in ED patients with (letter), 1706–1707
 unanticipated, succinylcholine in ED patient with, a case report, 2054.e3–2054.e5
- Hypnatremia**, comparison of serum sodium levels measured by blood gas analyzer and biochemistry autoanalyzer in patients with, 1473–1479
- Hyperosmolality**, and hyperglycemia, comprehensive interpretation on clinical outcomes among ischemic stroke patients, 2343–2350
- Hyperosmolar hyperglycemic state**
 bilateral above knee amputations, uncommon complication of, a case report, 341.e1–341.e2
 combined with pneumomediastinum, Hamman syndrome, a case report, 2058.e1–2058.e2
- Hypersensitivity reaction**, after administration of rivaroxaban (Xarelto), a case report, 1325.e1–1325.e2
- Hypertension**
 brainstem encephalopathy, mimicking central pontine myelinolysis, potential pitfall, a case report, 1918.e5–1918.e7
 chronic, uncontrolled, barriers to emergency physician diagnosis and treatment of (letter), 2241–2242
 emergencies in ED, utility of bolus intravenous nicardipine for, a case report, 2250.e1–2250.e3
 severe, in ED, effect of predischARGE blood pressure on follow-up outcomes in patients with, 834–839
- Hyperthermia**
 and severe rhabdomyolysis
 from synthetic cannabinoids, a case report, 121.e1–121.e2
 from synthetic cannabinoids (letter), 1690–1691
- Hypertrophic pachymeningitis**, of craniocervical junction mimicking acute epiglottitis, a case report, 2467.e3–2467.e4
- Hyperventilation**
 and intubated patients, cerebral oximetry with blood volume index and capnography in, 1102–1107
 mitigating, during CPR, 643–646
- Hypoalbuminemia**, predicts clinical outcome in patients with type B acute aortic dissection after endovascular therapy, 1369–1372
- Hypocalcemia**, refractory ionized, during cardiac arrest, failure of extracorporeal life support due to, a case report, 2251.e1–2251.e2
- Hyponatremia**, comparison of serum sodium levels measured by blood gas analyzer and biochemistry autoanalyzer in patients with, 1473–1479
- Hypotension**
 and bradycardia, after synthetic cannabinoid use, a case report, 2055.e1–2055.e2
 effects of acute intravenous amiodarone overdose in swine model, Intralipid™ administration attenuates, 1389–1393
 4-factor PCC-associated, a case report, 2059.e1–2059.e2
 propofol-induced, ultrasound measurement of IVC collapse predicts, 1125–1128
- Hypothermia**
 and cyclosporine A, protective effects on neuronal mitochondria in rat asphyxial cardiac arrest model, 1080–1085
 severe, and cardiac arrest successfully treated without external mechanical circulatory support, a case report, 1913.e5–1913.e6
 therapeutic
 for children surviving near-hanging, a case report, 1738.e5–1738.e7
 early neurologic examination is not reliable for prognostication in post-cardiac arrest patients who undergo, 975–979
 and extracorporeal CPR, for out-of-hospital cardiac arrest in patient with long QT syndrome, a case report, 1321.e1–1321.e3
 prehospital, after out-of-hospital cardiac arrest, 2209–2216
 protocol, introduction of esophageal heat transfer device into, 741–745
 after resuscitation, impact of body mass index on patient survival after, 722–725
 unexpected survival of sudden cardiac arrest patient with large multiple brain infarction after, a case report, 2462.e1–2462.e3
- Hypoventilation**, associated, and ethanol poisoning, with doxapram, treatment of, a case report, 2253.e1–2253.e2
- Hypovolemic shock**
 axillary artery laceration after anterior shoulder dislocation presenting as, a case report, 1739.e1–1739.e4
 vascular access technique chosen during (letter), 1886–1887
- Hypoxia**, brain, on CT after avalanche-induced cardiac arrest, Serum potassium concentration predicts, 856–860
- IABP**. *see* Intraaortic balloon pumping (IABP)
- Ibuprofen**, -induced thrombotic thrombocytopenic purpura, a case report, 942.e5–942.e7
- Icare**, vs Tono-Pen in ED (letter), 670–673
- Ice-fishing**, in United States, injury patterns and outcomes of, 1258–1261
- ICU**. *see* Intensive care unit (ICU)
- I-gel**
 as alternative airway tool for difficult airway in severely injured patients, a case report, 340.e1–340.e4
 and laryngeal mask airway, comparison for tracheal tube using ETTI during simulated resuscitation (letter), 106–107
- Illicit drug user**, purpuric skin eruption in, a case report, 1321.e5–1321.e6
- Illness**, severity, and racial disparities, in ED wait times, relationship between, 10–15
- Imaging**
 advanced, for pediatric patients with abdominal pain discharged from ED, variation in, 2320–2325
 unexpected findings, and shortness of breath, a case report, 937.e3–937.e4
 upper extremity deep vein thrombosis, diagnostic limitations of, Paget-Schroetter syndrome, a case report, 683.e1–683.e3
- Immune thrombocytopenic purpura (ITP)**, successful thrombolytic therapy for STEMI in patient with, a case report, 345.e1–345.e3
- Immunocompetence**, patient without rash, Varicella-zoster virus encephalitis in, a case report, 2257.e1–2257.e2
- Immunocompromised patient**, potentially, CSF/blood glucose ratio and bacterial meningitis without pleocytosis in (letter), 922
- Immunoglobulin E**, high values, at admission predict mortality in ED patients with sepsis, 1589–1594
- Impending paradoxical embolus**, bedside diagnosis in ED, a case report, 1917.e3–1917.e5
- Inborn errors of metabolism**, emergency medicine approach (letter), 317–318
- Infants**. *see* Pediatrics
- Infarct size**, smaller, STEMI notification by EMS predicts shorter door-to-balloon time and, 1610–1613
- Infection**
 cross-, copper-coated thermometer for prevention of, preliminary results (letter), 653–656
 at ED, predictive performance of qSOFA for mortality and ICU admission in patients with, 1788–1793
 heart, ventriculoseptal abscess and intracardiac fistulization, a case report, 678.e1–678.e3
 methicillin-resistant *Staphylococcus aureus*, after pediatric scald burn, toxic shock syndrome due to, a case report, 1735.e1–1735.e2
 and mortality, early predictors of, patients with cirrhosis in ED, 25–29
Pasteurella multocida, salvage of upper extremity in aggressive case of, a case report, 2470.e1–2470.e3
 patients, with DIC, recombinant human soluble thrombomodulin and short-term mortality of, 1876–1882
 pulmonary, and pulmonary infection with acute left ventricular heart failure, clinical application of rapid B-line score with lung ultrasonography in differentiating between, 278–281

- Streptococcus pneumoniae*, prognostic factors in adults, 202–206
systemic inflammatory response syndrome, and sepsis, in multiple-injured patients, fresh frozen plasma is permissive for, 1480–1485
in US EDs, changing epidemiology and management of, 1059–1065
- Inferior vena cava (IVC)
aorta, sonographic, cross-sectional area index for evaluation of dehydration in children, 1840–1844
collapse, predicts propofol-induced hypotension, ultrasound measurement of, 1125–1128
filters, duplicate, a case report, 115.e1–115.e4
spontaneous rupture, deadly right flank pain, a case report, 2050.e1–2050.e3
ultrasound, NT-proBNP, and bioelectrical impedance analysis for diagnosis of acute decompensated CHF, 1817–1822
- Inflammasome, NLRP3, activation, artemisinin protects mice against burn sepsis through inhibiting, 772–777
- Inflammation, biomarkers predicting prognosis in patients with acute dyspnea, 370–374
- Inflammatory response, and oxidative stress, in multiple-organ dysfunction syndrome rats with severe burns, effects of insulin combined with ethyl pyruvate on, 2154–2158
- Influenza, H1N1, presenting as severe acute pancreatitis and multiorgan dysfunction, a case report, 1911.e1–1911.e2
- Influenza A, and asthma in adult, retropharyngeal air and pneumomediastinum, rare complication of, a case report, 338.e1–338.e2
- Information technology, ED, proactive approach to, case of pediatric procedural sedation (letter), 1901–1902
- Infusion rate, fluid, affects rapidity of mean arterial pressure, restoration during controlled hemorrhage, 1743–1749
- Injection drug users
hand dominance in, does not affect peripheral venous access sites identified by ultrasound, 1088–1091
in urban ED, high-impact HCV testing for, 1108–1111
- Injury
American football, tackling causes and costs of ED presentation for, 1198–1204
ankle, orthopedic literature 2015, 1663–1667
elbow, comparison of bedside point-of-care ultrasound and computed tomography in, 2186–2190
electronic cigarette explosion, a case report, 2252.e1–2252.e3
finger tip, traumatic, closed, in patients with artificial nails, removal of UV gel and acrylic nails, 335–337
firearm, to head and neck, cervical spine immobilization may be of value following, 726–729
gastrointestinal, sustained in 2-hit porcine model of septic shock, ulinastatin ameliorates, 1497–1504
hand blast, ultrasound-guided forearm nerve blocks for, case series and multidisciplinary protocol (letter), 1895–1897
head, minor, comparison of near-infrared spectroscopy and head CT interpretations of ED patients with, 1364–1368
history of, accuracy obtained from caregiver in infantile head trauma, 1863–1867
intravenous morphine titration as rapid and efficient analgesia for adult patients with femoral shaft fractures after, 2107–2111
by mechanism, age effects on case fatality rates of patients, 515–520
moped, increase in, requiring emergency care, 2000–2002
multiple, fresh frozen plasma is permissive for systemic inflammatory response syndrome, infection, and sepsis in patients with, 1480–1485
multiple, patients, CT interpretations in, comparison of emergency physicians and on-call radiologists, 2331–2335
muscle, after tourniquet-induced ischemia-reperfusion injury in rats, postischemic conditioning does not reduce, 2065–2069
nail bed, in ED, value of point-of-care ultrasound for detecting, 1850–1854
ocult, and end-tidal carbon dioxide, in trauma patients, 2146–2149
of parotid duct, novel technique to diagnose, at bedside using fluorescein, 308–309
patients in ED, association between acute alcohol consumption and discharge against medical advice of, 464–468
patterns, and crash types, in MotoGP class, elite motorcycle racing, 1872–1875
patterns and outcomes of ice-fishing in United States, 1258–1261
severely, patients, I-gel as alternative airway tool for difficult airway in, a case report, 340.e1–340.e4
soft tissue, but also for improved laryngoscopy and tracheal intubation, lip pulling method for prevention of (letter), 1694
surfing, presenting to US EDs 2002 to 2013, epidemiological study of, 1491–1496
TASER penetrating ocular, emergent diagnosis and management of, a case report, 1740.e3–1740.e5
violent, recurrent, magnitude, risk factors, and opportunities for intervention from statewide analysis, 1823–1830
- INR. *see* International normalized ratio (INR)
- Insulin, combined with ethyl pyruvate, effects on inflammatory response and oxidative stress in multiple-organ dysfunction syndrome rats with severe burns, 2154–2158
- Intensive care unit (ICU)
admission
and mortality, in patients with infection at ED, predictive performance of qSOFA for, 1788–1793
routine, necessary for small subdural hemorrhages, 521–524
discharge, positive fluid balance as major predictor of clinical outcome of patients with sepsis/septic shock after, 2122–2126
unplanned transfer within 48 hours from ED admission as quality measure, critical analysis of, 1505–1510
- Interfacility transport, by outcome predictive scores in ED, prediction of en-route complications during, 877–882
- Interhospital transfer
first-door-to-balloon time delay in STEMI patients undergoing, 767–771
and point of injury, cumulative effective radiation dose received by blunt trauma patients arriving to military level I trauma center from, 2397–2401
- International normalized ratio (INR), PT, highly elevated, after dabigatran etexilate use in patient with acute kidney injury, successful hemostasis and reversal of, a case report, 758.e5–758.e6
- Interview, qualitative, patient, conducted within ED, 1169–1170
- Intestinal necrosis, baseline disease is more important predictor than CT findings in patients with acute mesenteric ischemia, 2261–2265
- Intoxication
alcohol, role in psychiatrists' recension of emergency physicians' involuntary admissions (letter), 2226–2227
CO, acute, dual-energy CT shows brain ischemia and hyperbaric oxygen therapy efficacy in, a case report, 1327.e3–1327.e4
with digoxin, Intravenous lipid emulsion prolongs survival in rats, 1112–1116
methanol, occult, pediatric, with hyperammonemia, a case report, 343.e5–343.e6
trauma patient, respiratory depression in, opioids to blame, 250–253
water, acute, drinking to near death—, leading to neurogenic stunned myocardium, a case report, 119.e3–119.e4
- Intraaortic balloon pumping (IABP), support, use for oozing-type cardiac rupture after acute MI, a case report, 120.e1–120.e3
- Intracompartmental pressure, comparison of 2 available methods with Bland-Altman analysis for measuring, 1765–1771
- Intracranial aneurysm, comorbid, acute ischemic stroke is dangerous to treat by thrombolytic therapy in patients with, 636–642
- Intracranial hematoma
perihematomal edema diameter as predictive indicator of 30-day mortality of (letter), 1701–1702
prognostic value of perihematomal edema area at initial ED presentation in patients with, 1241–1246
- Intracranial hemorrhage

- incidence and clinical features of, causing out-of-hospital cardiac arrest, 2326–2330
- traumatic, discontinuing oral anti-coagulants in patient with cardiac valve prosthesis after, a case report, 1736.e5–1736.e7
- Intracranial pressure
- in ED, bedside sonographic measurement of optic nerve sheath diameter as predictor of, 963–967
 - quantitative evaluation by optic nerve sheath diameter/eye diameter CT measurement, 2336–2342
- Intralipid™, administration attenuates hypotensive effects of acute intravenous amiodarone overdose in swine model, 1389–1393
- Intranasal administration, of cocaine abuse in young man, cardiac arrest most dangerous complication of intranasal cocaine abuse in young man, a case report, 1731.e5–1731.e7
- Intraosseous
- access
 - junior doctors trained to use (letter), 107
 - nurses' knowledge and attitudes toward (letter), 1724
 - in obese patient, assessing need for extended needle length, 1831–1834
 - in pediatric patients, nurses' attitudes and beliefs concerning (letter), 1890
 - for reversal of rivaroxaban, PCC administration through, a case report, 685.e1–685.e2
 - and start fluid resuscitation, in anaphylactic patient, firefighters able to perform (letter), 1707–1708
 - access devices
 - comparison of NIO and EZ-IO, in adult patients under resuscitation performed by paramedics, randomized crossover manikin trial (letter), 1166–1167
 - as method of vascular access during CPR (letter), 321–322
 - epinephrine, administration, effects of proximal and distal routes on short-term resuscitative outcome measures in adult swine model of ventricular fibrillation, randomized controlled study, 49–53
 - and intravenous administration, tibial, of vasopressin on kinetics and survivability in cardiac arrest, effects of, 429–432
 - new route to life in patients with circulatory shock (letter), 922–923
- Intrathecal pump, malfunction, and magnetic fields, a case report, 115.e5–115.e6
- Intravascular access
- comparison of two techniques when using CBRN-PPE, randomized crossover manikin trial (letter), 1170–1172
 - method, to choose during CPR (letter), 1160
- Intravenous access
- advanced, technique choices, pain scores, and failure rates in local registry, 553–557
 - difficult, ED patients, randomized controlled trial assessing use of ultrasound for nurse-performed intravenous placement in, 1950–1954
 - patients, difficult, prospective evaluation of ultrasound-guided short catheter placement in internal jugular veins of, 578–581
- Intravenous administration
- of amiodarone, acute overdose in swine model, Intralipid™ administration attenuates hypotensive effects of, 1389–1393
 - of dexketoprofen, ketoprofen gel improves low back pain in addition to, 1458–1461
 - home infusion therapy instituted from 24-hour clinical decision unit for patients with cellulitis, 1273–1275
 - and intraosseous, tibial, of vasopressin on kinetics and survivability in cardiac arrest, effects of, 429–432
 - of ketamine, optimal dosing, for procedural sedation in children in ED, 1347–1353
 - of lipid emulsion, in resuscitation of cocaine-induced cardiovascular arrest in rat model, 1452–1454
 - lipid emulsion therapy, use in acute methamphetamine toxicity, a case report, 1732.e3–1732.e4
 - of morphine
 - titration as rapid and efficient analgesia for adult patients with femoral shaft fractures after injury, 2107–2111
 - vs acupuncture, in management of acute pain in ED, 2112–2116
 - nicardipine, bolus, for hypertensive emergencies in ED, utility of, a case report, 2250.e1–2250.e3
- Intravenous fluids, for migraine, post hoc analysis of clinical trial data, 713–716
- Intravenous line placement, nurse-performed, in difficult access ED patients, randomized controlled trial assessing use of ultrasound for, 1950–1954
- Intubation
- blind, through supraglottic airway laryngopharyngeal tube with biohazard suit (letter), 2040
 - blind, using intubating supraglottic airway devices, effect of head position on success rate of, 1193–1197
 - body mass index associated with inappropriate tidal volumes in adults in ED (letter), 1682–1683
 - child, laryngoscope use for (letter), 1729–1730
 - delayed sequence, with ketamine in 2 critically ill children, a case report, 1190.e1–1190.e2
 - emergency, by experienced paramedics in standardized airway manikin with restricted access, comparison of VivaSight single lumen endotracheal tube and the Macintosh laryngoscope for, randomized, crossover trial (letter), 929–930
 - endotracheal
 - blind, via LMA Fastrach, firefighters are able to perform (letter), 2458
 - comparison of, performed with 3 devices by paramedics wearing chemical, biological, radiological, and nuclear personal protective equipment (letter), 1902–1903
 - during continuous chest compressions, ability of paramedics to perform, randomized cadaver study comparing Pentax AWS and Macintosh laryngoscopes, 1835–1839
 - difficult, retrograde intubation more successful than direct laryngoscopic technique in, 2384–2387
 - emergency, under fluoroscopy guidance for patients with acute dyspnea or asphyxia, 2177–2181
 - during manual inline cervical stabilization performed by nurses (letter), 2456–2457
 - pediatric, gold rules for (letter), 1711–1712
 - performed by nurses, comparison of GlideScope and Macintosh laryngoscopes for (letter), 2041
 - when using ETVIEW tube, use of semirigid stylet increase efficacy of (letter), 1908–1909
 - face-to-face, in manikin, randomized crossover trial comparing the C-MAC and Macintosh laryngoscopes for (letter), 920–922
 - fiberoptic, or video tube for trauma patient intubation, method to choose, randomized crossover manikin trial (letter), 751–753
 - first attempt, of trauma patient, Venner A.P. Advance video laryngoscope improve success of (letter), 315–316
 - and hyperventilated patients, cerebral oximetry with blood volume index and capnography in, 1102–1107
 - lung, single, utilization of gum elastic bougie to facilitate, 2408–2410
 - novice physicians, with ETVIEW tube without Macintosh laryngoscope (letter), 2242–2243
 - oro-tracheal, by nurses during resuscitation, comparison of Macintosh and Intubrite laryngoscopes for (letter), 1724–1725
 - in patients with inline manual neck stabilization, comparison of Airtraq and Miller laryngoscopes for (letter), 1721
 - protocols, drug-facilitated, prehospital, statewide, variability in United States (letter), 2459–2460
 - rapid sequence, in trauma resuscitation, integration of point-of-care sonography during (letter), 330
 - rapid sequence, medication dosing, accuracy in obese patients intubated in ED, 2423–2425
 - rapid sequence, utility of noninvasive nasal positive pressure ventilators for, optimizing oxygenation during, 1627–1630

- retrograde, more successful than direct laryngoscopic technique in difficult endotracheal intubation, 2384–2387
- during simulated resuscitation by novice physicians, comparison of AirTraq Avant® vs Macintosh laryngoscope for (letter), 1708–1709
- tracheal
- child, during CPR, McGrath-MAC vs Macintosh laryngoscopes for (letter), 2453
 - child, during CPR, videolaryngoscopy or direct laryngoscopy for (letter), 2453
 - child, during resuscitation by paramedics, comparison of McGrath-MAC and Macintosh laryngoscopes for, 1338–1341
 - face-to-face, comparison of ETView SL, Airtraq, and Macintosh laryngoscopes for, randomized crossover manikin trial (letter), 1893–1894
 - and laryngoscopy, lip pulling method for prevention of soft tissue injury but also for (letter), 1694
 - with Macintosh laryngoscope with and without chest compressions, performed by nurses (letter), 2448–2449
- Intubrite, laryngoscope, and Macintosh, for orotracheal intubation by nurses during resuscitation, comparison of (letter), 1724–1725
- Ischemia-modified albumin
- in cardiovascular disease (letter), 1699
 - pentraxin 3, and myeloperoxidase in predicting cardiac damage in acute carbon monoxide poisoning, a case report, 1927–1930
- Ischemia-reperfusion injury
- in experimental electrical burn model, effects of N-acetylcysteine and ethyl pyruvate on, 1217–1224
 - in experimental model of ischemic stroke, effect of ethyl pyruvate and N-acetylcysteine on, 1804–1807
 - in rats, postischemic conditioning does not reduce muscle injury after tourniquet-induced, 2065–2069
- Ischemic stroke
- acute
 - bradycardia caused by intravenous nicardipine in elderly patient with, a case report, 761.e1–761.e2
 - fatal hemopericardium after intravenous rt-PA for, a case report, 2462.e5–2462.e6
 - favorable outcome of blood urea nitrogen/creatinine-based hydration therapy 3 months after, 2414–2418
 - from hemorrhagic transformation during emergency evaluation, differentiating contrast staining after, a case report, 2255.e5–2255.e6
 - impact of emergency medicine pharmacist on time to thrombolysis in, 1997–1999
 - is dangerous to treat by thrombolytic therapy in patients with comorbid intracranial aneurysms, 636–642
 - serum albumin level is associated with recurrence of, 1812–1816
 - effect of ethyl pyruvate and N-acetylcysteine on ischemia-reperfusion injury in experimental model of, 1804–1807
 - patients, comprehensive interpretation of hyperglycemia and hyperosmolality on clinical outcomes among, 2343–2350
- ITP. *see* Immune thrombocytopenic purpura (ITP)
- IVC. *see* Inferior vena cava (IVC)
- Jamshidi intraosseous access devices, and Bone Injection Gun, by paramedics with and without chemical-biological-radiological-nuclear personal protective equipment, comparison of (letter), 1307–1308
- Jet injection, vs needle infiltration, in lumbar puncture, comparison of local anesthetic effect of lidocaine by, 1225–1229
- Karadeniz wrist rules, for the use of radiography in wrist trauma, 2074–2078
- Kayexalate, moving away from (letter), 1716
- Ketamine
- with delayed sequence intubation, in 2 critically ill children, a case report, 1190.e1–1190.e2
 - intravenous
 - finding optimal initial dose for pediatric procedural sedation is still challenging (letter), 1892
 - for procedural sedation in children in ED, optimal dosing of, 1347–1353
 - propofol combination (ketofol) vs propofol for procedural sedation and analgesia, 558–569
- Ketofol, ketamine-propofol combination vs propofol for procedural sedation and analgesia, 558–569
- Ketoprofen gel, improves low back pain in addition to IV dexamethasone, 1458–1461
- Keutel syndrome, rare cause of dyspnea in emergency medicine, a case report, 935.e3–935.e5
- Kidney failure, in US ED, mesoamerican nephropathy a novel case of, a case report, 1323.e5–1323.e6
- Kidney function, several parameters, correlation with clinical severity and prognosis of acute MI (letter), 1904–1907
- Kidney injury
- acute
 - and anuria, an uncommon case of bilateral synchronous ureteric calculi, a case report, 2256.e1–2256.e2
 - successful hemostasis and reversal of highly elevated PT/INR after dabigatran etexilate use in patient with, a case report, 758.e5–758.e6
- Kienbock disease, avascular necrosis of lunate bone, a case report, 1185.e5–1185.e6
- Knee dislocation, irreducible posterolateral, diagnosis, arthroscopic view, and treatment of, a case report, 2256.e3–2256.e4
- Kounis hypersensitivity, -associated ACS, and Takotsubo cardiomyopathy, myocardial injury through serum troponin I and echocardiography in anaphylaxis (letter), 650–651
- Lactate
- blood, concentration after cardiac arrest resulting from myocardial infarction and outcome (letter), 1311–1313
 - clearance and mortality in septic patients with hepatic dysfunction, 1011–1015
 - high plasma concentration, quantitative analysis in ED patients after alcohol intake, 825–829
 - levels, non-critical, prognostic role of, for in-hospital survival time among ED patients with sepsis, 170–173
 - plasma, and blood alcohol levels (letter), 1725–1726
 - point-of-care, and PSP-S, assessment of Bayesian clinical decision model integrating, acute care diagnostics collaboration, 193–196
 - serum
 - admission, predicts mortality in aneurysmal subarachnoid hemorrhage, 708–712
 - for predicting bacteremia in female patients with acute pyelonephritis, diagnostic performance of initial, 1359–1363
- Lactate dehydrogenase, predictive role in sepsis patients, 2167–2171
- Lactic acidosis, beta agonist-induced, evidence-based approach to critical question, 666–668
- Lance-Adams syndrome, peculiar case of, a case report, 342.e5–342.e7
- Laparoscopy, management for spontaneous rupture of left gastroepiploic vessel after forceful retching, a case report, 759.e1–759.e2
- Large-bowel disease, presenting as small-bowel obstruction is associated with poor prognosis, 477–479
- Laryngeal mask airway, and Igel, comparison for tracheal tube using ETTI during simulated resuscitation (letter), 106–107
- Laryngeal mask airway (LMA) Classic, for anesthetized, paralyzed children weighing 20 kg, application of (letter), 1697–1698
- Laryngeal tube
- or masks, used for out-of-hospital cardiac arrest patients (letter), 749–750
 - safety of (letter), 310–311
- Laryngoscope
- C-MAC and Macintosh, for face-to-face intubation in manikin, randomized crossover trial comparing (letter), 920–922

- ETView SL, Airtraq, and Macintosh, comparison for face-to-face tracheal intubation, randomized crossover manikin trial (letter), 1893–1894
- GlideScope and Macintosh, comparison of, for endotracheal intubation performed by nurses (letter), 2041
- Macintosh
 comparison with VivaSight, single lumen endotracheal tube, for emergency intubation by experienced paramedics in standardized airway manikin with restricted access, randomized, crossover trial (letter), 929–930
 double-lumen tube tracheal intubation in manikin model using the VivaSight Double Lumen, randomized controlled comparison with (letter), 103–104
 ETVIEW tube without, novice physicians intubate with (letter), 2242–2243
 and Intubrite, for orotracheal intubation by nurses during resuscitation, comparison of (letter), 1724–1725
 vs AirTraq Avant®, for intubation during simulated resuscitation by novice physicians, comparison of (letter), 1708–1709
 with and without chest compressions, performed by nurses, tracheal intubation with (letter), 2448–2449
- McGrath-MAC
 and Macintosh, for child tracheal intubation during resuscitation by paramedics, comparison of, 1338–1341
 vs Macintosh, for child tracheal intubation during CPR (letter), 2453
 Miller, and Airtraq, for intubation in patients with inline manual neck stabilization, comparison of (letter), 1721
 Pentax AWS and Macintosh, ability of paramedics to perform endotracheal intubation during continuous chest compressions, randomized cadaver study comparing, 1835–1839
 use for child intubation (letter), 1729–1730
- Laryngoscopy
 direct
 comparison in different head and neck positions in edentulous patients, 1855–1858
 in difficult endotracheal intubation, retrograde intubation more successful than, 2384–2387
 or videolaryngoscopy, for child tracheal intubation during CPR (letter), 2453
 and tracheal intubation, lip pulling method for prevention of soft tissue injury but also for (letter), 1694
 video, investigation of force received at upper teeth by (letter), 1695
- Lead perforation, late, presenting as acute shoulder pain, a case report, 2255.e1–2255.e3
- Leech bite, with tranexamic acid, treatment of patient in hemorrhagic shock due to, a case report, 2253.e3–2253.e4
- Left bundle-branch block, reversible, due to carbon monoxide poisoning, a case report, 342.e1–342.e3
- Lemierre syndrome, efficacy of non-vitamin K antagonist oral anticoagulants for, a case report, 2059.e3–2059.e5
- Length of stay
 in ED
 during mass casualty incident, factors associated with, 1462–1466
 variation across classifications of clinical condition and patient discharge disposition, 83–87
 prolonged, in ED psychiatric patients, multivariable predictive model, 133–139
- Lethargy, acute, in young woman due to latent disseminated cancer mimicking bacterial meningitis, diagnostic pitfall, a case report, 2050.e5–2050.e7
- Levamisole, -induced vasculitis, in an illicit drug user, a case report, 1321.e5–1321.e6
- Levosimendan, role in management of subarachnoid hemorrhage, 298–306
- Liberatory vertigo, new prognostic factor for repositioning maneuvers, 1548–1551
- Lidocaine
 as adjuvant to morphine improve pain relief in patients presenting to ED with acute renal colic, 443–448
- Hoigne syndrome, little-known adverse effect of, a case report, 679.e3–679.e4
 by jet injection vs needle infiltration in lumbar puncture, comparison of local anesthetic effect of, 1225–1229
- LifeLine ARM device, during simulated CPR, mechanical chest compression with (letter), 917
- LifeVac, assessment on human cadaver with complete airway obstruction (letter), 1673–1674
- Limb, measurements performed by emergency medicine nurses in simulated *Crotalinae* envenomation, variability of (letter), 1683–1685
- Linear transducer, high-frequency, improves detection of intrauterine pregnancy in first-trimester ultrasonography, 288–291
- Lipase
 and amylase
 normal, acute pancreatitis with, a case report, 2254.e3–2254.e6
 normal, with acute pancreatitis, a case report, 940.e5–940.e7
 normal, and scrotal swelling, rare presentation of acute pancreatitis, a case report, 763.e5–763.e6
- Lipid emulsion
 intravenous
 prolongs survival in rats intoxicated with digoxin, 1112–1116
 in resuscitation of cocaine-induced cardiovascular arrest in rat model, 1452–1454
- Lipid emulsion therapy, intravenous, use in acute methamphetamine toxicity, a case report, 1732.e3–1732.e4
- Lipid peroxidation
 and multiple-organ dysfunction, in rats with hemorrhagic shock, pyruvate alleviates, 525–530
 and multiple-organ dysfunction, in rats with hemorrhagic shock, pyruvate alleviates (erratum), 1330
- Lip pulling method, for prevention of soft tissue injury but also for improved laryngoscopy and tracheal intubation (letter), 1694
- Lipschütz ulcer, cause of misdiagnosis when suspecting child abuse, a case report, 1326.e1–1326.e2
- Liver injury, blunt traumatic, thrombotic thrombocytopenic purpura after, a case report, 939.e3–939.e4
- Liver metastasis, catastrophic pneumoperitoneum in patient with perforated colorectal cancer with, a case report, 344.e3–344.e5
- Liver transplantation, due to accidental alcohol ingestion in 7-month-old baby, fulminant hepatic failure requiring, a case report, 2466.e1–2466.e2
- LMA Classic. *see* Laryngeal mask airway (LMA) Classic
- LMA Fastrach, firefighters are able to perform endotracheal intubation via (letter), 2458
- Local anesthesia, effect of lidocaine by jet injection vs needle infiltration in lumbar puncture, comparison of, 1225–1229
- Long axis in-plane, vs short axis out-of-plane techniques, effects during ultrasound-guided vascular access, 778–783
- Long QT syndrome, extracorporeal CPR and therapeutic hypothermia for out-of-hospital cardiac arrest in patient with, a case report, 1321.e1–1321.e3
- Loop diuretics, efficacy in management of undocumented patients with end-stage renal disease, 1552–1555
- Looped-stockinette technique, distal radius fracture reduction using (letter), 2038–2040
- Lotion, IB1, skin protective, using as substitute for chemical protective gloves, 1986–1990
- Low back pain, in addition to IV dexketoprofen, ketoprofen gel improves, 1458–1461
- Low-energy mechanism, isolated dorsal dislocations of talonavicular and calcaneocuboid articulations (Chopart joints) from, a case report, 1733.e1–1733.e4

- L-selectin, and Mac-1, neutrophil adhesion molecules, inhaled corticosteroids increase blood neutrophil count by decreasing expression of, 1977–1981
- Luc abscess, extraordinary complication of acute otitis media, a case report, 117.e1–117.e3
- Lumbar puncture
 comparison of local anesthetic effect of lidocaine by jet injection vs needle infiltration in, 1225–1229
 ultrasonography-guided, vs landmark, perceived difficulty and success rate in novice study group, randomized crossover trial (letter), 2037–2038
- Lung injury, acute, transfusion-related, after blood products transfusions, risk of (letter), 1674–1675
- Lung protective ventilation, and mechanical ventilator settings, in ED, 1446–1451
- Mac-1, and L-selectin, neutrophil adhesion molecules, inhaled corticosteroids increase blood neutrophil count by decreasing expression of, 1977–1981
- Macintosh
 and C-MAC laryngoscopes, for face-to-face intubation in manikin, randomized crossover trial comparing (letter), 920–922
 comparison with GlideScope laryngoscope, for endotracheal intubation performed by nurses (letter), 2041
 direct laryngoscopy, comparison with video laryngoscopy, for airway management, 1886
- ETView SL, and Airtraq laryngoscopes, comparison for face-to-face tracheal intubation, randomized crossover manikin trial (letter), 1893–1894
- laryngoscope
 comparison with VivaSight, single lumen endotracheal tube, for emergency intubation by experienced paramedics in standardized airway manikin with restricted access, randomized, crossover trial (letter), 929–930
 double-lumen tube tracheal intubation in manikin model using the VivaSight Double Lumen, randomized controlled comparison with (letter), 103–104
- ETView tube without, novice physicians intubate with (letter), 2242–2243
 and Intubrite, for orotracheal intubation by nurses during resuscitation, comparison of (letter), 1724–1725
 vs AirTraq Avant®, for intubation during simulated resuscitation by novice physicians, comparison of (letter), 1708–1709
 with and without chest compressions, performed by nurses, tracheal intubation with (letter), 2448–2449
 and McGrath-MAC laryngoscopes, for child tracheal intubation during resuscitation by paramedics, comparison of, 1338–1341
 and Pentax AWS laryngoscopes, ability of paramedics to perform endotracheal intubation during continuous chest compressions, randomized cadaver study comparing, 1835–1839
 vs McGrath-MAC laryngoscopes, for child tracheal intubation during CPR (letter), 2453
- Macrocirculation, and microcirculation, during CPR, phosphodiesterase-5 inhibition improves, 162–166
- Magnesium sulfate
 beneficial action in bronchial asthma (letter), 1693–1694
 nebulized, effect in treatment of moderate to severe asthma attacks, randomized clinical trial, 883–886
- Magnetic field, and intrathecal pump malfunction, a case report, 115.e5–115.e6
- Magnetic resonance imaging (MRI)
 for pediatric burner and stinger syndrome, 1442–1445
 uncommon findings, and severe neurologic impairment, after carbon monoxide poisoning, a case report, 119.e1–119.e2
- Major surgery, atrial septal defect can mimic pulmonary thromboembolism after, a case report, 121.e3–121.e6
- Manual compression, on standard stretcher, versus mechanical compression on reducible stretcher, in small elevator, quality between, 1604–1609
- Mask
 bag-valve, with internal handle, single rescuer ventilation using, 1991–1996
 or laryngeal tubes, used for out-of-hospital cardiac arrest patients (letter), 749–750
- Mass casualty incident, factors associated with ED length of stay during, 1462–1466
- May-Thurner syndrome
 as cause of phlegmasia cerulea dolens, a case report, 1326.e3–1326.e4
 curious syndrome in ED, a case report, 1920.e1–1920.e3
- McGrath-MAC
 and Macintosh laryngoscopes, for child tracheal intubation during resuscitation by paramedics, comparison of, 1338–1341
 vs Macintosh laryngoscopes for child tracheal intubation during CPR (letter), 2453
- Mean arterial pressure, restoration, during controlled hemorrhage, infusion rate of fluid affects rapidity of, 1743–1749
- Mechanical compression, on reducible stretcher, versus manual compression on standard stretcher, in small elevator, quality between, 1604–1609
- Mechanical ventilation, lung, protective, in ED (letter), 1900
- Mechanical ventilator, settings, and lung protective ventilation in ED, 1446–1451
- Mediastinal mass, emergent radiation therapy as definite airway management for dyspnea with, a case report, 2258.e5–2258.e6
- Mediastinitis, poststernotomy, diagnosing, in ED, 618–622
- Medicaid
 beneficiaries, who continue to use ED, focus on Illinois Medical Home Network, 197–201
 expansion, 2006–2013, trends in hospital ED closures nationwide and across, 1262–1264
 and mental health disorders, with ED admissions for ambulatory care-sensitive condition conditions, association of, 820–824
- Medical center, quaternary, critical access hospital ED to, 1427–1430
- Medicare, beneficiaries with chest pain, increased observation services in, 16–19
- Medication
 controlled, prescribing, to patients who die of prescription drug abuse, 30–35
 dosing, rapid sequence intubation, accuracy in obese patients intubated in ED, 2423–2425
 overdose, oral, activated charcoal may not be necessary in all (letter), 319–321
 potentially inappropriate, in elderly patients at EDs, lack of awareness among emergency physicians of (letter), 1888–1890
- Mefloquine, -associated rhabdomyolysis, a case report, 2250.e5–2250.e6
- Meningitis
Plesiomonas shigelloides, in an adult in ED, a case report, 1329.e1–1329.e2
Streptococcus gallolyticus (bovis), rare presentation in ED, a case report, 677.e1
Streptococcus pneumoniae, without pleocytosis of CSF, a case report, 941.e1–941.e2
 validation of bacterial meningitis score in adults presenting to ED with, 1265–1267
- Meningoencephalitis, caused by septic arthritis of lumbar facet joint in patient without fever or risk factors, a case report, 762.e1–762.e2
- Mental foramen, and ED neuro-ultrasonography (letter), 2240–2241
- Mental health disorder, and Medicaid, with ED admissions for ambulatory care-sensitive condition conditions, association of, 820–824
- Mesoamerican nephropathy, a novel case of kidney failure in US ED, a case report, 1323.e5–1323.e6
- Metabolic acidosis
 anion gap
 normal, in salicylate overdose (letter), 2457–2458
 salicylate toxicity in absence of, a case report, 1328.e1–1328.e3

- Metabolite, and electrolyte results, measured by blood gas analyzer in ED, reliability of, 419–424
- Metacarpal fracture
effectiveness of bedside point-of-care ultrasonography in diagnosis and management of, contribution for management of metacarpal fractures (letter), 674
role of ultrasonography in diagnosis of, 1868–1871
- Metastasis, to heart, rare life-threatening condition, a case report, 1912.e3–1912.e4
- Metaxalone, therapeutic, associated with serotonin syndrome, in patient with cirrhosis, a case report, 346.e5–346.e6
- Methadone, overdose, causing acute cerebellitis and multi-organ damage, a case report, 343.e1–343.e3
- Methamphetamine, toxicity, acute, use of intravenous lipid emulsion therapy in, a case report, 1732.e3–1732.e4
- Methanol intoxication, occult, pediatric, with hyperammonemia, a case report, 343.e5–343.e6
- Methicillin-resistant *Staphylococcus aureus*, infection, after pediatric scald burn, toxic shock syndrome due to, a case report, 1735.e1–1735.e2
- Metoprolol in beta-blocker-naïve patients vs patients on chronic beta-blocker therapy, achieving ventricular rate control using, 606–608
- MHR. *see* Monocyte count to high-density lipoprotein ratio (MHR)
- MI. *see* Myocardial infarction (MI)
- Microcirculation, and macrocirculation, during CPR, phosphodiesterase-5 inhibition improves, 162–166
- Migraine, intravenous fluids for, post hoc analysis of clinical trial data, 713–716
- Miller, laryngoscope, and Airtraq, for intubation in patients with inline manual neck stabilization, comparison of (letter), 1721
- Misdiagnosis, of Lipschütz ulcer, when suspecting child abuse, a case report, 1326.e1–1326.e2
- Mitochondria, neuronal, in rat asphyxial cardiac arrest model, protective effects of cyclosporine A and hypothermia on, 1080–1085
- Monocyte count to high-density lipoprotein ratio (MHR), as new prognostic marker in patients with STEMI undergoing primary PCI, 240–244
- Monocyte to high density lipoprotein ratio, and no-reflow phenomenon, in patients with acute STEMI, relation between, 1542–1547
- Morbidity, or mortality, trauma patients with phencyclidine-positive urine drug screens increase, 1066–1068
- Morphine
improve pain relief in patients presenting to ED with acute renal colic, lidocaine as adjuvant to, 443–448
intravenous
titration, as rapid and efficient analgesia for adult patients with femoral shaft fractures after injury, 2107–2111
vs acupuncture, in management of acute pain in ED, 2112–2116
in setting of acute MI, pros and cons (letter), 746–748
- Mortality
in aneurysmal subarachnoid hemorrhage, admission serum lactate predicts, 708–712
association with triage time shock index, modified shock index, and age shock index, in emergency severity index level 2 patients, 63–68
considerable, spontaneous perforation of pyometra presenting as acute abdomen, rare condition with, a case report, 761.e3–761.e5
30-day, of intracranial hematoma, perihematoma edema diameter as predictive indicator of (letter), 1701–1702
in ED patients with sepsis, high immunoglobulin E values at admission predict, 1589–1594
in ED trauma patients, twenty-four-hour packed red blood cell requirement is strongest independent prognostic marker of, 1121–1124
in Emergency Severity Index level 3, shock index, modified and age shock index for prediction of, 2079–2083
and ICU admission, in patients with infection at ED, predictive performance of qSOFA for, 1788–1793
and infection, early predictors of, patients with cirrhosis in ED, 25–29
inhospital, due to pulmonary embolism in patients with Alzheimer disease, sex effect (letter), 665–666
and lactate clearance, in septic patients with hepatic dysfunction, 1011–1015
in low-risk cardiac patients with severe sepsis, elevated cardiac biomarkers are not associated with (letter), 1899–1900
or morbidity, trauma patients with phencyclidine-positive urine drug screens increase, 1066–1068
predictors in severe sepsis (letter), 1899
in sepsis, predictors of (letter), 2223–2224
short-term, and recombinant human soluble thrombomodulin, of infection patients with DIC, 1876–1882
- Mortality in Severe Sepsis in the Emergency Department (MISSED) score, usefulness in urban tertiary care hospital, 1117–1120
- Mounier-Kuhn syndrome, unusual cause of recurrent spontaneous pneumothorax, a case report, 122.e1–122.e2
- MRI. *see* Magnetic resonance imaging (MRI)
- Multidisciplinary team, for acute abdominal aortic aneurysm, 1519–1523
- Multi-organ damage, and acute cerebellitis, methadone overdose causing, a case report, 343.e1–343.e3
- Multiple-organ dysfunction
and lipid peroxidation, in rats with hemorrhagic shock, pyruvate alleviates, 525–530
and lipid peroxidation, in rats with hemorrhagic shock, pyruvate alleviates (erratum), 1330
and severe acute pancreatitis, H1N1 influenza presenting as, a case report, 1911.e1–1911.e2
- Multiple-organ dysfunction syndrome, in rats with severe burns, effects of insulin combined with ethyl pyruvate on inflammatory response and oxidative stress in, 2154–2158
- Multiple sclerosis, convergence spasm, very rare paroxysmal symptom in, a case report, 117.e5–117.e6
- Muscle hypertrophy, Paget-Schroetter syndrome secondary to thoracic outlet obstruction from, a case report, 1323.e1–1323.e3
- Muscle injury, after tourniquet-induced ischemia-reperfusion injury in rats, postischemic conditioning does not reduce, 2065–2069
- Muscle relaxant, effect on tracheal intubation with Pentax-AWS Airwayscope, prospective randomized controlled trial, 674–675
- Muscular fitness, as mediator of quality CPR, 1845–1849
- Mycobacterium tuberculosis*, disseminated, unusual presentation in ED, a case report, 1915.e5–1915.e7
- Myelinolysis, central pontine, hypertensive brainstem encephalopathy mimicking, potential pitfall, a case report, 1918.e5–1918.e7
- Myeloperoxidase, pentraxin 3, and ischemia-modified albumin, in predicting cardiac damage in acute carbon monoxide poisoning, 1927–1930
- Myocardial infarction (MI)
acute
in carbon monoxide poisoning, COHb levels predict long-term development of, 840–844
correlation of several kidney function parameters with clinical severity and prognosis of (letter) of, 1904–1907
effects of air pollution and weather conditions on incidence of, 449–454
emergency medical services annotated literature in review 2016, 2193–2199
inferior, new ST-segment algorithms to determine culprit artery location in, 1772–1778
morphine in setting of, pros and cons (letter), 746–748
in region with humid continental climate, influence of air pollutants on appearance of (letter), 1679–1682
straight back syndrome manifesting as, a case report, 1913.e1–1913.e3
after trauma, potency of PCI with transcatheter arterial embolization, a case report, 1186.e1–1186.e3
use of IABP support for oozing-type cardiac rupture after, a case report, 120.e1–120.e3

- anterior
 absent in early repolarization, terminal QRS distortion present in, 2182–2185
 impending, de Winter syndrome (letter), 2450–2451
 high lateral, South African flag sign, teaching tool for easier ECG recognition of (letter), 107–109
 hypercalcemia-induced pancreatitis presenting with acute ST-elevations mimicking, a case report, 1187.e1–1187.e2
 and outcome, blood lactate concentration after cardiac arrest resulting from (letter), 1311–1313
 paraoxanase as indicator of, and its utility in determining extension of ischemia, 45–48
 pseudo, celiac sprue presenting as, a case report, 1187.e3–1187.e5
 risk index, in acute heart failure, usefulness of thrombolysis in, 2351–2355
 type 2, and status epilepticus, a case report, 1735.e3–1735.e4
- Myocardial injury
 evaluation of, through serum troponin I and echocardiography in anaphylaxis, 140–144
 through serum troponin I and echocardiography in anaphylaxis, Takotsubo cardiomyopathy and Kounis hypersensitivity-associated ACS (letter), 650–651
- Myoglobin, higher serum level of, could predict more severity and poor outcome for patients with sepsis, 948–952
- N-acetylcysteine
 and ethyl pyruvate
 effect on ischemia-reperfusion injury in experimental model of ischemic stroke, 1804–1807
 effects on ischemia-reperfusion injury in experimental electrical burn model, 1217–1224
 investigation on contralateral testis tissue injury by experimental testicular torsion, 1069–1074
- Nasal cannula, high-flow, in ED for patients with solid malignancy, application of (letter), 2222–2223
- Nasal cannula therapy, high-flow, oxygen supply, as treatment in hypercapnic respiratory failure, a case report, 1914.e1–1914.e2
- Nasal positive pressure ventilators, noninvasive, for optimizing oxygenation during rapid sequence intubation, utility of, 1627–1630
- National Early Warning Score–Lactate (NEWS–L) score, validation in CAP patients, 536–541
- National Trauma Data Bank, suicide attempts presenting to, 1620–1624
- Near-hanging, therapeutic hypothermia for children survived from, a case report, 1738.e5–1738.e7
- Near-infrared spectroscopy, and head CT, interpretations of ED patients with minor head injury, comparison of, 1364–1368
- Neck stabilization, inline manual, comparison of Airtraq and Miller laryngoscopes for intubation in patients with (letter), 1721
- Needle aspiration
 in primary spontaneous pneumothorax, presenting hydrothorax predicts failure of, 1075–1079
 pulmonary contusion post, of primary spontaneous pneumothorax, a case report, 1734.e1–1734.e3
- Needle cricothyroidotomy, paramedics prepared to perform (letter), 1310–1311
- Needle infiltration, vs jet injection, in lumbar puncture, comparison of local anesthetic effect of lidocaine by, 1225–1229
- Needle thoracostomy, success rates, and chest wall thickness, determination at second and fifth intercostal spaces, cadaver-based study, 2310–2314
- Negative pressure wound therapy, for serious dog bites of extremities, prospective randomized trial, 1006–1010
- Nephropathy
 contrast, to adverse events in pulmonary emboli patients diagnosed with contrast CT, relation of, 1247–1250
 contrast-induced, related to adverse events in pulmonary embolism patients (letter), 2042–2043
 contrast-induced, related to adverse events in pulmonary embolism patients (reply), 2043–2044
 Neurogenic pulmonary edema, in patients with subarachnoid hemorrhage, ECG abnormalities predict, 79–82
 Neurogenic stunned myocardium, drinking to near death—acute water intoxication leading to, a case report, 119.e3–119.e4
 Neurologic examination, early, is not reliable for prognostication in post-cardiac arrest patients who undergo therapeutic hypothermia, 975–979
 Neurologic impairment, severe, and uncommon MRI findings, after carbon monoxide poisoning, a case report, 119.e1–119.e2
 Neutrophil count, by decreasing expression of neutrophil adhesion molecules Mac-1 and L-selectin, inhaled corticosteroids increase, 1977–1981
 Neutrophil-to-lymphocyte ratio, potential diagnostic marker for peptic ulcer perforation, 403–406
 NEWS–L. *see* National Early Warning Score–Lactate (NEWS–L)
 Nicardipine
 bolus intravenous, for hypertensive emergencies in ED, utility of, a case report, 2250.e1–2250.e3
 intravenous, in elderly patient with acute ischemic infarct, bradycardia caused by, a case report, 761.e1–761.e2
- Nitroglycerin
 corrected flow time, noninvasive ultrasound measure to detect preload reduction by, 1859–1862
 venous dilator, combined to clonidine, centrally acting antihypertensive, to handle severe pulmonary edema, a case report, 676.e5–676.e7
 Nitrous oxide, inhaled, use for analgesia in adult ED patients, 269–273
 NLRP3. *see* Nucleotide-binding domain and leucine-rich repeat containing family, pyrin containing 3 (NLRP3)
- Non-Hodgkin lymphoma, during remission period, bilateral facial paralysis in, rare but important condition to be investigated, a case report, 2463.e5–2463.e7
- Non-ST-segment elevation myocardial infarction (NSTEMI)
 association between prehospital transportation and clinical outcomes in patients with (letter), 1676–1677
 outcomes of patients by EMS vs private vehicle, 531–535
 by point-of-care echocardiography with speckle tracking, diagnosis of acute coronary occlusion in patients with, a case report, 1914.e3–1914.e6
 in teenage female desiring to lose weight, clenbuterol causing, a case report, 1739.e5–1739.e7
- NSTEMI. *see* Non-ST-segment elevation myocardial infarction (NSTEMI)
- N-terminal pro-B-type natriuretic peptide (NT-proBNP), IVC ultrasound, and bioelectrical impedance analysis for diagnosis of acute decompensated CHF, 1817–1822
- NT-proBNP. *see* N-terminal pro-B-type natriuretic peptide (NT-proBNP)
- Nucleotide-binding domain and leucine-rich repeat containing family, pyrin containing 3 (NLRP3), inflammasome activation, artemisinin protects mice against burn sepsis through inhibiting, 772–777
- Nurse
 attitudes and beliefs concerning intraosseous access in pediatric patients (letter), 1890
 comparison of GlideScope and Macintosh laryngoscopes for endotracheal intubation performed by (letter), 2041
 endotracheal intubation during manual inline cervical stabilization performed by (letter), 2456–2457
 knowledge and attitudes toward intraosseous access (letter), 1724
 palliative care, dedicated, improves access to palliative care and hospice services in urban ED (letter), 2440–2441
 -performed intravenous placement in difficult access ED patients, randomized controlled trial assessing use of ultrasound for, 1950–1954
 during resuscitation, comparison of Macintosh and Intubrite laryngoscopes for orotracheal intubation by (letter), 1724–1725
 in simulated *Crotalinae* envenomation, variability of limb measurements performed by emergency medicine (letter), 1683–1685
 tracheal intubation with Macintosh laryngoscope with and without chest compressions, performed by (letter), 2448–2449
 use, mechanical chest compression devices during CPR (letter), 2044–2045

- Nursery songs
 for infants and young children's anxiety, effects of, reporting mistake of effect measure of association (letter), 663
 for infants and young children's anxiety before and during head CT, randomized single-blinded clinical trial on effects of (letter), 663
- Nursing continuing education units, for ultrasound-guided intravenous education does not increase utilization, alteration of timing of (letter), 2248–2249
- OAR. *see* Ottawa Ankle Rules (OAR)
- Obesity
 intraosseous access in, assessing need for extended needle length, 1831–1834
 intubated in ED, accuracy of rapid sequence intubation medication dosing in, 2423–2425
 super, closed reduction of anterior shoulder dislocation in, a case report, 1181.e1–1181.e2
- Occult injury, and end-tidal carbon dioxide, in trauma patients, 2146–2149
- Ocular disease, use of ultrasound in (letter), 112
- Ocular injury, TASER penetrating, emergent diagnosis and management of, a case report, 1740.e3–1740.e5
- Omental infarction, unusual cause of abdominal pain in pregnancy, a case report, 1184.e3–1184.e4
- Omental torsion, mimicking perforated appendicitis in pediatric patient, emergency bedside sonography, a case report, 684.e3–684.e4
- Opioid
 to blame, respiratory depression in intoxicated trauma patient, 250–253
 deaths, shows startling findings, analysis of prescription patterns in (letter), 748–749
 prescribing, ED, is not associated with higher patient satisfaction scores (letter), 2032–2034
- Optic nerve, cysticercosis, retroorbital, a case report, 2461.e1–2461.e2
- Optic nerve sheath
 diameter, as predictor of intracranial pressure in ED, bedside sonographic measurement of, 963–967
 diameter, ultrasonography, knowledge, attitude, and practices of paramedics regarding, 1160–1161
 eye diameter, CT measurement, quantitative evaluation of intracranial pressure by, 2336–2342
- Optic sheath, posttraumatic pneumatization of, a case report, 1911.e3–1911.e4
- Orotracheal intubation, by nurses during resuscitation, comparison of Macintosh and Intubrite laryngoscopes for (letter), 1724–1725
- Orthopedic consultation, in ED, use of WhatsApp application for (letter), 1305–1307
- Ortner's syndrome, aortic dissection manifesting as dysphagia and hoarseness, a case report, 1185.e1–1185.e3
- Otitis media, acute, Luc abscess, extraordinary complication of, a case report, 117.e1–117.e3
- Ottawa Ankle Rules (OAR), impact of clinical decision support tool on adherence to, 412–418
- Outcomes
 adverse, and SHARE-FI, after ED visit, frailty defined by (letter), 2443–2445
 of 2005 and 2010 cardiopulmonary resuscitation guidelines for adults with in-hospital cardiac arrest, meta-analysis of, 1133–1139
 cardiovascular disease, in tertiary care centers in Japan (letter), 109–111
 and characteristics, for acute heart failure in elderly patients presenting to ED, 2159–2166
- clinical
 among ischemic stroke patients, comprehensive interpretation of hyperglycemia and hyperosmolality on, 2343–2350
 of patients with sepsis/septic shock after ICU discharge, positive fluid balance as major predictor of, 2122–2126
 in patients with type B acute aortic dissection after endovascular therapy, hypoalbuminemia predicts, 1369–1372
 and prehospital transportation, in patients with non-STEMI, association between (letter), 1676–1677
 and sepsis, C1 inhibitor in (letter), 1179–1180
 of ED visit, comparison with risk factors, nonsuicidal self-injury and suicide attempts among ED patients older than 50 years, 1016–1021
 of emergency department placement of transvenous pacemakers, 1411–1414
 favorable, of blood urea nitrogen/creatinine-based hydration therapy, 3 months after acute ischemic stroke, 2414–2418
 follow-up, in patients with severe hypertension in ED, effect of predischARGE blood pressure on, 834–839
 of in-hospital cardiac arrest, initial PETCO₂ predicts, 2367–2371
 measures, short-term resuscitative, in adult swine model of ventricular fibrillation, effects of proximal and distal routes of intraosseous epinephrine administration on, randomized controlled study, 49–53
 and myocardial infarction, blood lactate concentration after cardiac arrest resulting from (letter), 1311–1313
 neurologic
 after cardiopulmonary resuscitation in rats, activation of autophagy improved, 1511–1518
 of comatose survivors after hanging, 1467–1472
 favorable, by early epinephrine administration within 19 minutes after EMS call for out-of-hospital cardiac arrest patients, 2284–2290
 in post-cardiac arrest patients, severe acidemia on arrival not predictive of, 425–428
 and survival, after CPR in ventricular fibrillation model of rats, edaravone improves, 1944–1949
 and survival, in cardiac arrest rats induced by ventricular fibrillation, benefits of respective and combined use of green tea polyphenols and ERK inhibitor on, 570–575
 of nonsurgical intervention in patients with perforated peptic ulcers, 1556–1560
 of NSTEMI patients transported by EMS vs private vehicle, 531–535
 for out-of-hospital cardiac arrest patients, role of prehospital advanced airway management on, 2101–2106
 of patients receiving CPR in ED of urban academic hospital, 1595–1599
 poor, and more severity, for patients with sepsis, higher serum level of myoglobin could predict, 948–952
 predictive scores, in ED, prediction of en-route complications during interfacility transport by, 877–882
 prospective evaluation among geriatric chest pain patients in ED observation unit, 207–211
 and renal impairment, in patients with Takotsubo cardiomyopathy, 548–552
 survival, in out-of-hospital cardiac arrest, interaction of diabetes mellitus and cardiac diseases on, 702–707
 of warfarinized patients with minor head injury and normal initial CT scan, 75–78
- Out-of-hospital
 cardiac arrest
 assessment of oxidative stress after, 1561–1566
 drowning-associated, relationship between drowning location and outcome after, 1799–1803
 incidence and clinical features of intracranial hemorrhage causing, 2326–2330
 patients, favorable neurological outcomes by early epinephrine administration within 19 minutes after emergency medical service call for, 2284–2290
 in patient with congenital long QT syndrome, extracorporeal CPR and therapeutic hypothermia for, a case report, 1321.e1–1321.e3
 prehospital therapeutic hypothermia after, 2209–2216
 public-access automated external defibrillators and defibrillation for (letter), 2041–2042
 role of prehospital advanced airway management on outcomes for patients with, 2101–2106
 sodium bicarbonate dilemma in patients with (letter), 1314–1315
- Outpatient stress testing, follow-up, barriers for low-risk chest pain patients presenting to ED chest pain unit, 790–793

- Ovarian hyperstimulation syndrome, concurrent, with perforated appendicitis following induction ovulation with HMG and HCG, a case report, 2051.e5–2051.e6c
- Ovarian vein thrombosis, idiopathic, rare cause of abdominal pain, a case report, 935.e1–935.e2
- Overdose
antipsychotic, prehospital activated charcoal use in (letter), 316–317
of dapson, acute, incidence and patterns of hemolytic anemia in, 366–369
intravenous amiodarone, acute, in swine model, Intralipid™ administration attenuates hypotensive effects of, 1389–1393
of medication, oral, activated charcoal may not be necessary in all (letter), 319–321
of methadone, causing acute cerebellitis and multi-organ damage, a case report, 343.e1–343.e3
salicylate, normal anion gap metabolic acidosis in (letter), 2457–2458
simulated massive acute, accuracy of trans-abdominal ultrasound in, 1455–1457
- Oxidant, and antioxidant levels, investigation in patients with acute stroke in emergency service, 2379–2383
- Oxidative stress
and inflammatory response, in multiple-organ dysfunction syndrome rats with severe burns, effects of insulin combined with ethyl pyruvate on, 2154–2158
after out-of-hospital cardiac arrest, assessment of, 1561–1566
- Oxygen
comparison with EtCO₂, achieved by polypropylene and conventional polyvinyl chloride facemask in healthy individuals, 1157–1159
supply, nasal cannula, high-flow, as treatment in hypercapnic respiratory failure, a case report, 1914.e1–1914.e2
- Oxygenation, during rapid sequence intubation, utility of noninvasive nasal positive pressure ventilators for optimizing, 1627–1630
- Pacemakers, transvenous, outcomes of emergency department placement of, 1411–1414
- Packed red blood cells, twenty-four-hour, requirement, is strongest independent prognostic marker of mortality in ED trauma patients, 1121–1124
- Paget-Schroetter syndrome, diagnostic limitations of imaging upper extremity deep vein thrombosis, a case report, 683.e1–683.e3
- Paget-Schroetter syndrome, secondary to thoracic outlet obstruction from muscle hypertrophy, a case report, 1323.e1–1323.e3
- Pain
abdominal, in older adults in ED, effect of surgical consult in treatment of, 1524–1527
acute, in ED, acupuncture vs intravenous morphine in management of, 2112–2116
associated with venous catheterization, topical ethyl chloride to reduce, randomized crossover trial, 845–850
conditions, ED triage pain protocol reduces time to receiving analgesics in patients with, 2362–2366
in ED, patients' perspectives on insufficiency of scales to rate their (letter), 2246–2247
flank
acute, into ED, contrast-enhanced ultrasound as imaging technique for patients with (letter), 324–325
and fever, with renal vein thrombosis, unusual initial presentation of systemic lupus erythematosus, a case report, 937.e1–937.e2
without contrast in patients with, a case report, 765.e3–765.e5
low back, in addition to IV dexamethasone, ketoprofen gel improves, 1458–1461
relief, in patients presenting to ED with acute renal colic, lidocaine as adjuvant to morphine improve, 443–448
right flank, deadly, IVC spontaneous rupture, a case report, 2050.e1–2050.e3
scores, technique choices, and and failure rates in local registry, advanced intravenous access, 553–557
severity, and CT imaging, effects on analgesia prescription in acute appendicitis, 36–39
shoulder, acute, late lead perforation presenting as, a case report, 2255.e1–2255.e3
- Pain management
acute, treatment of acute renal colic in US and French EDs, simulated cases and real cases in, 1955–1958
renal colic, desmopressin effectiveness in, 1535–1541
- Pain protocol, ED, triage, reduces time to receiving analgesics in patients with painful conditions, 2362–2366
- Palliative care, and hospice services in urban ED, dedicated palliative care nurse improves access to (letter), 2440–2441
- Pancreatitis
acute
with normal amylase and lipase, ED dilemma, a case report, 940.e5–940.e7
with normal lipase and amylase, a case report, 2254.e3–2254.e6
scrotal swelling and normal lipase, rare presentation of, a case report, 763.e5–763.e6
severe, and multiorgan dysfunction, H1N1 influenza presenting as, a case report, 1911.e1–1911.e2
hypercalcemia-induced, presenting with acute ST-elevations mimicking MI, a case report, 1187.e1–1187.e2
- Paradoxical embolism, ultrasound-accelerated rheolytic thrombectomy for impending (letter), 1711
- Paramedics
ability to perform endotracheal intubation during continuous chest compressions, randomized cadaver study comparing Pentax AWS and Macintosh laryngoscopes, 1835–1839
able to confirm endotracheal tube placement using ultrasonography (letter), 923–924
able to perform endotracheal intubation with access to patient through back seat of car, randomized crossover manikin study (letter), 1161–1163
assessment and treatment of upper airway obstruction in pediatric patients, exploratory analysis by Children's Safety Initiative-Emergency Medical Services, 599–601
in cervical immobilized manikin, Tracheal intubation with VivaSight-SL endotracheal tube by (letter), 309–310
comparison of McGrath-MAC and Macintosh laryngoscopes for child tracheal intubation during resuscitation by, 1338–1341
comparison of NIO and EZ-IO intraosseous access devices in adult patients under resuscitation performed by, randomized crossover manikin trial (letter), 1166–1167
experienced, in standardized airway manikin with restricted access, comparison of VivaSight single lumen endotracheal tube and the Macintosh laryngoscope for emergency intubation by, randomized, crossover trial (letter), 929–930
knowledge, attitude, and practices of, regarding optic nerve sheath diameter ultrasonography, 1160–1161
prepared to perform needle cricothyroidotomy (letter), 1310–1311
students' knowledge and attitudes regarding automated external defibrillators (letter), 1887–1888
wearing chemical, biological, radiological, and nuclear personal protective equipment, comparison of endotracheal intubation performed with 3 devices by (letter), 1883–1910
with and without chemical-biological-radiological-nuclear personal protective equipment, comparison of Bone Injection Gun and Jamshidi intraosseous access devices by (letter), 1307–1308
- Paraoxanase, as indicator of myocardial ischemia and its utility in determining extension of ischemia, 45–48
- Paraplegia, caused by aortic coarctation complicated with spinal epidural hemorrhage, a case report, 680.e1–680.e2
- Parapneumonic effusion, comparing posteroanterior with lateral and anteroposterior chest radiography in initial detection of, 2402–2407

- Paroxysmal cold hemoglobinuria, in child with RSV, a case report, 2252.e5–2252.e6
- Partial pressure of end-tidal carbon dioxide (PETCO₂), initial, predicts outcomes of in-hospital cardiac arrest, 2367–2371
- Pasteurella multocida*, infection, salvage of upper extremity in aggressive case of, a case report, 2470.e1–2470.e3
- Patella fracture, accuracy of point-of-care ultrasound as diagnostic tool for, 1576–1578
- Patient activation, and health literacy, assessing, in ED, 93–96
- Patient satisfaction
higher scores, ED opioid prescribing is not associated with (letter), 2032–2034
and provider, patient throughput, and revenue, effect of scribes on, 2018–2028
- PCC. *see* Prothrombin complex concentrate (PCC)
- PCI. *see* Percutaneous coronary intervention (PCI)
- Pediatrics
acute hemorrhagic edema of, and common mimics, a case report, 936.e3–936.e6
and adult academic EDs, impact of scribes on patient throughput in, 1982–1985
anxiety, effects of nursery songs for, reporting mistake of effect measure of association (letter), 663
anxiety before and during head CT, randomized single-blinded clinical trial on effects of nursery songs for (letter), 663
burner and stinger syndrome, magnetic resonance imaging for, 1442–1445
can teach their parents CPR and teaching does influence retention of their knowledge, 1159–1160
cardiopulmonary resuscitation safe and effective, current guideline recommendations for, 1658–1660
characterizing, presenting to three freestanding EDs (letter), 2045–2048
clotiapine poisoning, clinical manifestations and toxicokinetics, a case report, 2469.e1–2469.e2
critically ill, delayed sequence intubation with ketamine in, a case report, 1190.e1–1190.e2
in ED
with abdominal pain who are at low risk for acute appendicitis, prospective validation of biomarker panel to identify, 1373–1382
cost-effectiveness of point-of-care testing for dehydration in, 1573–1575
experience, REDCap (letter), 2048–2049
optimal dosing of intravenous ketamine for procedural sedation, 1347–1353
treatment of septic hip in, retrospective case series analysis, 602–605
emergent anticoagulation reversal, using 4-factor PCC, a case report, 1182.e1–1182.e2
endotracheal intubation, gold rules for (letter), 1711–1712
esophageal food impaction, point-of-care ultrasonography in detection of, a case report, 763.e1–763.e3
evaluation of proadrenomedullin as diagnostic or prognostic biomarker of acute appendicitis in, 2298–2305
forearm fracture, ultrasound-guided, reductions in resource-limited ED, 40–44
head trauma, accuracy of history of injury obtained from caregiver in, 1863–1867
intubation, laryngoscope use for (letter), 1729–1730
7-month-old baby, fulminant hepatic failure requiring liver transplantation due to accidental alcohol ingestion in, a case report, 2466.e1–2466.e2
new F(ab')₂ antivenom for treatment of crotaline envenomation in, 2003–2006
not moving her leg, a case report, 756.e1–756.e2
nurses' attitudes and beliefs concerning intraosseous access in (letter), 1890
occult methanol intoxication with hyperammonemia, a case report, 343.e5–343.e6
omental torsion mimicking perforated appendicitis in, emergency bedside sonography, a case report, 684.e–684.e4
paramedic assessment and treatment of upper airway obstruction in, exploratory analysis by Children's Safety Initiative–Emergency Medical Services, 599–601
with patient rhabdomyolysis as presenting feature of acute HIV-1 seroconversion in, a case report, 760.e3–760.e5
patients with abdominal pain discharged from ED, variation in advanced imaging for, 2320–2325
posttraumatic pulmonary hernia in, a case report, 759.e3–759.e4
procedural sedation
finding optimal initial dose of intravenous ketamine is still challenging for (letter), 1892
proactive approach to ED information technology, case of (letter), 1901–1902
psychiatric patients, deemed to be in need of inpatient admission, prior authorization for (letter), 915–916
with RSV, paroxysmal cold hemoglobinuria in, a case report, 2252.e5–2252.e6
scald burn, toxic shock syndrome due to methicillin-resistant *Staphylococcus aureus* infection after, a case report, 1735.e1–1735.e2
small bowel obstruction, point-of-care ultrasound in, ED case series, a case report, 2464.e1–2464.e2
sonographic aorta/IVC cross-sectional area index for evaluation of dehydration in, 1840–1844
survived from near-hanging, therapeutic hypothermia for, a case report, 1738.e5–1738.e7
with suspected appendicitis, usefulness of new and traditional serum biomarkers in, 871–876
tracheal intubation, during CPR, McGrath-MAC vs Macintosh laryngoscopes for (letter), 2453
tracheal intubation, during CPR, videolaryngoscopy or direct laryngoscopy for (letter), 2453
tracheal intubation, during resuscitation by paramedics, comparison of McGrath-MAC and Macintosh laryngoscopes for, 1338–1341
ventilation, with 3 different airway modalities, use of capnographs to assess quality of, 69–74
weighing 20 kg, application of LMA Classic for anesthetized, paralyzed (letter), 1697–1698
1-year-old patient, spontaneously rapid and complete resolution of acute subdural hemorrhage in, a case report, 2056.e1–2056.e4
Pentax AWS, and Macintosh laryngoscopes, ability of paramedics to perform endotracheal intubation during continuous chest compressions, randomized cadaver study comparing, 1835–1839
Pentax-AWS Airwayscope, muscle relaxant effect on tracheal intubation with, prospective randomized controlled trial, 674–675
Pentraxin 3, ischemia-modified albumin, and myeloperoxidase in predicting cardiac damage in acute carbon monoxide poisoning, a case report, 1927–1930
Peptic ulcer perforation, neutrophil-to-lymphocyte ratio, potential diagnostic marker for, 403–406
Percutaneous catheter intervention, vs thrombolysis, ST elevation in massive pulmonary embolism, a case report, 2466.e3–2466.e4
Percutaneous coronary intervention (PCI)
primary
fragmented QRS complexes have predictive value of imperfect ST-segment resolution in patients with STEMI after, 398–402
MHR as new prognostic marker in patients with STEMI undergoing, 240–244
with transcatheter arterial embolization, acute MI after, potency of, a case report, 1186.e1–1186.e3
Perforated peptic ulcers, outcome of nonsurgical intervention in patients with, 1556–1560
Perforation, of pyometra, spontaneous, presenting as acute abdomen, rare condition with considerable mortality, a case report, 761.e3–761.e5
Pericarditis
acute
handy echocardiographic marker in, echo probe sign (letter), 1883–1884
potentially missed, atypical pericarditis (letter), 2451–2453
role of combined ECG criteria in differential diagnosis of (letter), 1309
ST depression in lead aVL differentiates inferior STEMI from, 149–154
ST depression in lead aVL differentiates inferior STEMI from (letter), 1310

- Perihematomal edema
 area at initial ED presentation in patients with intracranial hematoma, prognostic value of, 1241–1246
 diameter as predictive indicator of 30-day mortality of intracranial hematoma (letter), 1701–1702
- Peripheral intravenous catheter, Delphi method validation of procedural performance checklist for insertion of ultrasound-guided (letter), 2227–2230
- Peripheral neuropathy, progressive, copper deficiency presenting as, a case report, 340.e5–340.e6
- Peripheral vein, cannulation, ultrasound-guided (letter), 1675
- Peripheral venous access, sites, identified by ultrasound, hand dominance in intravenous drug using patients does not affect, 1088–1091
- Peripheral venous catheter, size, and resident postgraduate years, influence on success of peripheral venous catheterization (letter), 1716–1718
- Peripheral venous catheterization, influence of peripheral venous catheter size and resident postgraduate years on success of (letter), 1716–1718
- Personal protective equipment, chemical-biological-radiological-nuclear, comparison of Bone Injection Gun and Jamshidi intraosseous access devices by paramedics with and without (letter), 1307–1308
- PETCO₂. *see* Partial pressure of end-tidal carbon dioxide (PETCO₂)
- Phalanx fractures, accuracy of bedside ultrasonography for diagnosis of (letter), 1698
- Pharmacist
 ED, monitoring of provider antibiotic selection aids appropriate treatment for outpatient UTI, 1600–1603
 emergency medicine, impact on antibiotic dosing adjustment, 980–984
 emergency medicine, impact on time to thrombolysis in acute ischemic stroke, 1997–1999
 impact on time to antibiotic administration in patients with sepsis in ED, 2117–2121
- Pharyngeal wire bristle foreign body, ED removal of, a case report, 1328.e5–1328.e6
- Phencyclidine,-positive urine drug screens, increase morbidity or mortality, trauma patients with, 1066–1068
- Phenylephrine
 push-dose, use, impact on subsequent preload expansion in ED setting, 2419–2422
 and terbutaline, effect on ischemic priapism, 222–224
- Phlegmasia cerulea dolens, May-Thurner syndrome as cause of, a case report, 1326.e3–1326.e4
- Phosphodiesterase-5, inhibition, improves macrocirculation and microcirculation during cardiopulmonary resuscitation, 162–166
- Physical examination, along with point-of-care echocardiography, indisputable right path (letter), 673
- Physicians
 are able to recognize ineffective (agonal) breathing as element of cardiac arrest (letter), 1165
 novice, comparison of AirTraq Avant® vs Macintosh laryngoscope for intubation during simulated resuscitation by (letter), 1708–1709
 novice, intubate with ETView tube without Macintosh laryngoscope (letter), 2242–2243
 and patient gestalt lead to shared decision to reduce unnecessary radiography in extremity trauma, 349–356
 and patient gestalt lead to shared decision to reduce unnecessary radiography in extremity trauma (erratum), 348
 young, prepared to perform focused assessment with sonography in trauma examination (letter), 314–315
- Placebo spray, vs vapocoolant spray, in adults undergoing venipuncture, prospective, randomized, double-blind controlled trial comparing, 798–804
- Placenta percreta, during pregnancy, spontaneous uterine rupture due to, a case report, 1918.e1–1918.e3
- Platypnea-orthodeoxia syndrome, simple clinical observation that heralds critical but treatable disease, a case report, 2053.e3–2053.e4
- Pleocytosis
 bacterial meningitis without (letter), 2443
 of CSF, *Streptococcus pneumoniae* meningitis without, a case report, 941.e1–941.e2
 in potentially immunocompromised host, CSF/blood glucose ratio and bacterial meningitis without (letter), 922
- Plesiomonas shigelloides*, meningitis in an adult in ED, a case report, 1329.e1–1329.e2
- Pneumatization, posttraumatic, of optic sheath, a case report, 1911.e3–1911.e4
- Pneumococcal meningitis, wider implications of normal cerebrospinal fluid cellularity in, 657
- Pneumomediastinum
 combined with hyperosmolar hyperglycemic state, Hamman syndrome, a case report, 2058.e1–2058.e2
 and retropharyngeal air, rare complication of influenza A and asthma in adult, a case report, 338.e1–338.e2
 still lung point, new sonographic evidence for, a case report, 344.e1–344.e2
- Pneumopericardium
 simple, due to blunt trauma progressing to tension pneumopericardium during transportation, a case report, 933.e3–933.e5
- Pneumoperitoneum
 catastrophic, in patient with perforated colorectal cancer with liver metastasis, a case report, 344.e3–344.e5
- Pneumorrhachis, spontaneous, a case report, 2465.e3–2465.e4
- Pneumothorax
 in patients with high positive end-expiratory pressure, absence of lung sliding is not reliable sign of (letter), 2034–2036
 primary spontaneous, pulmonary contusion post needle aspiration of, a case report, 1734.e1–1734.e3
 spontaneous
 primary, presenting hydrothorax predicts failure of needle aspiration in, 1075–1079
 recurrent, Mounier-Kuhn syndrome, unusual cause of, a case report, 122.e1–122.e2
 spontaneous bilateral, unusual complication of Behcet disease, a case report, 1320.e1–1320.e2
 tension, ultrasonography as a tool for prehospital recognition of (letter), 1302–1303
- PocketCPR, as educational tool according to smartphone attachment site, assessment of chest compression depth obtained using (letter), 2243–2246
- Point-of-care
 bedside, ultrasonography, effectiveness in diagnosis and management of metacarpal fractures, contribution for management of metacarpal fractures (letter), 674
 echocardiography
 physical examination along with, indisputable right path (letter), 673
 with speckle tracking, diagnosis of acute coronary occlusion in patients with NSTEMI, a case report, 1914.e3–1914.e6
 lactate, and PSP-S, assessment of Bayesian clinical decision model integrating, acute care diagnostics collaboration, 193–196
 sonography, integration during rapid sequence intubation in trauma resuscitation (letter), 330
 strain echocardiography in acute heart failure, (letter), 2234–2236
 testing, for dehydration in pediatric ED, cost-effectiveness of, 1573–1575
 transthoracic echocardiography, before CT angiogram of chest, saddle embolism diagnosed by, a case report, 2467.e1–2467.e2
 ultrasonography
 in detection of pediatric esophageal food impaction, a case report, 763.e1–763.e3
 at ED maximizes patient confidence in emergency physicians, 657–659
 impact on ED time to disposition for patients with nontraumatic shock, 1022–1030

- for management of shoulder dislocation in ED, 866–870
- ultrasound
 bedside, and computed tomography in elbow injuries, comparison of, 2186–2190
 as diagnostic tool for patella fractures, accuracy of, 1576–1578
 fellowship application process, innovation in (letter), 1303–1305
 in pediatric small bowel obstruction, ED case series, a case report, 2464.e1–2464.e2
 value for detecting nail bed injury in ED, 1850–1854
- Poison control, can be automate using webPOISONCONTROL, 1614–1619
- Poisoning
 with carbon monoxide
 ability of firefighters to recognize (letter), 1710
 acute, pentraxin 3, ischemia-modified albumin, and myeloperoxidase in predicting cardiac damage in, 1927–1930
 in acute and chronic periods, cardiac effects of (letter), 1303
 COHb levels predict long-term development of acute myocardial infarction in, 840–844
 deaths in United States, 1999 to 2012 (letter), 904
 myth busting in, 295–297
 patients with headache, treatment in, 2140–2145
 reversible left bundle-branch block due to, a case report, 342.e1–342.e3
 severe neurologic impairment and uncommon MRI findings after, a case report, 119.e1–119.e2
 with clonidine, pediatric, clinical manifestations and toxicokinetics, a case report, 2469.e1–2469.e2
 ethanol, and associated hypoventilation, with doxapram, treatment of, a case report, 2253.e1–2253.e2
 honey, heart stopping, not just Turkish honey, a case report, 1915.e1–1915.e3
 scambroid from tilapia, a case report, 339.e1–339.e2
 self
 patients, routine biological tests in, results from observational prospective multicenter study (letter), 2036–2037
 patients, routine biological tests in, results from observational prospective multicenter study (reply), 2037
- Polymorphonuclear neutrophil, and C5a2intra/C5a2 ratio level, for early sepsis in ED, prognostic significance of C5a2 on, 2084–2089
- Polypropylene, and conventional polyvinyl chloride facemask in healthy individuals, comparison of O₂ and EtCO₂ achieved by, 1157–1159
- Polyvinyl chloride, conventional, and polypropylene, facemask, in healthy individuals, comparison of O₂ and EtCO₂ achieved by, 1157–1159
- Positive end-expiratory pressure, high, absence of lung sliding is not reliable sign of pneumothorax in patients with (letter), 2034–2036
- Posterior reversible encephalopathy syndrome, 322–324
 after bevacizumab treatment presenting to ED as chest pain and headache, a case report, 1916.e1–1916.e2
- Posterior tibial nerve block, ultrasound-guided, ED, for calcaneal fracture analgesia, a case report, 1183.e1–1183.e3
- Postischemic conditioning, does not reduce muscle injury after tourniquet-induced ischemia-reperfusion injury in rats, 2065–2069
- Postpartum coma
 isolated hyperammonemia due to urea cycle disorder, rare cause of (letter), 1894–1895
 isolated hyperammonemia due to urea cycle disorder rare cause of, a case report, 1324.e3–1324.e4
- Postresuscitation, normal, 12-lead ECG, resuscitated out-of-hospital cardiac arrest with, a case report, 346.e1–346.e3
- Postsplinting x-rays, of nondisplaced hand, wrist, ankle, and foot fractures, 1625–1626
- Potassium
 blood, after avalanche-induced cardiac arrest (letter), 1317–1318
 measurement in ED
 interpret with caution (letter), 753
 interpret with caution (reply), 753–754
 as a potential predictive biomarker of brain hypoxia in avalanche victims (letter), 1315–1316
 serum, agreement of, measured by blood gas and biochemistry analyzer in patients with moderate to severe hyperkalemia, 794–797
 serum, cutoff for in-hospital triage after avalanche-induced cardiac arrest (letter), 1317
 serum concentration, predicts brain hypoxia on CT after avalanche-induced cardiac arrest, 856–860
- Pregnancy
 ectopic
 causing splenic rupture, a case report, 1184.e1–1184.e2
 partial resuscitative endovascular balloon occlusion of the aorta as a hemorrhagic shock adjunct for, a case report, 1917.e1–1917.e2
 intrauterine, in first-trimester ultrasonography, high-frequency linear transducer improves detection of, 288–291
 omental infarction, unusual cause of abdominal pain in, a case report, 1184.e3–1184.e4
 after out-of-hospital cardiac arrest, successful use of targeted temperature management in, a case report, 122.e3–122.e4
 patients in ED practice, treatment for maternal supraventricular tachyarrhythmia in (letter), 1702–1704
 seatbelt use during (letter), 326–328
 spontaneous uterine rupture due to placenta percreta during, a case report, 1918.e1–1918.e3
- Prehospital
 activated charcoal use in antipsychotic overdose (letter), 316–317
 advancing research, ethical and regulatory challenges in, focus on sepsis, 623–625
 drug-facilitated intubation protocols, statewide, variability in United States (letter), 2459–2460
 emergency medicine, use and implementation of SOPs and checklists in, 2432–2439
 endotracheal intubation and survival after out-of-hospital cardiac arrest, results from the Korean nationwide registry, 128–132
 and hospital costs of emergency care for frequent ED patients, 459–463
 screening tool utilizing end-tidal carbon dioxide predicts sepsis and severe sepsis, 813–819
 situations, GCS used as prognostic factor in unconscious patients following cardiac arrest (letter), 1178–1179
 vital signs, chart sharing system, development of, 88–92
- Prehospital delay, socioeconomic status, and acuity of illness, on presentation (letter), 1712–1714
- Prehospital recognition, of tension pneumothorax, ultrasonography as a tool for (letter), 1302–1303
- Prehospital Sepsis Project Score (PSP-S), and point-of-care lactate, assessment of Bayesian clinical decision model integrating, acute care diagnostics collaboration, 193–196
- Prehospital transportation, and clinical outcomes, in patients with non-STEMI, association between (letter), 1676–1677
- PRES. *see* Posterior reversible encephalopathy syndrome (PRES)
- Prescription
 analgesia, in acute appendicitis, effects of pain severity and CT imaging on, 36–39
 drug abuse, prescribing controlled medications to patients who die of, 30–35
 patterns, in opioid deaths shows startling findings, analysis of (letter), 748–749
- Prescription drug monitoring program, data, and prescription-related deaths, medical examiner, description of, 510–514
- Priapism, ischemic, effect of phenylephrine and terbutaline on, 222–224
- Primary care linkage, for patients discharged from ED, transitional care clinics for follow-up and, 1230–1235
- Primary care provider, enrollment with, does not preclude ED visits for patients with woman's health-related problems, 266–268

- Proadrenomedullin, evaluation as diagnostic or prognostic biomarker of acute appendicitis in children, 2298–2305
- Procalcitonin
 -guided therapy for initiation of antibiotics in ED, 1286–1293
 levels, predict identification of bacterial strains in blood cultures of septic patients (letter, 2453–2454)
 levels predict to identify bacterial strains in blood cultures of septic patients, 2150–2153
- Prophylaxis, venous thromboembolism, in ED observation units, (letter), 2238–2240
- Propofol
 administration for procedural sedation in ED (letter), 1675–1676
 -induced hypotension, ultrasound measurement of IVC collapse predicts, 1125–1128
 vs ketamine, -propofol combination (ketofol), for procedural sedation and analgesia, 558–569
- Prospective validation of biomarker panel to identify pediatric ED patients with abdominal pain who are at low risk for acute appendicitis, 1373–1382
- Prothrombin complex concentrate (PCC)
 administration through intraosseous access for reversal of rivaroxaban, a case report, 685.e1–685.e2
 4-factor
 associated hypotension, a case report, 2059.e1–2059.e2
 emergent pediatric anticoagulation reversal using, a case report, 1182.e1–1182.e2
 high-dose, for acute warfarin reversal, thrombosis induced by, a case report, 1736.e1–1736.e3
 issues with treating superwarfarin ingestion with (letter), 1168
 superwarfarin ingestion treated successfully with, a case report, 116.e1–116.e2
- Prothrombin time (PT), INR, highly elevated, after dabigatran etexilate use in patient with acute kidney injury, successful hemostasis and reversal of, a case report, 758.e5–758.e6
- Providence Pedi cast-sling, technique and application, 651–653
- Pseudoaneurysm, radial, after puncture for blood gas analysis, a case report, 757.e1–757.e2
- Psychiatric patients
 child and adolescent, deemed to be in need of inpatient admission, prior authorization for (letter), 915–916
 in ED, prolonged length of stay in, multivariable predictive model, 133–139
- Psychiatrists, recension of emergency physicians' involuntary admissions, role of alcohol intoxication in (letter), 2226–2227
- PT. *see* Prothrombin time (PT)
- Public health, more than 15 minutes of resuscitation prior to termination of resuscitation results in undue harm to (letter), 1689–1690
- Pulmonary artery, unilateral absence of, concomitant with chronic pulmonary thromboembolism in adult patient, a case report, 1916.e3–1916.e5
- Pulmonary contusion, post needle aspiration of primary spontaneous pneumothorax, a case report, 1734.e1–1734.e3
- Pulmonary decongestion, in acute heart failure, resolution of sonographic B-lines as measure of, 1129–1132
- Pulmonary edema
 postobstructive, that developed immediately after emergency surgical cricothyroidotomy, a case report, 936.e1–936.e2
 severe, centrally acting antihypertensive, clonidine, combined to venous dilator, nitroglycerin, to handle, a case report, 676.e5–676.e7
- Pulmonary embolism
 acute
 high-sensitivity troponin and right ventricular function in, 1579–1582
 in pulmonary trunk and main pulmonary arteries, ECG characteristics of, 212–217
 right-sided electrocardiogram usage in, 1437–1441
 simultaneous recording of right- and left-sided electrocardiogram in, a case report, 1183.e5–1183.e7
 syncope and collapse in, 1251–1257
 catheter-directed option for treatment of (letter), 1691–1693
 comparison between systemic and catheter thrombolysis in patients with, 985–988
 contrast-induced nephropathy related to adverse events in (letter), 2042–2044
 endobronchial ultrasound, unusual diagnostic tool for, a case report, 684.e1–684.e2
 epipericardial fat necrosis clinically mimicking, CT angiography findings, a case report, 2056.e5–2056.e6
 intermediate- to high-risk, with B-type BNP, a case report, 2463.e1–2463.e3
 with low dose of tissue plasminogen activator after thoracic surgery, successful treatment of, a case report, 2259.e5–2259.e6
 massive
 during distal femur fracture surgery, salvage thrombolysis and ECMO for, a case report, 1189.e3–1189.e5
 mimicking ECG pattern of Brugada syndrome, a case report, 933.e1–933.e2
 ST elevation in, thrombolysis vs percutaneous catheter intervention, a case report, 2466.e3–2466.e4
 in patients diagnosed with contrast CT, relation of contrast nephropathy to adverse events in, 1247–1250
 in patients with Alzheimer disease, in-hospital mortality due to, sex effect (letter), 665–666
 predicting prognosis in (letter), 1170
 risk after prior negative CT pulmonary angiogram, 1968–1972
 saddle, diagnosed by point-of-care transthoracic echocardiography before CT angiogram of chest, a case report, 2467.e1–2467.e2
 spectrum of disturbances in cerebral function in elderly patients with, 651
 thrombolysis during continuous chest compression in patient with cardiac arrest due to, prolonged CPR-induced spinal cord injury, a case report, 681.e3–681.e5
 thrombolysis in (letter), 1720–1721
 treated with thrombolysis, pulseless electrical activity in, 1963–1967
- Pulmonary thromboembolism
 acute, prognostic importance of thiol/disulfide homeostasis in patients with, 2315–2319
 chronic, concomitant with unilateral absence of pulmonary artery, in adult patient, a case report, 1916.e3–1916.e5
 after major surgery, atrial septal defect can mimic, a case report, 121.e3–121.e6
- Pulseless electrical activity, in pulmonary embolism treated with thrombolysis, 1963–1967
- Purpuric skin eruption, in an illicit drug user, a case report, 1321.e5–1321.e6
- Pyelonephritis
 acute
 diagnostic performance of initial serum lactate for predicting bacteremia in female patients with, 1359–1363
 health care-associated, associated with inappropriate empiric antibiotic therapy in ED, 1415–1420
- Pyometra, spontaneous perforation of, presenting as acute abdomen, rare condition with considerable mortality, a case report, 761.e3–761.e5
- Pyruvate
 alleviates lipid peroxidation and multiple-organ dysfunction in rats with hemorrhagic shock, 525–530
 alleviates lipid peroxidation and multiple-organ dysfunction in rats with hemorrhagic shock (erratum), 1330
- QRS analysis, high-frequency, superior to conventional ST-segment analysis of women with chest pain, 437–442
- QRS complexes, fragmented, have predictive value of imperfect ST-segment resolution in patients with STEMI after primary PCI, 398–402
- qSOFA. *see* quick Sepsis-related Organ Failure Assessment (qSOFA)
- quick Sepsis-related Organ Failure Assessment (qSOFA), predictive performance for mortality and ICU admission in patients with infection at ED, 1788–1793

- Racial disparity, and illness severity, in ED wait times, relationship between, 10–15
- Radial artery catheterization, in adults, optimal wrist positioning for, 2372–2378
- Radiation dose, cumulative effective, received by blunt trauma patients arriving to military level I trauma center from point of injury and interhospital transfers, 2397–2401
- Radiation therapy, emergent, as definite airway management for dyspnea with mediastinal mass, a case report, 2258.e5–2258.e6
- Radiography
injuries, after blunt elbow trauma, performance of 4-way ROM test for, 235–239
unnecessary, in extremity trauma, physician and patient gestalt lead to shared decision to reduce, 349–356
unnecessary, in extremity trauma, physician and patient gestalt lead to shared decision to reduce (erratum), 348
in wrist trauma, Karadeniz wrist rules for the use of, 2074–2078
- Radiologists, on-call, comparison with emergency physicians, CT interpretations in multiply injured patients, 2331–2335
- Radiotherapy, carotid blowout and cerebral gas embolism related to bidirectional carotid-esophageal fistula, serious complication of esophageal cancer under, a case report, 683.e5–683.e6
- Range of motion (ROM), test, 4-way, for radiographic injuries after blunt elbow trauma, performance of, 235–239
- Rapid ventricular response, or tachycardia, three faces of atrial fibrillation (letter), 931–932
- Rash
and fever, acute generalized exanthematous pustulosis, uncommon cause of, a case report, 681.e1–681.e2
seventeen-year-old sexually active male with, a case report, 764.e1–764.e3
Varicella-zoster virus encephalitis in immunocompetent patient without, a case report, 2257.e1–2257.e2
- Recombinant human soluble thrombomodulin, and short-term mortality of infection patients with DIC, 1876–1882
- Recombinant transplasminogen activator (rt-PA), intravenous, for acute ischemic stroke, fatal hemopericardium after, a case report, 2462.e5–2462.e6
- REDCap. *see* Research Electronic Data Capture (REDCap)
- Renal artery aneurysm, left, rupture, rare abdominal emergency, a case report, 347.e3–347.e4
- Renal colic
acute
lidocaine as adjuvant to morphine improve pain relief in patients presenting to ED with, 443–448
treatment in US and French EDs, simulated cases and real cases in acute pain management, 1955–1958
pain management, desmopressin effectiveness in, 1535–1541
uncomplicated, suspected, validity of STONE scores in younger patients presenting with, 230–234
- Renal failure, acute, and rhabdomyolysis, synthetic cannabinoid hyperemesis resulting in, a case report, 765.e1–765.e2
- Renal impairment, and outcome in patients with Takotsubo cardiomyopathy, 548–552
- Renal replacement therapy, continuous, new treatment of seriously hyperglycemia, a case report, 2469.e3–2469.e4
- Renal vein thrombosis, with fever and flank pain, unusual initial presentation of systemic lupus erythematosus, a case report, 937.e1–937.e2
- Reperfusion, should be revisited (editorial), 1086–1087
- Repolarization, early, terminal QRS distortion present in anterior myocardial infarction but absent in, 2182–2185
- Repositioning maneuvers, liberatory vertigo new prognostic factor for, 1548–1551
- Rescue equipment, assessing efficacy in lifeguard resuscitation efforts for drowning, 480–485
- Rescuer
factors predict high-quality CPR—manikin-based study of health care providers, 20–24
novice, body weight, relationship with chest compression depth, during CPR, 2411–2413
single, ventilation, using bag-valve mask with internal handle, randomized crossover trial, 1991–1996
- Research Electronic Data Capture (REDCap), pediatric ED experience (letter), 2048–2049
- Residents, postgraduate years and peripheral venous catheter size influence on success of peripheral venous catheterization (letter), 1716–1718
- Respiratory depression, in intoxicated trauma patient, opioids to blame, 250–253
- Respiratory failure
acute, persistent hiccups followed by, a case report, 1181.e3–1181.e4
hypercapnic, high-flow nasal cannula oxygen supply as treatment in, a case report, 1914.e1–1914.e2
- Respiratory function, effects of spinal immobilization at 20° on, 1959–1962
- Respiratory syncytial virus (RSV), paroxysmal cold hemoglobinuria in child with, a case report, 2252.e5–2252.e6
- Resuscitation
of cocaine-induced cardiovascular arrest in rat model, intravenous lipid emulsion in, 1452–1454
comparison of Macintosh and Intubrite laryngoscopes for orotracheal intubation by nurses during (letter), 1724–1725
endovascular balloon occlusion of aorta, usefulness in detecting source of hemorrhage due to abdominal blunt trauma, a case report, 2057.e1–2057.e3
fluid, in anaphylactic patient, firefighters able to perform intraosseous access and start (letter), 1707–1708
impact of body mass index on patient survival after therapeutic hypothermia after, 722–725
lifeguard, efforts for drowning, assessing efficacy of rescue equipment in, 480–485
15 minutes duration, appropriate before termination of traumatic cardiac arrest, 505–509
more than 15 minutes, prior to termination of resuscitation results in undue harm to public health (letter), 1689–1690
ongoing, is associated with increased return of spontaneous circulation, sodium bicarbonate administration during, 225–229
outcome measures, short-term, in adult swine model of ventricular fibrillation, effects of proximal and distal routes of intraosseous epinephrine administration on, randomized controlled study, 49–53
out-of-hospital cardiac arrest with normal postresuscitation 12-lead ECG, a case report, 346.e1–346.e3
by paramedics, comparison of McGrath-MAC and Macintosh laryngoscopes for child tracheal intubation during, 1338–1341
partial, endovascular balloon occlusion of aorta, as hemorrhagic shock adjunct for ectopic pregnancy, a case report, 1917.e1–1917.e2
performed by paramedics, comparison of NIO and EZ-IO intraosseous access devices in adult patients under, randomized crossover manikin trial (letter), 1166–1167
simulated
comparison of exchange of laryngeal mask airway and Igel for tracheal tube using ETTI during (letter), 106–107
effect of bed width on quality of compressions in, randomized crossover manikin study, 2272–2276
simulated, by novice physicians, comparison of AirTraq Avant® vs Macintosh laryngoscope for intubation during (letter), 1708–1709
successful, in dose-dependent fashion in pig model of cardiac arrest, 2,3-butanedione monoxime facilitates, 1053–1058
tracheostomy patient in ED, 1148–1155
trauma, integration of point-of-care sonography during rapid sequence intubation in (letter), 330
- Return of spontaneous circulation, increased, sodium bicarbonate administration during ongoing resuscitation is associated with, 225–229
- Rhabdomyolysis, acute, in healthy woman, a case report, 113.e1–113.e2

Rhabdomyolysis

- and acute renal failure, synthetic cannabinoid hyperemesis resulting in, a case report, 765.e1–765.e2
- mefloquine-associated, a case report, 2250.e5–2250.e6
- nontraumatic, coma blister in, a case report, 1324.e1–1324.e2
- as presenting feature of acute HIV-1 seroconversion in pediatric patient, a case report, 760.e3–760.e5
- and profound acidosis, due to dietary supplement use, survival after, a case report, 2259.e1–2259.e3
- severe
 - and hyperthermia, from synthetic cannabinoids, a case report, 121.e1–121.e2
 - and hyperthermia, from synthetic cannabinoids (letter), 1690–1691
 - with minor trauma, sickle cell trait to blame, a case report, 341.e3–341.e4

Right ventricular function, and high-sensitivity troponin, in acute pulmonary embolism, 1579–1582

Risk factors

- comparison with ED visit outcomes, nonsuicidal self-injury and suicide attempts among ED patients older than 50 years, 1016–1021
- or embolic source, acute bilateral lower extremity arterial occlusion without, a case report, 762.e3–762.e4
- or fever, meningococcal meningitis caused by septic arthritis of a lumbar facet joint in patient without, a case report, 762.e1–762.e2
- and prevalence, for central diabetes insipidus in cardiac arrest survivor treated with targeted temperature management, 1400–1405
- of transfusion-related acute lung injury after blood products transfusions (letter), 1674–1675

Risk stratification, and management of patients with febrile neutropenia, provider familiarity with specialty society guidelines for (letter), 1704–1705

Rivaroxaban

- hypersensitivity reaction after administration of, a case report, 1325.e1–1325.e2
- left atrial thrombus formation after brief interruption of, a case report, 116.e3–116.e4
- PCC administration through intraosseous access for reversal of, a case report, 685.e1–685.e2
- superior mesenteric artery thrombosis after abrupt discontinuation of, a case report, 764.e5–764.e7

Rocuronium, succinylcholine, and hyperkalemia (letter), 1705–1706

Routine biological tests, in self-poisoning patients, 1383–1388

RSV. *see* Respiratory syncytial virus

Rupture

- bilateral spontaneous atraumatic, of Achilles tendon in athlete, a case report, 114.e1–114.e2
- cardiac, oozing-type, after acute MI, use of IABP support for, a case report, 120.e1–120.e3
- IVC, spontaneous, deadly right flank pain, a case report, 2050.e1–2050.e3
- of left gastroepiploic vessel, spontaneous, after forceful retching, laparoscopic management for, a case report, 759.e1–759.e2
- of left renal artery aneurysm, rare abdominal emergency, a case report, 347.e4–347.e4
- left ventricular free-wall, early, non-STEMI never to be neglected, a case report, 113.e3–113.e4
- of noncoronary sinus of Valsalva into right atrium, unusual cause of acute chest pain, a case report, 2052.e1–2052.e3
- splenic
 - ectopic pregnancy causing, a case report, 1184.e1–1184.e2
 - nontraumatic, as initial manifestation of acute myeloid leukemia in ED, a case report, 1731.e1–1731.e3
- spontaneous, of right gastroepiploic artery aneurysm, fibromuscular dysplasia presenting as hemorrhagic shock due to, a case report, 677.e3–677.e5

Salicylate

- overdose, normal anion gap metabolic acidosis in (letter), 2457–2458
- toxicity, in absence of anion gap metabolic acidosis, a case report, 1328.e1–1328.e3

Salvage, of upper extremity in aggressive case of *Pasteurella multocida* infection, a case report, 2470.e1–2470.e3

Scald burn, pediatric, toxic shock syndrome due to methicillin-resistant *Staphylococcus aureus* infection after, a case report, 1735.e1–1735.e2

Scaphoid fractures, orthopedic literature 2015, 1663–1667

Scorpion venom, atrial fibrillation, rare complication of, a case report, 937.e3–937.e4

Screening, brief intervention, and referral to treatment (SBIRT), 5-year comparison of staff buy-in (letter), 2232–2234

Scribes

- effect on patient throughput, revenue, and patient and provider satisfaction, 2018–2028

Impact on patient throughput in adult and pediatric academic EDs, 1982–1985

Scrotal pain, acute, clinical findings in young adults with, 1931–1933

SCUBE1. *see* Signal peptide-CUB-EGF domain-containing protein 1 (SCUBE1)

Seatbelt, use during pregnancy (letter), 326–328

Sedation

- of agitated patients in ED, chemical agents for, 2426–2431
- procedural

- and analgesia, ketamine-propofol combination (ketofol) vs propofol for, 558–569

- in children in ED, optimal dosing of intravenous ketamine for, 1347–1353

- in ED, propofol administration for (letter), 1675–1676

- pediatric, finding optimal initial dose of intravenous ketamine is still challenging for (letter), 1892

- pediatric, proactive approach to ED information technology, case of (letter), 1901–1902

Seizure, is not real seizure, a case report, 757.e3–757.e5

Self-injury, nonsuicidal, and suicide attempts among ED patients older than 50 years, comparison of risk factors and ED visit outcomes, 1016–1021

Self-poisoning, routine biological tests in patients with, 1383–1388

Sepsis

- biomarkers in (letter), 924–925

- burn, through inhibiting NLRP3 inflammasome activation, artemisinin protects mice against, 772–777

- and clinical outcome, C1 inhibitor in (letter), 1179–1180

- clinical value of soluble uPAR in diagnosis, prognosis, and therapeutic guidance of, 375–380

- confirmed, presenting hemodynamic phenotypes in ED patients with, 2291–2297

- early, in ED, prognostic significance of C5a2 on polymorphonuclear neutrophil and C5a2intra/C5a2 ratio level for, 2084–2089

- in ED, pharmacist impact on time to antibiotic administration in patients with, 2117–2121

- emergency medical services annotated literature in review 2016, 2193–2199

- ethical and regulatory challenges in advancing prehospital research, focus on, 623–625

- forgotten predictors in (letter), 1714

- higher serum level of myoglobin could predict more severity and poor outcome for patients with, 948–952

- high immunoglobulin E values at admission predict mortality in ED patients with, 1589–1594

- hip, treatment in pediatric ED, retrospective case series analysis, 602–605

- after ICU discharge, positive fluid balance as major predictor of clinical outcome of patients with, 2122–2126

- infection, and systemic inflammatory response syndrome, in multiple-injured patients, fresh frozen plasma is permissive for, 1480–1485

- patients with hepatic dysfunction, lactate clearance and mortality in, 1011–1015
- predictive role of lactate dehydrogenase in patients with, 2167–2171
- predictors of mortality in (letter), 2223–2224
- procalcitonin levels predict identification of bacterial strains in blood cultures of (letter), 2453–2454
- procalcitonin levels predict to identify bacterial strains in blood cultures of patients with, 2150–2153
- prognostic role of non-critical lactate levels for in-hospital survival time among ED patients with, 170–173
- serial change of C1 inhibitor in patients with, 594–598
- serial change of C1 inhibitor in patients with (erratum), 1741–1742
- severe
- among ED patients, effect of electronic medical record alert for, 185–188
 - elevated cardiac biomarkers are not associated with mortality in low-risk cardiac patients with (letter), 1899–1900
 - predictors of mortality in (letter), 1899
- and severe sepsis, prehospital screening tool utilizing end-tidal carbon dioxide predicts, 813–819
- triage, alert, and sepsis protocol lower times to fluids and antibiotics in ED, 1–9
- Septic arthritis**
- of lumbar facet joint, in patient without fever or risk factors, meningococcal meningitis caused by, a case report, 762.e1–762.e2
 - spontaneous, of pubic symphysis in afebrile patient, a case report, 934.e5–934.e6
 - and transient synovitis, in ED, guideline for differential diagnosis between, 1631–1636
- Septic embolus, and endocarditis, Hampton's hump in patient with, a case report, 942.e1–942.e3**
- Septic shock**
- due to urinary infection, urinary obstruction is important complicating factor in patients with, 694–696
 - after ICU discharge, positive fluid balance as major predictor of clinical outcome of patients with, 2122–2126
 - ulinastatin ameliorates gastrointestinal injury sustained in 2-hit porcine model of, 1497–1504
- Serotonin syndrome, associated with therapeutic metaxalone in patient with cirrhosis, a case report, 346.e5–346.e6**
- Sex, effect, in-hospital mortality due to pulmonary embolism in patients with Alzheimer disease (letter), 665–666**
- Sex differences**
- in frequent ED use among those with multimorbid chronic diseases, 2127–2131
 - in STEMI activation for patients presenting to ED 1939, 1939–1943
- Sexual assault, and domestic violence, prevalence of strangulation in survivors of, 1281–1285**
- Sexually transmitted diseases, compliance with centers for disease control guidelines for ED patients with (letter), 1727–1729**
- Sexually transmitted infection (STI), from 2001 to 2010, retrospective cross-sectional study of patients treated in US EDs and ambulatory care clinics with, 1808–1811**
- SHARE-FI. see Survey of Health, Ageing and Retirement in Europe Frailty Instrument (SHARE-FI)**
- Shen-Fu injection, effects on coagulation-fibrinolysis disorders in porcine model of cardiac arrest, 469–476**
- Shock**
- electric, implantable cardiac defibrillator, ED evaluation and management of, 1140–1147
 - nontraumatic, impact of point-of-care ultrasonography on ED time to disposition for patients with, 1022–1030
- Shock index**
- modified and age shock index, for prediction of mortality in Emergency Severity Index level 3, 2079–2083
 - triage time, modified, and age, association with mortality in emergency severity index level 2 patients, 63–68
- Short axis out-of-plane, vs long axis in-plane techniques, effects during ultrasound-guided vascular access, 778–783**
- Shortness of breath**
- acute, *Brucella*-induced sublingual abscess causes, a case report, 1737.e3–1737.e4
 - BRIPPED scan for evaluation of ED patients with, 386–391
 - and unexpected imaging findings, a case report, 937.e3–937.e4
- Shoulder dislocation**
- anterior
 - bilateral, sequential, a case report, 939.e1–939.e2
 - effectiveness of newly developed reduction method of, 1406–1410
 - GONAIIS method, new autoreduction method for, a case report, 120.e5–120.e7
 - presenting as hypovolemic shock, axillary artery laceration after, a case report, 1739.e1–1739.e4 - atraumatic bilateral posterior, rare case of sleep deprivation, a case report, 1322.e3–1322.e4
 - in ED, point-of-care ultrasonography for management of, 866–870
 - posterior, traumatic, with large engaging Hill-Sachs lesion, splinting technique, a case report, 682.e1–682.e3
 - reduction
 - Cunningham technique for (letter), 1890–1890
 - Sool's method of, Cunningham technique redux (letter), 1891
- Shoulder pain, acute, late lead perforation presenting as, a case report, 2255.e1–2255.e3**
- Sick leave, paid, is associated with fewer ED visits among US private sector working adults, 784–789**
- Sickle cell trait, to blame, severe rhabdomyolysis with minor trauma, a case report, 341.e3–341.e4**
- Signal peptide-CUB-EGF domain-containing protein 1 (SCUBE1), and coronary artery disease, relation between (letter), 647–648**
- SimMon, simulation application, efficacy for emergency response training (letter), 927–928**
- Skin infections, bacterial, acute, failure of outpatient antibiotics among patients hospitalized for, 957–962**
- Sleep deprivation, atraumatic bilateral posterior shoulder dislocations rare case of, a case report, 1322.e3–1322.e4**
- Sling, and swathe modification for immobilization of posterior glenohumeral dislocations, 668–670**
- Small bowel obstruction**
- is associated with poor prognosis, large-bowel disease presenting as, 477–479
 - pediatric, point-of-care ultrasound in, ED case series, a case report, 2464.e1–2464.e2
- Smartphone**
- attachment site, assessment of chest compression depth obtained using PocketCPR as educational tool according to (letter), 2243–2246
 - reading, diagnostic performance in coronary CT angiography in patients with acute chest pain at ED, 1794–1798
- Sniffing position**
- elevated, for endotracheal intubation, need to use, 928
 - for good glottic view (letter), 1156
- Socioeconomic status, prehospital delay, and acuity of illness, on presentation (letter), 1712–1714**
- Sodium, serum levels, measured by blood gas analyzer and biochemistry autoanalyzer in patients with hyponatremia, eunatremia, and hypernatremia, comparison of, 1473–1479**
- Sodium bicarbonate**
- administration during ongoing resuscitation is associated with increased return of spontaneous circulation, 225–229
 - dilemma in patients with out-of-hospital cardiac arrest (letter), 1314–1315
- Sodium polystyrene sulfate, moving away from (letter), 1716**
- Soft tissue, foreign bodies, ultrasound for evaluation of, before and after addition of fluid to surrounding interstitial space in cadaveric model, 1779–1782**

- Soft tissue injury, but also for improved laryngoscopy and tracheal intubation, lip pulling method for prevention of (letter), 1694
- Solid malignancy, application of high-flow nasal cannula in ED for patients with (letter), 2222–2223
- Sonography
bedside, emergency, omental torsion, mimicking perforated appendicitis in pediatric patient, a case report, 684.e3–684.e4
point-of-care, integration during rapid sequence intubation in trauma resuscitation (letter), 330
- Sool's method, of dislocated shoulder reduction, Cunningham technique redux (letter), 1891
- Sotalol, upper extremity edema caused by, a case report, 1920.e5–1920.e6
- South African flag sign, teaching tool for easier ECG recognition of high lateral infarct (letter), 107–109
- Specialist, availability in emergencies, contributions of response times and use of ad hoc coverage in New York State, 687–693
- Spinal cord compression, thoracic, masquerading as cauda equina syndrome, a case report, 756.e3–756.e5
- Spinal cord injury
CPR-induced, prolonged, thrombolysis during continuous chest compression inpatient with cardiac arrest due to pulmonary embolism, a case report, 681.e3–681.e5
secondary prevention, in-water, out-of-water cervical immobilization saves time (letter), 1172–1174
- Spinal epidural hemorrhage, paraplegia caused by aortic coarctation complicated with, a case report, 680.e1–680.e2
- Spinal immobilization, effects at 20° on respiratory functions, 1959–1962
- Spine board, long, does not reduce lateral motion during transport, randomized healthy volunteer crossover trial, 717–721
- Spine fixation, post-cervical, patients, percutaneous tracheostomy in, challenging the dogma (letter), 910–911
- Splenectomy, hemoperitoneum after blunt abdominal trauma 27 years after, a case report, 1325.e3–1325.e4
- Splenic rupture
ectopic pregnancy causing, a case report, 1184.e1–1184.e2
nontraumatic, as initial manifestation of acute myeloid leukemia in ED, a case report, 1731.e1–1731.e3
- Splint, modified coaptation, with sugar tong intrinsic plus extension for initial management of wrist drop, 659–663
- Sports medicine, orthopedic literature 2015, 1663–1667
- Standard operating procedures (SOPs), and checklists, use and implementation in prehospital emergency medicine, 2432–2439
- Status epilepticus, and type 2 myocardial infarction, a case report, 1735.e3–1735.e4
- ST depression
in lead aVL differentiates inferior STEMI from pericarditis, 149–154
in lead aVL differentiates inferior STEMI from pericarditis (letter), 1310
- ST elevation
in massive pulmonary embolism, thrombolysis vs percutaneous catheter intervention, a case report, 2466.e3–2466.e4
with spontaneous coronary artery dissection, in aVR and posterior lead, a case report, 2470.e5–2470.e7
- STEMI. *see* ST-segment elevation myocardial infarction (STEMI)
- Sternotomy, post, mediastinitis, diagnosing, in ED, 618–622
- Sternum, anatomic variations of, may mimic traumatic complications, a case report, 1912.e1–1912.e2
- Stevens-Johnson syndrome
Fuchs syndrome, rare and unique variant of (letter), 910
after treatment with azithromycin, uncommon culprit, a case report, 676.e1–676.e3
- STI. *see* Sexually transmitted infection (STI)
- STONE score
and derivation of the modified STONE score, external validation of, 1567–1572
validity in younger patients presenting with suspected uncomplicated renal colic, 230–234
- Straight back syndrome, manifesting as acute MI, a case report, 1913.e1–1913.e3
- Strangulation, prevalence in survivors of sexual assault and domestic violence, 1281–1285
- Streptococcus gallolyticus (bovis)*, rare presentation of meningitis in ED, a case report, 677.e1
- Streptococcus pneumoniae*
infection, prognostic factors in adults, 202–206
meningitis without pleocytosis of CSF, a case report, 941.e1–941.e2
rare pathogen in epidural abscess, a case report, 1919.e3–1919.e4
- Stretcher, reducible and standard, on mechanical compression versus manual compression, in small elevator, quality between, 1604–1609
- Stroke
acute
CT angiography in (revisiting the 4Ps of imaging), 282–287
in emergency service, investigation of oxidant and antioxidant levels in patients with, 2379–2383
a case report, 1189.e1–1189.e2
emergency medical services annotated literature in review 2016, 2193–2199
prevention for patients with atrial fibrillation (letter), 1685–1687
prevention in real-life practice (letter), 1714–1715
suspects in rural state, emergency transport of, 1640–1644
- ST-segment
new algorithms, to determine culprit artery location in acute inferior MI, 1772–1778
resolution, imperfect, in patients with STEMI after primary PCI, fragmented QRS complexes have predictive value of, 398–402
- ST-segment analysis, conventional, of women with chest pain, high-frequency QRS analysis superior to, 437–442
- ST-segment elevation
acute, mimicking MI, hypercalcemia-induced pancreatitis presenting with, a case report, 1187.e1–1187.e2
high-risk electrocardiographic presentations lacking anatomically oriented, electrocardiogram in ACS patient, 611–617
- ST-segment elevation myocardial infarction (STEMI)
activation, for patients presenting to ED 1939, sex differences in, 1939–1943
acute, relation between monocyte to HDL ratio and no-reflow phenomenon in patients with, 1542–1547
emergency coronary angiography in out-of-hospital cardiac arrest patients without, a case report, 118.e1–118.e3
inferior, from pericarditis, ST depression in lead aVL differentiates, 149–154
inferior, from pericarditis, ST depression in lead aVL differentiates (letter), 1310
non, early left ventricular free-wall rupture in, never to be neglected, a case report, 113.e3–113.e4
notification by EMS predicts shorter door-to-balloon time and smaller infarct size, 1610–1613
patients, undergoing interhospital transfer, first-door-to-balloon time delay in, 767–771
in patient with ITP, successful thrombolytic therapy for, a case report, 345.e1–345.e3
after primary PCI, fragmented QRS complexes have predictive value of imperfect ST-segment resolution in patients with, 398–402
relationship between fibrinogen to albumin ratio and severity of coronary artery disease in patients with, 1037–1042
undergoing primary PCI, MHR as new prognostic marker in patients with, 240–244
unusual adverse event of acute urinary retention, a case report, 347.e1–347.e2
- Stuttering, after minor head trauma, a case report, 685.e3–685.e4
- Stylet, semirigid, use of, increase efficacy of endotracheal intubation when using EView tube (letter), 1908–1909
- Subarachnoid hemorrhage

- aneurysmal, admission serum lactate predicts mortality in, 708–712
 –associated out-of-hospital cardiac arrest, bispectral index monitoring in, a case report, 934.e1–934.e3
 retroclival, posttraumatic, a case report, 2257.e3–2257.e5
 role of levosimendan in management of, 298–306
 traumatic, complicated by cerebral vasospasm, hemodynamic challenges in (letter), 904–906
- Subdural hemorrhage
 acute, spontaneously rapid and complete resolution, in 1-year-old patient, a case report, 2056.e1–2056.e4
 small, routine intensive care unit admission is necessary, 521–524
- Sublingual abscess, causes acute shortness of breath, *Brucella*-induced, a case report, 1737.e3–1737.e4
- Subtalar dislocation, –fracture, lateral, with maintenance of talonavicular joint, case study, diagnosis and management of, a case report, 2055.e3–2055.e5
- Succinylcholine
 in ED patients with unanticipated hyperkalemia (letter), 1706–1707
 in ED patient with unanticipated hyperkalemia, a case report, 2054.e3–2054.e5
 rocuronium, and hyperkalemia (letter), 1705–1706
- Sugar tong splint, modified, to avoid skin breakdown at the posterior elbow, 332–335
- Suicidal ideation, in ED, demographic characteristics of individuals admitted to hospital for (letter), 1174–1175
- Suicide, risk stratification, and violence screening, on psychiatric emergency service, development and testing of procedures for, 499–504
- Suicide attempts
 and nonsuicidal self-injury, among ED patients older than 50 years, comparison of risk factors and ED visit outcomes, 1016–1021
 presenting to National Trauma Data Bank, 1620–1624
- Superior mesenteric artery dissection, does not necessarily mean acute mesenteric ischemia (letter), 2457
- Superwarfarin
 ingestion treated successfully with prothrombin complex concentrate, a case report, 116.e1–116.e2
 ingestion with PCC, issues with treating (letter), 1168
- Supracondylar nerve block
 ultrasound-guided
 for reduction of distal radius fractures, novel technique in ED, 912–913
 for reduction of distal radius fractures, novel technique in ED (letter), 1172
- Supracondylar radial nerve block, ultrasound-guided, for Colles fractures in ED (letter), 1718–1720
- Supraglottic airway devices, effect of head position on success rate of blind intubation using intubating, 1193–1197
- Supraglottic airway laryngopharyngeal tube, with biohazard suit, blind intubation through (letter), 2040
- Suprasternal notch, as landmark of chest compression depth in CPR, 433–436
- Supraventricular tachyarrhythmia, maternal, in pregnant patients in ED practice, treatment for (letter), 1702–1704
- Supraventricular tachycardia
 novel vagal maneuver technique for termination of, a case report, 118.e5–118.e7
 valsalva maneuver position for treatment of (letter), 2247
- Surfing injury, presenting to US EDs 2002 to 2013, epidemiological study of, 1491–1496
- Surgery
 distal femur fracture, salvage thrombolysis and ECMO for massive pulmonary embolism during, a case report, 1189.e3–1189.e5
 weight loss, and subsequent emergency care use, population-based cohort study, 861–865
- Surgical consult, effect in treatment of abdominal pain in older adults in ED, 1524–1527
- Survey of Health, Ageing and Retirement in Europe Frailty Instrument (SHARE-FI), and adverse outcomes, after ED visit, frailty defined by (letter), 2443–2445
- Swelling, scrotal, and normal lipase, rare presentation of acute pancreatitis, a case report, 763.e5–763.e6
- Swyer-James-Macleod syndrome, in ED, a case report, 1329.e3–1329.e4
- Syncope, and collapse, in acute pulmonary embolism, 1251–1257
- Systemic inflammatory response syndrome, infection, and sepsis, in multiple-injured patients, fresh frozen plasma is permissive for, 1480–1485
- Systemic lupus erythematosus, fever and flank pain with renal vein thrombosis, unusual initial presentation of, a case report, 937.e1–937.e2
- Tachycardia, or rapid ventricular response, three faces of atrial fibrillation (letter), 931–932
- Tackle, causes and costs of ED presentation for American football injuries, 1198–1204
- Taenia asiatica*, role in human cases of cysticercosis (letter), 2030
- Takotsubo broken heart syndrome, my heart is burning, (letter), 1305
- Takotsubo cardiomyopathy
 and Kounis hypersensitivity-associated ACS, myocardial injury through serum troponin I and echocardiography in anaphylaxis (letter), 650–651
 renal impairment and outcome in patients with, 548–552
 warm-up Brugada phenocopy associated with, a case report, 2051.e1–2051.e3
- Takotsubo syndrome
 anaphylaxis-triggered, prevalence of (letter), 748
 diagnosis, in ED, pivotal role of pre-admission electrocardiogram for (letter), 903–904
- Talonavicular dislocation, isolated dorsal, from low-energy mechanism, a case report, 1733.e1–1733.e4
- Tamsulosin, for urolithiasis, 2217–2221
- Targeted temperature management
 prevalence and risk factors for central diabetes insipidus in cardiac arrest survivor treated with, 1400–1405
 successful use in pregnancy after out-of-hospital cardiac arrest, a case report, 122.e3–122.e4
- Target organ damage, and adropin levels, secondary to high blood pressure in ED, 2061–2064
- TASER. *see* Thomas A. Swift Electric Rifle
- Tattoo, posttraumatic, effective prevention using hydrosurgical debridement in ED, a case report, 759.e3–759.e4
- TBSS. *see* Traumatic Bleeding Severity Score (TBSS)
- Terbutaline, and phenylephrine, effect on ischemic priapism, 222–224
- Terminal QRS distortion, present in anterior myocardial infarction but absent in early repolarization, 2182–2185
- Tertiary care hospital
 in Japan, cardiovascular disease outcomes in (letter), 109–111
 urban, usefulness of MISSED score in, 1117–1120
- Testicular torsion, experimental, investigation of N-acetylcysteine on contralateral testis tissue injury by, 1069–1074
- Testis tissue injury, contralateral, by experimental testicular torsion, investigation of N-acetylcysteine on, 1069–1074
- Thermometer, copper-coated, for prevention of cross-infections, preliminary results (letter), 653–656
- Thiol, disulfide, homeostasis, prognostic importance in patients with acute pulmonary thromboembolism, 2315–2319
- Thomas A. Swift Electric Rifle (TASER), penetrating ocular injury, emergent diagnosis and management of, a case report, 1740.e3–1740.e5
- Thoracic outlet obstruction, from muscle hypertrophy, Paget-Schroetter syndrome secondary to, a case report, 1323.e1–1323.e3
- Thoracic surgery, successful treatment of pulmonary embolism with low dose of tissue plasminogen activator after, a case report, 2259.e5–2259.e6
- Thoracostomy
 trocar, new, development and experimental application of, 917–920
 tube, bougie-assisted, placement, novel technique, 101–102

- Thrombectomy**
ultrasound-accelerated rheolytic, for impending paradoxical embolism (letter), 1711
- Thromboembolism, complications, and atrial standstill (letter), 932**
- Thrombolysis**
in acute ischemic stroke, impact of emergency medicine pharmacist on time to, 1997–1999
during continuous chest compression in patient with cardiac arrest due to pulmonary embolism, prolonged CPR–induced spinal cord injury, a case report, 681.e3–681.e5
door to needle, higher prehospital blood pressure prolongs, 1268–1272
of out-of-hospital cardiac arrest patients (letter), 916–917
in pulmonary embolism (letter), 1720–1721
pulseless electrical activity in pulmonary embolism treated with, 1963–1967
salvage, and ECMO, for massive pulmonary embolism during distal femur fracture surgery, a case report, 1189.e3–1189.e5
systemic and catheter, in patients with pulmonary embolism, comparison between, 985–988
usefulness in MI risk index, in acute heart failure, 2351–2355
vs percutaneous catheter intervention, ST elevation in massive pulmonary embolism, a case report, 2466.e3–2466.e4
- Thrombolytic therapy**
in patients with comorbid intracranial aneurysms, acute ischemic stroke is dangerous to treat by, 636–642
successful, for STEMI in patient with ITP, a case report, 345.e1–345.e3
- Thrombosis**
induced by high-dose prothrombin complex concentrate for acute warfarin reversal, a case report, 1736.e1–1736.e3
partial segmental, of corpus cavernosum presenting to ED, a case report, 1182.e3–1182.e5
superior mesenteric artery, after abrupt discontinuation of rivaroxaban, a case report, 764.e5–764.e7
- Thrombotic thrombocytopenic purpura**
after blunt traumatic liver injury, a case report, 939.e3–939.e4
ibuprofen-induced, a case report, 942.e5–942.e7
- Thrombus, formation, left atrial, after brief interruption of rivaroxaban, a case report, 116.e3–116.e4**
- Thyroid testing, utility in ED (letter), 2029–2030**
- Tidal volumes, in adults intubated in ED, body mass index associated with inappropriate (letter), 1682–1683**
- Tissue plasminogen activator, low dose, after thoracic surgery, successful treatment of pulmonary embolism with, a case report, 2259.e5–2259.e6**
- Tono-Pen, vs Icare, in ED (letter), 670–673**
- Tourniquet**
-induced ischemia-reperfusion injury, in rats, postischemic conditioning does not reduce muscle injury after, 2065–2069
use of
performance, prospectively comparing effects of 2 amounts of practice on, 2356–2361
skill decay or maintenance, among first aid caregivers, exploration in manikin model (letter), 1897–1899
- Toxicity**
methamphetamine, acute, use of intravenous lipid emulsion therapy in, a case report, 1732.e3–1732.e4
salicylate, in absence of anion gap metabolic acidosis, a case report, 1328.e1–1328.e3
- Toxic shock syndrome, due to methicillin-resistant Staphylococcus aureus infection after pediatric scald burn, a case report, 1735.e1–1735.e2**
- Tracheal intubation**
during chest compressions in manikin, utility of gum-elastic bougie for, randomized crossover trial, 54–56
child, during CPR, McGrath-MAC vs Macintosh laryngoscopes for (letter), 2453
child, during CPR, videolaryngoscopy or direct laryngoscopy for (letter), 2453
child, during resuscitation by paramedics, comparison of McGrath-MAC and Macintosh laryngoscopes for, 1338–1341
double-lumen tube, in manikin model using VivaSight Double Lumen, randomized controlled comparison with Macintosh laryngoscope (letter), 103–104
face-to-face, comparison of ETView SL, Airtraq, and Macintosh laryngoscopes for, randomized crossover manikin trial (letter), 1893–1894
and laryngoscopy, lip pulling method for prevention of soft tissue injury but also for (letter), 1694
with Macintosh laryngoscope with and without chest compressions, performed by nurses (letter), 2448–2449
with Pentax-AWS Airwayscope, muscle relaxant effect on, prospective randomized controlled trial, 674–675
with VivaSight-SL endotracheal tube by paramedics in cervical immobilized manikin (letter), 309–310
- Tracheostomy**
percutaneous, in post-cervical spine fixation patients, challenging the dogma (letter), 910–911
resuscitating, patient in ED, 1148–1155
- Tranexamic acid**
is used to treat upper gastrointestinal bleeds, in emergency medicine (letter), 1892–1893
treatment of patient in hemorrhagic shock due to leech bite with, a case report, 2253.e3–2253.e4
- Transcatheter arterial embolization, with PCI, acute MI after, potency of, a case report, 1186.e1–1186.e3**
- Transcatheter lumbar arterial embolization, successful, of traumatic hemothorax, a case report, 1188.e3–1188.e5**
- Transcutaneous cardiac pacing, efficacy in ED, 2090–2093**
- Transesophageal echocardiography, for extracorporeal life support vascular cannula placement, emergency physician-performed, 1637–1639**
- Transfusion**
massive
bleeding sites in elderly trauma patients who required, comparison with younger patients, 123–127
modified TBSS, early determination of need for, 1097–1101
-related acute lung injury, after blood products transfusions, risk of (letter), 1674–1675
- Transient synovitis, and septic arthritis, in ED, guideline for differential diagnosis between, 1631–1636**
- Transitional care clinics, for follow-up and primary care linkage for patients discharged from ED, 1230–1235**
- Transplantation, complications, emergency medicine approach to, 2200–2208**
- Transthoracic echocardiography, point-of-care, before CT angiogram of chest, saddle embolism diagnosed by, a case report, 2467.e1–2467.e2**
- Trauma**
within accountable care organization, cost savings associated with transfer of, 455–458
acute MI after, potency of PCI with transcatheter arterial embolization, a case report, 1186.e1–1186.e3
arrests, and hemorrhagic shock, need for abdominal only CPR in treatment of (letter), 1156–1157
arterial base deficit is still useful prognostic marker in, 626–635
blunt
abdominal, 27 years after splenectomy, hemoperitoneum after, a case report, 1325.e3–1325.e4
liver injury, thrombotic thrombocytopenic purpura after, a case report, 939.e3–939.e4
patients, arriving to military level I trauma center from point of injury and interhospital transfers, cumulative effective radiation dose received by, 2397–2401

- progressing to tension pneumopericardium during transportation, simple pneumopericardium due to, a case report, 933.e3–933.e5
- complications, anatomic variations of sternum may mimic, a case report, 1912.e1–1912.e2
- ED, twenty-four-hour packed red blood cell requirement is strongest independent prognostic marker of mortality in, 1121–1124
- in ED, video laryngoscope as new standard of care in (letter), 1313–1314
- elderly patients, who required massive transfusion, bleeding sites in, comparison with younger patients, 123–127
- emergency medical services annotated literature in review 2016, 2193–2199
- end-tidal carbon dioxide and occult injury in patients with, 2146–2149
- examination, young physicians prepared to perform focused assessment with sonography in (letter), 312–314
- extremity
- physician and patient gestalt lead to shared decision to reduce unnecessary radiography in, 349–356
 - physician and patient gestalt lead to shared decision to reduce unnecessary radiography in (erratum), 348
- focused assessment with computed tomography for (letter), 1721–1723
- hemothorax, successful transcatheter lumbar arterial embolization of, a case report, 1188.e3–1188.e5
- intoxicated patient, respiratory depression in, opioids to blame, 250–253
- intubation, fiberoptic intubation or video tube for, method to choose, randomized crossover manikin trial (letter), 751–753
- major, patients, retrospective study of compliance with secondary survey standards in management of (letter), 2446–2448
- minor, with severe rhabdomyolysis, sickle cell trait to blame, a case report, 341.e3–341.e4
- with phencyclidine-positive urine drug screens increase morbidity or mortality, 1066–1068
- posterior shoulder dislocation with large engaging Hill-Sachs lesion, splinting technique, a case report, 682.e1–682.e3
- resuscitation, integration of point-of-care sonography during rapid sequence intubation in (letter), 330
- subarachnoid hemorrhage, complicated by cerebral vasospasm, hemodynamic challenges in (letter), 904–906
- undertriage and overtriage rates, 2191–2192
- Venner A.P. Advance video laryngoscope improve success of first intubation attempt of (letter), 315–316
- wrist, Karadeniz wrist rules for the use of radiography in, 2074–2078
- Traumatic Bleeding Severity Score (TBSS), modified, early determination of need for massive transfusion, 1097–1101
- Triage
- ED
- of cardiology patients, direct cardiologist involvement in (letter), 325–326
 - pain protocol, reduces time to receiving analgesics in patients with painful conditions, 2362–2366
- inhospital, after avalanche-induced cardiac arrest, serum potassium cutoff for (letter), 1317
- sepsis alert and sepsis protocol lower times to fluids and antibiotics in ED, 1–9
- time shock index, modified shock index, and age shock index association with mortality in emergency severity index level 2 patients, 63–68
- Triage Risk Screening Tool (TRST), predictive validity of different versions of (letter), 2454–2456
- Trigeminal nerve foramina, in human cadaveric model, emergency physician sonographers can identify and inject regions of (letter), 312–314
- Trocar, thoracostomy, new, development and experimental application of, 917–920
- Troponin
- cardiac, high-sensitivity, and copeptin, to exclude severe coronary stenosis in patients with chest pain and coronary artery disease, 493–498
 - high-sensitivity, and right ventricular function, in acute pulmonary embolism, 1579–1582
 - leak, coronary anatomy, and prognosis, electrocardiography changes in acute aortic dissection—association with, 1431–1436
 - negative, and acute chest pain, clinical score to obviate the need for cardiac stress testing in patients with, 1421–1426
- Troponin I
- levels, elevated, meaning of, not always ACS, 145–148
 - serum
 - and echocardiography in anaphylaxis, evaluation of myocardial injury through, 140–144
 - and echocardiography in anaphylaxis, myocardial injury through, Takotsubo cardiomyopathy and Kounis hypersensitivity-associated ACS (letter), 650–651
- TRST. *see* Triage Risk Screening Tool (TRST)
- Tuberculosis, advanced disseminated, in patient perceived to be at low risk (letter), 2248
- Tularemia, predicting with clinical, laboratory and demographical findings in ED, 218–221
- Ulinastatin, ameliorates gastrointestinal injury sustained in 2-hit porcine model of septic shock, 1497–1504
- Ulnar nerve block
- ultrasound-guided
 - for boxer fractures, anatomical challenges in (letter), 1903–1904
 - for boxer fractures (letter), 1726–1727
- Ultrasonography
- accelerated rheolytic thrombectomy for impending paradoxical embolism (letter), 1711
 - versus anatomic landmark, approach to cryothyroidotomy (letter), 1696–1697
 - aorta/IVC cross-sectional area index for evaluation of dehydration in children, 1840–1844
- bedside
- accuracy for diagnosis of finger fractures, 809–812
 - for diagnosis of phalanx fractures, accuracy of (letter), 1698
 - emergency, to change diagnosis of peritonsillar abscess using, a case report, 1186.e5–1186.e7
 - for management of shoulder dislocation in ED, 866–870
 - measurement of optic nerve sheath diameter as predictor of intracranial pressure in ED, 963–967
 - point-of-care, effectiveness in diagnosis and management of metacarpal fractures, contribution for management of metacarpal fractures (letter), 674
- B-lines, resolution of, as measure of pulmonary decongestion in acute heart failure, 1129–1132
- cardiac, emergency physician focused, improves diagnosis of ascending aortic dissection, 486–492
- comparison with surface landmarks, in detecting localization for cricothyroidotomy, 254–256
- contrast-enhanced, as imaging technique for patients with acute flank pain into ED (letter), 324–325
- C-reactive protein, and Alvarado score, predictive value in acute appendicitis, 189–192
- and CT scan, in assessment of acute appendicitis, Bayesian comparative model of, 2070–2073
- in ED, bibliometric analysis of top 100 most-cited clinical studies on, 1210–1216
- emergency physicians, can identify and inject regions of trigeminal nerve foramina in human cadaveric model (letter), 312–314
- endobronchial, unusual diagnostic tool for pulmonary embolism, a case report, 684.e1–684.e2
- for evaluation of soft tissue foreign bodies before and after addition of fluid to surrounding interstitial space in cadaveric model, 1779–1782

- first-trimester, high-frequency linear transducer improves detection of intrauterine pregnancy in, 288–291
- guided central venous access, which probe is preferred for viewing subclavian vein using supraclavicular approach?, 1761–1764
 - guided forearm nerve blocks for hand blast injuries, case series and multidisciplinary protocol (letter), 1895–1897
 - guided forearm nerve blocks vs landmark-based wrist blocks for hand anesthesia in healthy volunteers, 730–734
 - guided intravenous education does not increase utilization, alteration of timing of nursing continuing education units for (letter), 2248–2249
 - guided lumbar punctures, vs landmark, perceived difficulty and success rate in novice study group, randomized crossover trial (letter), 2037–2038
 - guided nerve blocks for intracapsular and extracapsular hip fractures, 586–589
 - guided nerve blocks for intracapsular and extracapsular hip fractures (erratum), 1921
 - guided pediatric forearm fracture reductions in resource-limited ED, 40–44
 - guided peripheral intravenous catheter, Delphi method validation of procedural performance checklist for insertion of (letter), 2227–2230
 - guided peripheral vein cannulation (letter), 1675
 - guided posterior tibial nerve blocks, ED, for calcaneal fracture analgesia, a case report, 1183.e1–1183.e3
 - guided short catheter placement, prospective evaluation in internal jugular veins of difficult venous access patients, 578–581
 - guided supracondylar nerve block, for reduction of distal radius fractures, novel technique in ED, 912–913
 - guided supracondylar nerve block, for reduction of distal radius fractures, novel technique in ED (letter), 1172
 - guided supracondylar radial nerve block for Colles fractures in ED (letter), 1718–1720
 - guided ulnar nerve block, for boxer fractures, anatomical challenges in (letter), 1903–1904
 - guided ulnar nerve block for boxer fractures (letter), 1726–1727
 - guided vascular access, effects of long axis in-plane vs short axis out-of-plane techniques during, 778–783
- hand dominance in intravenous drug using patients does not affect peripheral venous access sites identified by, 1088–1091
- IVC, NT-proBNP, and bioelectrical impedance analysis, for diagnosis of acute decompensated CHF, 1817–1822
- lung
- diagnostic and prognostic tool at every step in pathway of care for acute heart failure, 656–657
 - in differentiating between pulmonary infection and pulmonary infection with acute left ventricular heart failure, clinical application of rapid B-line score with, 278–281
- measurement of IVC collapse predicts propofol-induced hypotension, 1125–1128
- neuro, and mental foramen, in ED (letter), 2240–2241
- noninvasive, corrected flow time, measure to detect preload reduction by nitroglycerin, 1859–1862
- for nurse-performed intravenous placement in difficult access ED patients, randomized controlled trial assessing use of, 1950–1954
- optic nerve sheath diameter, knowledge, attitude, and practices of paramedics regarding, 1160–1161
- paramedics are able to confirm endotracheal tube placement using (letter), 923–924
- point-of-care
- impact on ED time to disposition for patients with nontraumatic shock, 1022–1030
 - bedside, and computed tomography in elbow injuries, comparison of, 2186–2190
 - in detection of pediatric esophageal food impaction, a case report, 763.e1–763.e3
 - as diagnostic tool for patella fractures, accuracy of, 1576–1578
 - at ED maximizes patient confidence in emergency physicians, 657–659
 - fellowship application process, innovation in (letter), 1303–1305
 - in pediatric small bowel obstruction, ED case series, a case report, 2464.e1–2464.e2
 - value for detecting nail bed injury in ED, 1850–1854
- role in diagnosis of metacarpal fractures, 1868–1871
- still lung point, new evidence for pneumomediastinum, a case report, 344.e1–344.e2
- for supraclavicular central venous catheterization, influence of caudal traction of ipsilateral arm on, 851–855
- as a tool for prehospital recognition of tension pneumothorax (letter), 1302–1303
- tracheal, and lung sliding, for confirming endotracheal tube placement, speed and reliability, 953–956
- trans-abdominal, in simulated massive acute overdose, accuracy of, 1455–1457
- in trauma examination, young physicians prepared to perform focused assessment with (letter), 314–315
- use in ocular diseases (letter), 112
- vs anatomic landmark, to locate cricothyroid membrane, comparison of (letter), 1696
- uPAR. *see* Urokinase-type plasminogen activator receptor (uPAR)
- Upper teeth, by video laryngoscopy, investigation of force received at (letter), 1695
- Urea cycle disorder
- isolated hyperammonemia due to, rare cause of postpartum coma (letter), 1894–1895
 - rare cause of postpartum coma, isolated hyperammonemia due to, a case report, 1324.e3–1324.e4
- Ureteric calculi, synchronous, bilateral, anuria and acute kidney injury an uncommon case of, a case report, 2256.e1–2256.e2
- Urinalysis, automated, and routine tests, in patients with suspected urinary tract infection at ED, 1528–1534
- Urinary obstruction, is important complicating factor in patients with septic shock due to urinary infection, 694–696
- Urinary retention, acute, STEMI, unusual adverse event of, case report, 347.e1–347.e2
- Urinary tract infections
- ED pharmacist monitoring of provider antibiotic selection aids appropriate treatment for outpatient, 1600–1603
 - suspected, at ED, routine tests and automated urinalysis in patients with, 1528–1534
 - urinary obstruction is important complicating factor in patients with septic shock due to, 694–696
- Urine drug screen, phencyclidine-positive, increase morbidity or mortality, trauma patients with, 1066–1068
- Urokinase-type plasminogen activator receptor (uPAR), soluble, clinical value in diagnosis, prognosis, and therapeutic guidance of sepsis, 375–380
- Urolithiasis, tamsulosin for, 2217–2221
- Uterine rupture, spontaneous, due to placenta percreta during pregnancy, a case report, 1918.e1–1918.e3
- UTI. *see* Urinary tract infections
- Vagal maneuver, technique, novel, for termination of supraventricular tachycardias, a case report, 118.e5–118.e7
- Valsalva maneuver, position for treatment of supraventricular tachycardia (letter), 2247
- Vapocoolant spray, vs placebo spray, in adults undergoing venipuncture, prospective, randomized, double-blind controlled trial comparing, 798–804
- Varicella-zoster virus encephalitis, in immunocompetent patient without rash, a case report, 2257.e1–2257.e2

- Vascular access
 during CPR, intraosseous access devices as method of (letter), 321–322
 technique, chosen during hypovolemic shock (letter), 1886–1887
 ultrasound-guided, effects of long axis in-plane vs short axis out-of-plane techniques during, 778–783
- Vascular cannula, placement of, extracorporeal life support, emergency physician-performed transesophageal echocardiography for, 1637–1639
- Vasculitis
 ANCA-positive, with diffuse alveolar hemorrhage related to spider bite, a case report, 339.e3–339.e4
 levamisole-induced, in an illicit drug user, a case report, 1321.e5–1321.e6
- Vasopressin, on kinetics and survivability in cardiac arrest, effects of tibial intraosseous and intravenous administration of, 429–432
- Venipuncture, prospective, randomized, double-blind controlled trial comparing vapocoolant spray vs placebo spray in adults, 798–804
- Venner A.P. Advance, video laryngoscope, improve success of first intubation attempt of trauma patient (letter), 315–316
- Venous catheterization, during chest compressions, needle guides for, crossover simulation trial, 989–992
- Venous thromboembolism, prophylaxis in ED observation units (letter), 2238–2240
- Ventilation
 in ED, attention must be paid (letter), 1901
 pediatric, with 3 different airway modalities, use of capnographs to assess quality of, 69–74
 single rescuer, using bag-valve mask with internal handle, randomized crossover trial, 1991–1996
- Ventricular assist devices, emergency management of, 1294–1301
- Ventricular fibrillation
 benefits of respective and combined use of green tea polyphenols and ERK inhibitor on the survival and neurologic outcomes in cardiac arrest rats induced by, 570–575
 combination of chest compressions and interposed abdominal compressions in swine model of, 968–974
 effects of proximal and distal routes of intraosseous epinephrine administration on short-term resuscitative outcome measures in adult swine model of, randomized controlled study, 49–53
 firefighters able to recognize (letter), 1885–1886
 model of rats, edaravone improves survival and neurological outcomes after CPR in, 1944–1949
 self-termination during transport by EMS, a case report, 940.e1–940.e3
- Ventricular rate control, achieving, using metoprolol in beta-blocker-naïve patients vs patients on chronic beta-blocker therapy, 606–608
- Ventricular tachycardia, cardiac arrest, during central line placement, a case report, 114.e3–114.e4
- Video laryngoscope
 as new standard of care in trauma ED (letter), 1313–1314
 Venner A.P. Advance, improve success of first intubation attempt of trauma patient (letter), 315–316
- Videolaryngoscopy
 comparison with Macintosh direct laryngoscopy, for airway management, 1886
 or direct laryngoscopy for child tracheal intubation during CPR (letter), 2453
- Video tube, or fiberoptic intubation, for trauma patient intubation, method to choose, randomized crossover manikin trial (letter), 751–753
- Violence, screening, and suicide risk stratification on psychiatric emergency service, development and testing of procedures for, 499–504
- Visceral artery dissection, spontaneous, in ED, characteristics and diagnostic pitfalls of, 1092–1096
- Visual recognition, of anatomical structures in circulated and in non-circulated airway, 1236–1240
- Vital signs, prehospital, chart sharing system, development of, 88–92
- VivaSight, single lumen endotracheal tube, comparison with Macintosh laryngoscope for emergency intubation by experienced paramedics in standardized airway manikin with restricted access, randomized, crossover trial (letter), 929–930
- VivaSight DLT, Double-lumen tube tracheal intubation in manikin model using, randomized controlled comparison with Macintosh laryngoscope (letter), 103–104
- VivaSight-SL, endotracheal tube, tracheal intubation with, by paramedics in cervical immobilized manikin (letter), 309–310
- Vortex tube, rapid and selective brain cooling method using, 887–894
- Wait time, ED, relationship between racial disparities and illness severity in, 10–15
- Wallenberg's lateral medullary syndrome, in adolescent, a case report, 2254.e1–2254.e2
- Warfarin
 patients with minor head injury and normal initial CT scan, outcomes of, 75–78
 reversal, acute, thrombosis induced by high-dose prothrombin complex concentrate for, a case report, 1736.e1–1736.e3
- Weather, conditions, and air pollution, effects on incidence of acute MI, 449–454
- WebPOISONCONTROL, poison control can be automate using, 1614–1619
- Weight loss
 clenbuterol causing NSTEMI in teenage female desiring to, a case report, 1739.e5–1739.e7
 surgery, and subsequent emergency care use, population-based cohort study, 861–865
- Wellens syndrome
 late diagnosis in patient presenting with atypical ACS, a case report, 338.e3–338.e5
 in 22-year-old man, a case report, 937.e3–937.e4
- WhatsApp application, use for orthopedic consultations in ED (letter), 1305–1307
- Wrist block, landmark-based, vs forearm ultrasound-guided nerve blocks, for hand anesthesia in healthy volunteers, 730–734
- Wrist drop, modified coaptation splint with sugar tong intrinsic plus extension for initial management of, 659–663
- Wrist fracture, postsplinting x-rays of, 1625–1626
- Wrist positioning, optimal, for radial arterial cannulation in adults, 2372–2378
- Wrist trauma, Karadeniz wrist rules for the use of radiography in, 2074–2078